



2019, Volume 13, Issue 1 (Volume 13, Issue 1, January-July 2019)

Effect of crude extracts of the peels of *Musa acuminata* L. Banana plant in some biological aspects of *Culex Molestus* Forskal (Diptera: Culicidae) with an estimation of the enzymatic effectiveness of Tyrosinase

Sarah Kadhim Al-Rahimy, Aseel Kariem Al-Sultany, Zahraa Raheem Murshidy, Rafid Abbas Al-Essa, Ayam Razaq Abdul Kadhim

[Download Full Article - PDF format](#)

[Abstract](#)

Physico-chemical parameters, chlorophyll a and phytoplankton community as trophic state indices of two tropical lakes, southwestern Nigeria

Adedolapo Abeke Ayoadé, Blessing O. Osuala, Temitayo A. Adedapo

[Download Full Article - PDF format](#)

[Abstract](#)

Cannibalism of the medical leeches *Hirudo verbana*

Ruslan Aminov

[Download Full Article - PDF format](#)

[Abstract](#)

Hydrothermal assisted biological synthesis of Silver nanoparticles by using honey and gomutra (Cow Urine) for qualitative determination of its antibacterial efficacy against *Pseudomonas sp.* isolated from contact lenses

Nishita Jain, Preeti Bhosale, Vidya Tale, Rabinder Henry, Jayant Pawar

[Download Full Article - PDF format](#)

[Abstract](#)

Biochemical study of amino acid profile of *Kappaphycus alvarezii* and *Gracilaria salicornia* seaweeds from Gerupuk Waters, West Nusa Tenggara (NTB)

S. Y. Lumbessy, S. Andayani, H. Nursyam, M. Firdaus

[Download Full Article - PDF format](#)

[Abstract](#)

Determination of congenital cytomegalovirus infection and virus genotypes in newborn urine samples

Tariq M. Quasim, Abdel Wahab A R. Alshaikhly

[Download Full Article - PDF format](#)

[Abstract](#)

Operator model to control process of obtaining vermicompost

Evgeny V. Kuznetsov, Anna E. Khadzhidi, Yan A. Poltorak, Margarita Kuznetsova

[Download Full Article - PDF format](#)

[Abstract](#)

Eco-friendly flame retardant additives for epoxy resin nanocomposites

Nguyen Tuan Anh, Nguyen Quang Tung, Nguyen Xuan Canh, Nguyen Van Hoan

[Download Full Article - PDF format](#)

[Abstract](#)

Development new multiplex reverse transcription quantitative real time polymerase reaction to detect human norovirus genogroups I and II

Nadira Salman Mohamed, Shaymaa Naji Dahham, Mohammed Nadhir Maarof

[Download Full Article - PDF format](#)

[Abstract](#)

Justification parameters of the ripper tooth of combined unit for minimum tillage

B. Kalimbetov, B. Kalymbetov, G. Kenzhibayeva, D. Karmanov, A. Tukhtakuziyev, B. Kablan

[Download Full Article - PDF format](#)

[Abstract](#)

Compilation and revision of the laws and regulations of protected areas in accordance with international conventions and obligations

Atefeh Mahdi, Dariush Karimi, Parvin Farshchi, Mostafa Panahi

[Download Full Article - PDF format](#)

[Abstract](#)



Biochemical study of amino acid profile of *Kappaphycus alvarezii* and *Gracilaria salicornia* seaweeds from Gerupuk Waters, West Nusa Tenggara (NTB)

S.Y. Lumbessy^{1,2*}, S. Andayani³, H. Nursyam³, M. Firdaus³

¹ Doctoral Program, Faculty of Fisheries and Marine Science, Brawijaya University, INDONESIA

² Study Program of Aquaculture, University of Mataram, West Nusa Tenggara, INDONESIA

³ Faculty of Fisheries and Marine Science, Brawijaya University, INDONESIA

*Corresponding author: alyachali@gmail.com

Abstract

This study aimed to analyze the amino acid profile of *Kappaphycus alvarezii* and *Gracilaria salicornia* seaweeds from Gerupuk Waters, West Nusa Tenggara (NTB). Samples were collected in January using a transect method, then dried and ground to powder to determine the amino acid content. Amino acid analysis was performed using high-performance liquid chromatography (HPLC). An amino acid standard curve was used to calculate amino acid content of each sample. The results of the analysis showed that there were 15 amino acids present in *K. alvarezii* seaweed, consisting of seven essential amino acids and eight non-essential amino acids. Meanwhile, *G. salicornia* seaweed contained 13 amino acids, consisting of seven essential amino acids and six non-essential amino acids. In *K. alvarezii*, the amino acid with the highest level was glutamate (13.73%), while in *G. salicornia*, the amino acid with the highest level was serine (13.02%). The results of the nutritional evaluation indicate that lysine is a limiting amino acid in *K. alvarezii* seaweed and methionine is a limiting amino acid in *G. salicornia* seaweed.

Keywords: amino acid, *Gracilaria salicornia*, Gerupuk Waters, *Kappaphycus alvarezii*, seaweed

Lumbessy SY, Andayani S, Nursyam H, Firdaus M (2019) Biochemical study of amino acid profile of *Kappaphycus alvarezii* and *Gracilaria salicornia* seaweeds from Gerupuk Waters, West Nusa Tenggara (NTB). Eurasia J Biosci 13: 303-307.

© 2019 Lumbessy et al.

This is an open-access article distributed under the terms of the Creative Commons Attribution License.

INTRODUCTION

Seaweed is an important marine biological resource with high nutritional value, and thus it becomes one of the most promising plants of the future. With its unique biochemical structure and composition, seaweed is widely used in various industries, such as those engaged in the foods, energy, medicine and cosmetic sectors, as well as utilized for biotechnology (Komalavalli and Lalitha 2015, Norziah and Ching 2000, Ommee and Payap 2012). Most seaweed contains essential amino acids which are almost identical to the essential amino acids found in eggs (Norziah and Ching 2000). In addition, most seaweed contains quite a lot of aspartic acid, glutamic acid, and arginine in the composition of total amino acids (Rajasulochana *et al.* 2010). The varied nutritional composition of seaweed is affected by the species, habitat, and environmental conditions (Zawawi *et al.* 2014).

Kappaphycus alvarezii and *Gracilaria salicornia* are two species of seaweed that are in the class of Rhodophyceae, largely found in Gerupuk Waters, West Nusa Tenggara (NTB). *K. alvarezii* is a species of seaweed with a high economic value that has been widely cultivated as a source of carrageenan

(Madhavarani and Ramanibai 2014). Meanwhile, *G. salicornia* has the potential to produce jelly which, due to its high chlorophyll content, has potential as antioxidant properties (Vijayavel and Martinez 2010, Vimala and Poonghuzhali 2015).

The results of Xiren and Aminah's (2017) study suggest that *K. alvarezii* originating from Langkawi and Sabah Waters, Malaysia contains 19 amino acids, with aspartic acid, glutamic acid, alanine, and leucine being the dominant amino acids. On another side, the information on the amino acid content of *Gracilaria sp.* genus suggests that the aspartic acid and glutamic acid of this species of seaweed are present in a dominant number (Fateminasab 2014, Kumar and Kaladharan 2007, Norziah and Ching 2000). This research provides limited information on the sea surrounding Malaysia. In fact, *K. alvarezii* and *G. salicornia* are widely spread in several marine areas in Southeast Asia, including Indonesia. Biochemical composition of seaweed is reported to be generally affected by geographical locations and conditions of the surrounding environment

Received: August 2018

Accepted: January 2019

Printed: May 2019

Table 1. Amino acid composition of *K. alvarezii* and *G. salicornia* seaweeds in Gerupuk Waters, NTB

No.	Amino acids	<i>K. alvarezii</i>		<i>G. salicornia</i>	
		mg/g	(% b/b)	mg/g	(% b/b)
<i>Essential amino acids</i>					
1	Histidine	16.01 ± 0.36	1.01	22.92 ± 0.67	7.92
2	Threonine	24.90 ± 0.57	9.9	22.51 ± 0.63	7.51
3	Methionine	16.86 ± 0.67	1.86	16.11 ± 0.67	1.11
4	Valine	24.94 ± 0.59	9.94	22 ± 0.59	7
5	Phenylalanine	21.05 ± 0.15	6.05	17.94 ± 0.25	2.94
6	Isoleucine	21.37 ± 0.30	6.37	19.85 ± 0.42	4.85
7	Leucine	23.43 ± 0.04	8.43	21.60 ± 0.56	6.6
	<i>Total EAA</i>	148.55 ± 3.62	43.56	142.93 ± 2.57	37.93
	<i>% of total AA</i>	33.64	47.58	49.58	46.49
<i>Non-essential amino acids</i>					
8	Aspartic acid	33.26 ± 0.09	12.26	ND	ND
9	Glutamic acid	116.73 ± 0.16	13.73	27.93 ± 0.11	11.93
10	Serine	26.82 ± 0.27	4.82	29.02 ± 0.12	13.02
11	Glycine	29.71 ± 0.31	7.71	21.85 ± 0.58	5.85
12	Arginine	24.04 ± 0.26	2.04	24.04 ± 0.30	2.34
13	Alanine	29.33 ± 0.17	7.33	25.11 ± 0.85	9.11
14	Tyrosine	18.06 ± 0.10	0.06	17.40 ± 0.21	1.4
15	Lysine	15.05 ± 0.01	0.05	ND	ND
	<i>Total non-EAA</i>	293 ± 32.94	48	145.53 ± 11.77	43.65
	<i>% of total AA</i>	66.36	52.42	50.42	53.51
	<i>Total AA</i>	441.55	91.56	288.28	81.58
	<i>EAA/non-EAA</i>	0.51	0.91	0.98	0.87
	<i>EAA/total AA</i>	0.34	0.48	0.50	0.46

*AA: amino acid; EAA: essential amino acid; Non-EAA: non-essential amino acid; ND: not determined

(Boyaci et al. 2014, Davoobadi and Aghajani 2013, Rohani 2012).

There is an abundance of *K. alvarezii* and *G. salicornia* in Gerupuk, NTB, but the utilization of this seaweed is limited due to limited knowledge of its nutritional value and limited human resources. Therefore, this research was intended to analyze the amino acid profile of *K. alvarezii* and *G. salicornia* seaweeds from Gerupuk Waters, West Nusa Tenggara (NTB), in order to provide information on the nutritional value of both species.

MATERIALS AND METHODS

Sampling Methods

Seaweed samples used in this research were *K. alvarezii*, obtained from Gerupuk Waters, NTB, at the ordinates of 8°55'3,46"S Lat. and 116°20'3,49" E Lon. and *G. Salicornia*, collected from the same waters at the ordinates of 8°55'1,4" S Lat. and 116°19'57,22 E Lon. Sampling of both seaweeds was conducted during the month of January. The collected sample was then identified in the Indonesian Science Center (LIPI), North Jakarta.

Samples of both seaweed species were collected in a fresh condition. Samples were placed inside plastic polybags, stored on ice, and placed in a cool box. The seaweed samples were immediately taken to the laboratory. In the laboratory, the seaweed samples were washed with clean water to remove dirt. After washing, samples were wiped with tissue paper to remove any excess water. The seaweed samples were then dried in an oven at 37 °C until reaching a constant weight. The dried samples were mashed using a blender and sieved to obtain the sample powder which was then stored in a refrigerator (Lima-Filho et al. 2002).

Analysis of Amino Acids

The amino acid analysis was performed using high-performance liquid chromatography (HPLC) schimadzu LC 10 A (column lichrospher 100 Rp-18 (sum). The eluent used was 50 mM Natrium asetat and sodium dibasic phospat : THF : Methanol (96 : 2 : 2), pH 6.8.

One gram each sample was put into the reaction tube, 4 ml of HCl 6 N was then added and the mixture refluxed for 24 hours at 110 °C. After that, the samples were cooled and neutralized with 6 N NaOH to pH 7, then filtered using 0.2 µm Whatman filter paper. As much as 10 µL of each sample was taken and 300 µL of OPA was added. The samples were then left for 1 minute for perfect derivation. After that, 20 µL of the sample was injected into the HPLC column for separation of the different amino acids. The standard amino acid solution used in HPLC analysis consisted of 20 types of standard amino acid. HPLC analysis performed twice for each sample. The amount of each amino acid contained in both seaweed samples was determined using a standard amino acid curve. Meanwhile, the amino acid scores were determined using the following formula:

$$\text{Amino Acid Score of the Sample} = \frac{\text{Essential Amino Acid Content of the Sample}}{\text{FAO / WHO Standard Amino Acid Pattern}}$$

Data Analysis

The amino acid data were analyzed descriptively and presented in the form of narration and table.

RESULTS

The results of the amino acid analysis of *K. alvarezii* and *G. salicornia* seaweeds in Gerupuk Waters, NTB, can be seen in **Table 1** and **Fig. 1**. Based on the results of the amino acid analysis using HPLC (**Table 1**), there were 15 amino acids in *K. alvarezii* seaweed, consisting




Fig. 1. Chromatograms of amino acids in seaweeds, (A) *G. Salicornia* and (B) *K. Alvarezii*

Table 2. Essential amino acids (mg/g protein) in *K. alvarezii* and *G. salicornia* compared to the amino acid reference pattern recommended by FAO (1991)

Essential amino acids	<i>K. alvarezii</i>	<i>G. salicornia</i>	FAO/WHO (1991)
Isoleucine	22.75	17.32	28
Leucine	12.77	10	66
Lysine	0.08	ND*	58
Phenylalanine + Tyrosine	9.70	6.89	63
Methionine	7.44	4.44	25
Threonine	29.11	22.09	34
Valine	28.4	20.00	35

*ND, not determined

of seven essential amino acids and eight non-essential amino acids. Meanwhile, in *G. salicornia* seaweed, 13 amino acids were identified, consisting of seven essential amino acids and six non-essential amino acids. Furthermore, the percentages of the total essential amino acids contained in *K. alvarezii* and *G. salicornia* seaweeds were 47.58% and 46.49% of the total amino acids. Meanwhile, the percentages of the total non-essential amino acids contained in *K. alvarezii* and *G. salicornia* seaweeds were 52.42% and 53.51%, respectively.

After the amino acid composition of *K. alvarezii* and *G. salicornia* (Table 1) were determined, the next step was to evaluate the nutritional value by comparing the essential amino acid content of the samples with the amino acid reference pattern recommended by Food Agriculture Organization of United Nations (FAO) in 1991. This is particularly helpful in initial estimation of the nutrient content of seaweed protein (Kumar and Kaladharan 2007, Siddique et al. 2013, Xiren and Aminah 2017).

Amino acid score data presented in Table 2 show that lysine is a limiting amino acid in *K. alvarezii* seaweed and methionine is a limiting amino acid in *G. salicornia* seaweed due to its lowest chemical value. Meanwhile, threonine is the amino acid with the highest chemical score in both *K. alvarezii* and *G. salicornia* seaweeds.

DISCUSSION

The samples of *K. alvarezii* and *G. salicornia* seaweed had the same essential amino acid profile, which included histidine, leucine, threonine, valine, methionine, isoleucine, and phenylalanine. However, both seaweed samples differed in their non-essential amino acid profile; aspartate and lysine were present only in *K. alvarezii*, but did not exist in *G. salicornia*. According to previous studies on the amino acid profile of *K. alvarezii* seaweed originating from Brazil (Gressler et al. 2010) and Malaysia (Matanjan et al. 2009), 16 amino acids were identified, consisting of nine essential amino acids and seven non-essential amino acids. Gressler et al. (2010) stated that the profile of amino acids can determine the nutritional quality of food, especially in the evaluation of new protein sources. Amino acid content is linearly dependent on seaweed protein content in which low protein content causes low amino acid content. Therefore, the factors affecting amino acid content are similar to those affecting protein content, involving species, environment, and harvested period (Hardjani et al. 2017).

The result of essential amino acid percentages in *K. alvarezii* and *G. salicornia* in Table 1 were consistent with findings of Dawczynski et al. (2007), namely that the essential amino acid content of some types of seaweed was 30% of the total amino acids. Several previous studies have reported that the total essential amino acids of *Caulerpa lentillifera* and *Ulva Reticulata* accounted for nearly 40% of the total amino acids (Ratana-arporn and Chirapart, 2006), those of *Kappaphycus alvarezii* dan *Hypnea musciformis* reached 45–49% (Dawczynski et al. 2007, Kumar and Kaladharan 2007), and those of *Gelidium pusillum* reach 52.08% (Siddique et al. 2013).

The results in the Table 1 also showed that glutamic acid (13.73%) and aspartic acid (12.26%) contained in *K. alvarezii* seaweed were the amino acids that occurred at the highest level. On the other hand, the amino acids in *G. salicornia* seaweed that occurred at the highest

level were glutamic acid (11.93%) and serine (13.02%). Several previous studies have also shown that aspartic acid and glutamic acid content of red seaweed ranged from 15% to 33% (Gressler *et al.* 2009, Matanjun *et al.* 2009, Ortiz *et al.* 2009). The content of aspartic acid and glutamic acid in red and green seaweeds were significantly higher than that of brown seaweed. The high content of aspartic acid and glutamic acid contributes to providing a distinctive flavor in seaweed; its savory flavor (*umami* flavor), for example, leads to its use as food seasoning to stimulate appetite in elderly people. Meanwhile, serine can provide a sweet taste that results in the complex flavor of seaweed (Yaich *et al.* 2011, Helaly 2013). Furthermore, Holds and Kran (2011) reported that red algae contain many glutamic acids. The taste of *nori* (dried or edible seaweed often used in Japanese cookery) is the result of a large number of different amino acids, including alanine, glutamic acid, and glycine.

The high percentage of glutamic acid, aspartic acid, and serine in *K. alvarezii* and *G. salicornia* seaweeds can be used as the consideration in the utilization of other potentials of both seaweeds, especially in the health sector. Khanifar *et al.* (2011) mentioned that glutamic acid is a precursor of glutamine and GABA. Aspartic acid helps to protect the liver and is involved in the metabolism of DNA and RNA. Moreover, it plays a role in the immune system by increasing the production of immunoglobulins and the formation of antibodies. Meanwhile, serine is essential in maintaining blood

sugar levels, plays a role in the central nervous system including the development of the myelin sheath, and is necessary for muscle growth and maintenance.

The nutritional evaluation in **Table 2** indicate that lysine is a limiting amino acid in *K. alvarezii* seaweed and methionine is a limiting amino acid in *G. salicornia* seaweed. Meanwhile, threonine is the highest amino acids in both *K. alvarezii* and *G. salicornia* seaweeds. This result is proportional with the result of the study by Kumar and Kaladharan (2017). They suggest that leucine and lysine are limiting amino acids in *K. alvarezii*, while threonine and tryptophane had the highest chemical score. In contrast, lysine and methionine are limiting amino acids in *Gracilaria sp.*, while threonine and tryptophane had the highest score. Furthermore, the study of Xiren and Aminah (2017) showed that lysine is a limiting amino acid in *K. alvarezii* seaweed species from both Langkawi and Sabah Waters in Malaysia.

In general, it can be seen that the content of *K. alvarezii* and *G. salicornia* seaweeds is rich in aromatic amino acids (i.e. threonine) and has limited sulfur amino acids (i.e. lysine). This suggests that *K. alvarezii* and *G. salicornia* seaweeds can be a source of complementary protein for humans and animals.

ACKNOWLEDGEMENT

The author thank the DP2M DIKTI for funding this research through Doctoral Dissertaion Research Funding Program.

REFERENCES

- Alwarsamy M, Ravichandran R (2014) In-Vitro antibacterial Activity of Kappaphycus alvarezii extracts collected from Mandapam Coast, Rameswaram, Tamil Nadu. International Journal of Innovative Research in Science, Engineering and Technology, 3(1): 8436-40.
- Benjama O, Masniyom P (2012) Biochemical composition and physicochemical properties of two red seaweeds (*Gracilaria fisheri* and *G. tenuistipitata*) from the Pattani Bay in Southern Thailand, Songklanakarin. Journal of Science and Technology, 34(2): 223-30.
- Boyacı A, Tutoğlu A, Boyacı N, Koca İ, Dokumacı DŞ, Kocatürk Ö, ... Şakalar A (2014) An evaluation of radial and ulnar artery flow characteristics in diabetic patients with carpal tunnel syndrome and the diagnostic value of ultrasonography in these patients. J Clin Exp Invest, 5(2). <https://doi.org/10.5799/ahinjs.01.2014.02.0386>
- Davooabadi FM, Aghajani H (2013) Identification of Potential Groundwater Zones Using RS and GIS. UCT Journal of Research in Science, Engineering and Technology, 1(1): 4-6.
- Dawczynski C, Schubert R, Jahreis G (2007) Amino Acid, fatty acid, and dietary fibre in edible seaweed products. Food chemistry, 103: 891–9. <https://doi.org/10.1016/j.foodchem.2006.09.041>
- FAO (1991) Protein quality evaluation: Report of joint FAO/WHO expert consultation. Rome, Italy: 1-66.
- Fateminasab A (2014) Investigating the challenges and barriers of convergence between Iran and republic of Azerbaijan. UCT Journal of Social Sciences and Humanities Research, 1: 18-24.
- Gressler V, Yokoya NS, Fujii MT, Colepicolo P, Filho JM, Torres RP, Pinto E (2010) Lipid, fatty acid, protein, amino acid and ash contents in four Brazilian red algae species. Food Chemistry, 120(2): 585-90. <https://doi.org/10.1016/j.foodchem.2009.10.028>
- Hardjani DK, Suantika G, Aditiawati P (2017) Nutritional Profile of Red Seaweed *Kappaphycus alvarezii* after Fermentation using *Saccharomyces cerevisiae* as a Feed Supplement for White Shrimp *Litopenaeus vannamei*

- Nutritional Profile of Fermented Red Seaweed. *Journal of Pure and Applied Microbiology*, 11(4): 1637-45. <https://doi.org/10.22207/JPAM.11.4.01>
- Helaly MAH, Hatata ESZ, Abu-Elmagd M, Ibrahim EF, Alsaied A, El-Aal A, ... Settin A (2013) Association of IL-10 and IL-6 Gene Polymorphisms with Type 2 Diabetes Mellitus among Egyptian Patients. *European Journal of General Medicine*, 10(3): 158-62. <https://doi.org/10.29333/ejgm/82250>
- Holdt SL, Kraan S (2011) Bioactive compounds in seaweed: functional food applications and legislation. *Journal of Applied Phycology*, 23(3): 543-97. <https://doi.org/10.1007/s10811-010-9632-5>
- Khanifar J, Ghoorchian H, Ahmadi AR, Hajihosaini R (2011) Comparison of essential and non Essential amino acids in the single cell Protein (scp) Of White Rot Fungi From Wheat Straw. *African Journal of Agricultural Research*, 6(17): 3994-9.
- Komalavalli N, Lalitha N (2015) Proximate Composition and Amino Acid Profile of Five Green Algal Seaweeds from Mandapam Coastal Regions, Tamil Nadu, India. *International Journal of Advances in Interdisciplinary Research*, 2(2): 37-40.
- Kumar V, Kaladharan P (2007) Amino acids in the seaweeds as an alternate source of protein for animal feed. *Journal of the Marine Biological Association of India*, 49(1): 35-40.
- Lima-Filho JVM, Carvalho AFFU, Freitas SM, Melo VMM (2002) Antibacterial Activity of Extracts of Six Macro Algae from the Northeastern Brazilian Coast. *Brazilian Journal of Microbiology*, 33(4): 311-3. <https://doi.org/10.1590/S1517-83822002000400006>
- Matanjan P, Mohamed S, Mustapha NM, Muhammad K (2009) Nutrient content of tropical edible seaweeds, *Euclima cottonii*, *Caulerpa lentillifera* and *Sargassum polycystum*. *Journal of Applied Phycology*, 21(1): 75-80. <https://doi.org/10.1007/s10811-008-9326-4>
- Norziah MH, Ching CY (2000) Nutritional composition of edible seaweed *Gracilaria changgi*. *Journal of Food Chemistry*, 68(1): 69-76. [https://doi.org/10.1016/S0308-8146\(99\)00161-2](https://doi.org/10.1016/S0308-8146(99)00161-2)
- Ortiz J, Romero N, Robert P, Araya J, Hernandez JL, Bozzo C, Navarrete E, Osario A, Rios A (2006) Dietary fiber, amino acid, fatty acid and tocopherol contents of the edible seaweeds *Ulva lactuca* and *Durvillaea antarctica*. *Food chemistry*, 99(1): 98-104. <https://doi.org/10.1016/j.foodchem.2005.07.027>
- Rajasulochana P, Krishnamoorthy P, Dhamotharan R (2010) Amino acids, Fatty acids and minerals in *Kappaphycus* sps. *Journal of Agricultural and Biological Science*, 5(5): 1990-6145.
- Ratana-arporn P, Chirapart A (2006) Nutritional Evaluation of Tropical Green Seaweeds *Caulerpa lentillifera* and *Ulva reticulata*. *Journal Natural Science* 40: 75-83.
- Rohani GK, Abdulalian E, Ng WK (2012) Evaluation of the proximate, fatty acid and mineral composition of representative green, brown and red seaweeds from the Persian Gulf of Iran as potential food and feed resources. *Journal of Food Science and Technology*, 49(6): 774-80. <https://doi.org/10.1007/s13197-010-0220-0>
- Siddique MAM, Khan MSK, Bhuiyan MKA (2013) Nutritional composition and amino acid profile of a sub-tropical red seaweed *Gelidium pusillum* collected from St. Martin's Island, Bangladesh. *International Food Research Journal*, 20(5): 2287-92. <https://doi.org/10.3153/jfscm.2013018>
- Vijayavel K, Martinez JA (2010) In vitro antioxidant and antimicrobial activities of two Hawaiian marine Limu: *Ulva fasciata* (Chlorophyta) and *Gracilaria salicornia* (Rhodophyta). *Journal of Medicinal Food*, 13(6): 1494-9. <https://doi.org/10.1089/jmf.2009.0287>
- Vimala T, Poonghuzhali TV (2015) Estimation of Pigments from Seaweeds by Using Acetone and DMSO. *International Journal of Science and Research (IJSR)*, 4(10): 1850-4.
- Xiren GK, Aminah A (2017) Proximate composition and total amino acid composition of *Kappaphycus alvarezii* found in the waters of Langkawi and Sabah, Malaysia. *International Food Research Journal*, 24(3): 1255-60.
- Yaich H, Garna H, Besbes S, Paquot M, Blecker C, Attia H (2011) Chemical composition and functional properties of *Ulva lactuca* Seaweed Collected in Tunisia. *Food Chemistry*, 128(4): 895-901. <https://doi.org/10.1016/j.foodchem.2011.03.114>
- Zawawi MH, Idris MH, Kamal AHM, Wong SK (2014) Seaweed composition from Bintulu Coast of Sarawak, Malaysia. *Pakistan Journal of Biological Sciences*, 17(8): 1007-14. <https://doi.org/10.3923/pjbs.2014.1007.1014>