



Digital Receipt

This receipt acknowledges that Turnitin received your paper. Below you will find the receipt information regarding your submission.

The first page of your submissions is displayed below.

Submission author: Herpan Syafii Harahap
Assignment title: Quick Submit
Submission title: RELATIONSHIP BETWEEN LEVEL OF EDUCATION AND POST-S...
File name: ARTIKEL_2_-_1.docx
File size: 193.6K
Page count: 4
Word count: 2,845
Character count: 16,979
Submission date: 19-May-2021 12:09PM (UTC+0700)
Submission ID: 1589314959

<http://doi.org/10.21769/mj.v15n1.1111>
DOI: 10.21769/mj.v15n1.1111 (filed by Editor)
eISSN: 2542-5001 pISSN: 2407-0724
Assessing by DOI | Issue No. 215/KP7/2015

MALANG
NEUROLOGY
JOURNAL
MNJ
OPEN ACCESS

RESEARCH ARTICLE

RELATIONSHIP BETWEEN LEVEL OF EDUCATION AND POST-STROKE COGNITIVE STATUS IN HOSPITAL-BASED ISCHEMIC STROKE SURVIVORS

Herpan Syafii Harahap¹, Yanna Indrayana², Setyawati Ashi Putri³

Correspondence: herpanharahap@ubooa.co.id
¹Department of Neurology, Faculty of Medicine, Universitas Mataram, Mataram, Indonesia
²Department of Cardiology, Faculty of Medicine, Universitas Mataram, Mataram, Indonesia
³Department of Neurology Mataram General Hospital, Mataram, Indonesia

Article History (filed by Editor)
Received: 11.11.2020
Accepted: 11.11.2020
Published: 11.11.2020

Cite this as Vancouver style
First Author's last name followed by the initials of first and/or second name, Second Author's last name followed by the initials of first and/or second name, Third author's last name followed by the initials of first and/or second name, Article's title, Publisher, Year of published, Volume/Issue/page, DOI/PubMed link

ABSTRACT
Background: Post-ischemic stroke cognitive decline is significantly affecting the quality of life of its survivors. Its prevalence was about 74.1% which was mostly determined by the existing of vascular risk factors and cognitive reserve of the subjects. Level of education is one of determinants of cognitive reserve, a factor that affect the susceptibility of subjects to cognitive decline after experiencing ischemic stroke-related neuronal damage. Since level of education is protective for cognitive function, the intervention on it can reduce the occurrence of cognitive decline.
Objective: To investigate the relationship between level of education and cognitive status among hospital-based ischemic stroke survivors.
Methods: This cross-sectional study involved post-ischemic stroke outpatients in two hospitals. The data collected in this study were demographic data, including level of education, and clinical data as well. The level of education was categorized into <12 years and >12 years groups. Cognitive status was assessed using Montreal Cognitive Assessment in Indonesia version (MoCA-Ind) and subjects with scores of 26-30 were normal. The relationship between level of education as well as clinical data and cognitive status were analyzed using Chi-square test.
Results: There were 166 subjects eligible for this study (n=166). The mean age of subjects was 58 years and 68.67% of them were male. Cognitive decline were found 80.12% of subjects (n=133). The level of education was significantly associated with cognitive status of the subjects and hypertension as well.
Conclusion: The level of education had significant relationship with cognitive decline in the hospital-based population of ischemic stroke survivors.
Keywords: Post-ischemic stroke cognitive decline, level of education, cognitive reserve

Introduction

Cognitive decline is currently becoming an important complication of ischemic stroke. Post-stroke cognitive decline caused a significant decrease in quality of life and increase of mortality of the ischemic stroke survivors.¹ A systematic review and meta-analysis conducted by Pendlbury and Rothwell showed that the prevalence of cognitive decline among stroke survivors was about 74.1%.² Its prevalence was highest in the first year after onset of stroke.³ The wide range of its prevalence was mostly determined by several factors, including the population of the subjects, the time of the assessment of cognitive status, the instrument of the assessment of cognitive status used, the characteristics of vascular risk factors of the subjects, and cognitive reserve possessed by the subjects.⁴ Cognitive reserve is a concept, besides brain reserve, which theoretically determines whether an individual is susceptible to cognitive decline after experiencing neuronal damage.^{5,6} In the context of ischemic stroke, the concept of cognitive reserve may account for the discrepancy between the extent of ischemic stroke-related neuronal damage and the cognitive status and/or the significant cognitive improvement among stroke survivors who have similar clinical conditions. Several factors recognized as the determinants of cognitive reserve are occupation, routine physical activity, routine social activity, and level of education.⁷ The study focused on investigation of the significant relationship between level education as the representation of cognitive reserve and the occurrence of post-ischemic stroke cognitive decline has never been conducted, including in Indonesia. Most of the previous studies were focused on the investigation of the significant relationship between well-identified vascular risk factors, including age, hypertension, diabetes mellitus, dyslipidemia, and atrial fibrillation, and decrease of cognitive status among stroke survivors.^{8,9,10} A cohort study conducted by Levine et al. in overall stroke survivors, both ischemic and haemorrhagic subtypes of stroke, showed that low level of education was significantly decrease their cognitive status, prominently on executive domain.⁹

Page 1 of 4

RELATIONSHIP BETWEEN LEVEL OF EDUCATION AND POST- STROKE COGNITIVE STATUS IN HOSPITAL-BASED ISCHEMIC STROKE SURVIVORS

by Herpan Syafii Harahap

Submission date: 19-May-2021 12:09PM (UTC+0700)

Submission ID: 1589314959

File name: ARTIKEL_2_-_1.docx (193.6K)

Word count: 2845

Character count: 16979

RELATIONSHIP BETWEEN LEVEL OF EDUCATION AND POST-STROKE COGNITIVE STATUS IN HOSPITAL-BASED ISCHEMIC STROKE SURVIVORS

Herpan Syafii Harahap¹, Yanna Indrayana², Setyawati Asih Putri³

Correspondence: herpanharahap@yahoo.co.id

¹Department of Neurology, Faculty of Medicine, Universitas Mataram, Mataram, Indonesia

²Department of Cardiology, Faculty of Medicine, Universitas Mataram, Mataram, Indonesia

³Department of Neurology Mataram General Hospital, Mataram, Indonesia

Article History: (filled by Editor)

Received: xx x, xxxx

Accepted: xx x, xxxx

Published: xx x, xxxx

Cite this as: Vancouver style

First Author's last name followed by the initials of first and/or second name, Second Author's last name followed by the initials of first and/or second name, Third Author's last name followed by the initials of first and/or second name. Article's title. Publisher. Time of published; Volume(Issue);page. DOI/PubMed link

ABSTRACT

Background: Post-ischemic stroke cognitive decline is significantly affecting the quality of life of its survivors. Its prevalence was about 7-41% which was mostly determined by the existing of vascular risk factors and cognitive reserve of the subjects. Level of education is one of determinants of cognitive reserve, a factor that affect the susceptibility of subjects to cognitive decline after experiencing ischemic stroke-related neuronal damage. Since level of education is protective for cognitive function, the intervention on it can reduce the occurrence of cognitive decline.

Objective: To investigate the relationship between level of education and cognitive status among hospital-based ischemic stroke survivors.

Methods: This cross-sectional study involved post-ischemic stroke outpatients in two hospitals. The data collected in this study were demographic data, including level of education, and clinical data as well. The level of education was categorized into ≥ 12 years and < 12 years groups. Cognitive status was assessed using Montreal Cognitive Assessment in Indonesia version (MoCA-Indo) and subjects with score of > 20 were normal. The relationship between level of education as well as clinical data and cognitive status were analyzed using Chi-square test.

Results: There were 166 subjects eligible for this study (n=166). The mean age of subjects was 58 years and 68.67% of them were male. Cognitive decline were found 80.12% of subjects (n=133). The level of education was significantly associated with cognitive status of the subjects and hypertension as well.

Conclusion: The level of education had significant relationship with cognitive decline in the hospital-based population of ischemic stroke survivors.

Keywords: Post-ischemic stroke cognitive decline, level of education, cognitive reserve

Introduction

Cognitive decline is currently becoming an important complication of ischemic stroke. Post-stroke cognitive decline caused a significant decrease in quality of life and increase of mortality of the ischemic stroke survivors.¹ A systematic review and meta-analysis conducted by Pendlebury and Rothwell showed that the prevalence of cognitive decline among stroke survivors was about 7-41%.² Its prevalence was highest in the first year after onset of stroke.³ The wide range of its prevalence was mostly determined by several factors, including the population of the subjects, the time of the assessment of cognitive status, the instrument of the assessment of cognitive status used, the characteristics of vascular risk factors of the subjects, and cognitive reserve possessed by the subjects.⁴

Cognitive reserve is a concept, besides brain reserve, which theoretically determines whether an individual is susceptible to cognitive decline after experiencing neuronal damage.^{5,6} In the context of ischemic stroke, the concept of cognitive reserve may account for the discrepancy between

the extent of ischemic stroke-related neuronal damage and the cognitive status and/or the significant cognitive improvement among stroke survivors who have similar clinical conditions. Several factors recognized as the determinants of cognitive reserve are occupation, routine physical activity, routine social activity, and level of education.^{7,8}

The study focused on investigation of the significant relationship between level education as the representation of cognitive reserve and the occurrence of post-ischemic stroke cognitive decline has never been conducted, including in Indonesia. Most of the previous studies were focused on the investigation of the significant relationship between well-identified vascular risk factors, including age, hypertension, diabetes mellitus, dyslipidemia, and atrial fibrillation, and decrease of cognitive status among stroke survivors.^{9,10,11} A cohort study conducted by Levine et al. in overall stroke survivors, both ischemic and haemorrhagic subtype of stroke, showed that low level of education was significantly decrease their cognitive status, prominently on executive domain.¹²

This was the first study conducted to investigate the significant relationship between level of education and cognitive status in ischemic stroke survivors. The ischemic stroke survivors included in this study represented hospital-based population rather than community-based population.

4

Methods

This was a cross-sectional study involving all of ischemic stroke outpatients attending to two main referral hospital in West Nusa Tenggara, Mataram General Hospital and Mutiara Sukma Mental Hospital during period of January – June 2019 after fulfilling the inclusion but not exclusion criteria. The inclusion criteria were subjects with prior history of ischemic stroke, literate, fully conscious, voluntary participated, and speaking in Indonesia Language. The exclusion criteria were previous diagnosis of pre-stroke cognitive impairment, the evidence of clinically significant depression, and aphasia and motor dysfunctions that interfere cognitive assessment. Subjects were considered to have clinically significant depression if the results of screening test of depression using Beck Depression Inventory-II (BDI-II) showed score of 14-14.^{3,14} This study was conducted in January – June 2019. Written informed consent was obtained from all of the subjects. This study was conducted after getting approval from Komisi Etik Penelitian Kesehatan Universitas Mataram with the ethical clearance number 401/UN18.F7/ETIK/2018.

The data collected in this study were demographic and clinical characteristics, level of education and cognitive status of the subjects. The demographic data of subjects consisted of age and gender. The clinical data of subjects consisted of stroke onset, location of lesion, hypertension, diabetes mellitus, dyslipidemia, and atrial fibrillation. The level of education was categorized into two groups, i.e ≥ 12 years and < 12 years according to previous studies.^{15,16} This categorization of level education was aimed to distinguish subjects with higher level of education (≥ 12 years) and those with lower level of education (< 12 years), so that the relationship between level of education and cognitive status of these subjects could be analyzed. The instrument used in the assessment of cognitive status was Montreal Cognitive Assessment in Indonesia version (MoCA-Ina) scores ranged between 0-30. Based on this instrument, subjects with score of 26-30 were considered to have normal cognitive status, while those with score of 0-25 were considered to have cognitive impairment.¹⁷

The stroke onset was graded into four categories of time, i.e ≤ 3 months, > 3 to 12 months, > 12 to 24 months, and > 24 months. The location of lesion was determined by clinical examination and head CT scan results and categorized into location, right and left hemispheres respectively. All clinical data of subjects were obtained from questionnaire and their medical records.

The data were presented as mean (95%CI) unless otherwise stated. The relationship between level of education and cognitive status were analyzed using Chi-square test. The association between clinical data and cognitive status was also analyzed using chi-square test. Statistical significance was set to be $p < 0.05$. All data was analyzed using a computer program.

Results

Characteristics of the Subjects

There were 191 ischemic stroke outpatients recruited at the baseline survey during January – July 2019 and 166 of them fulfilled the inclusion but not exclusion criteria ($n=166$), as shown in Figure 1. These eligible subjects were obtained from Mataram General Hospital ($n=114$) and Mutiara Sukma Mental Hospital ($n=52$). The mean age of the subjects was 58 years and 68.7% of them were male. The demographic and clinical data as well as level of education is presented in Table 1.




Figure 1. Diagram of subjects recruitment in the present study

The Association between Level of Education as well as Clinical Characteristics and Cognitive Status

Cognitive decline in this study were found 80.1% of the total subjects ($n=133$). After grouping subjects into two groups, namely cognitive decline and normal cognitive status groups, level of education was significantly associated with cognitive function of the subjects and so was hypertension (Table 2).

Discussion

Cognitive decline among ischemic stroke survivors are now getting attention, since its prevalence is consistently quite high in several studies conducted in different population, either in community-based or hospital-based ischemic stroke survivors population. Epidemiologic studies conducted in several countries showed that the prevalence of post-ischemic stroke cognitive decline was in the range of 38-56.6%.^{18,19,20} Similar studies conducted in some regions of Indonesia, including Jakarta and Bandung using Montreal Cognitive Assessment in Indonesia version (MoCA-Ina) as cognitive assessment instrument, showed higher prevalence of post-ischemic stroke cognitive decline, in the range of 60.44-72%.^{17,21} Consistent with the studies conducted in some Jakarta and Bandung, the result of this study showed that its prevalence was about 80.1%.

Not all of ischemic stroke patients will be suffered from cognitive decline. The vulnerability of the stroke survivors to be suffered from cognitive decline is mainly determined by the existing vascular risk factors. Previous studies, both

Table 1. Demographic and Clinical Characteristics of the Subjects

Variables	Mean (95% CI), unless otherwise stated (n=166)
Age, years	58,67 (57.22 – 60.17)
Male, n (%)	114 (68.7)
Level of education, n (%)	
<12 years	74 (44.6)
≥12 years	92 (55.4)
Stroke onset, n (%)	
≤3 months	36 (21.7)
>3 – 12 months	74 (44.6)
>12 – 24 months	32 (19.3)
>24 months	24 (14.4)
Location of lesion, n (%)	
Right hemisphere	85 (51.2)
Left hemisphere	81 (48.8)
Hypertension, n (%)	128 (77.1)
Diabetes mellitus, n (%)	37 (22.3)
Dyslipidemia, n (%)	85 (51.2)
Atrial fibrillation, n (%)	19 (11.4)
MoCA-Ina score, n (%)	
Normal (0-25)	33 (19.88)
Cognitive decline (26-30)	133 (80.1)

MoCA-Ina=Montreal Cognitive Assessment in Indonesia version.

Table 2. The demographic and clinical characteristics between cognitive decline and normal cognitive status groups

Variables	Cognitive Status		p-value
	Cognitive Decline (n=133)	Normal (n=33)	
Gender, n (%)			
Male	92 (55.4)	22 (13.3)	0.781
Female	41 (24.7)	11 (6.6)	
Level of education, n (%)			
<12 years	66 (39.8)	8 (4.8)	0.009*
≥12 years	67 (40.4)	25 (15.1)	
Stroke onset, n (%)			
≤3 months	32 (19.3)	4 (2.4)	
>3 – 12 months	53 (31.9)	21 (12.7)	0.099
>12 – 24 months	27 (16.3)	5 (3.0)	
>24 months	21 (12.7)	3 (1.8)	
Location of lesion, n (%)			
Right hemisphere	66 (39.8)	19 (11.4)	0.413
Left hemisphere	67 (40.4)	14 (8.4)	
Hypertension, n (%)			
Yes	108 (65.1)	20 (12.0)	0.012*
No	25 (15.1)	13 (7.8)	
Diabetes mellitus			
Yes	30 (18.1)	7 (4.2)	0.868
No	103 (62.0)	26 (15.7)	
Dyslipidemia			
Yes	70 (42.2)	15 (9.0)	0.460
No	63 (38.0)	18 (10.8)	
Atrial fibrillation, n (%)			
Yes	17 (10.2)	2 (1.2)	0.371
No	116 (69.9)	31 (18.7)	

*indicate p<0.05 (significant)

original articles and systematic reviews, were more focused on the identification of the significant association of vascular risk factors of ischemic stroke and the prevalence of post-ischemic stroke cognitive decline.^{22,23,24} In general, vascular risk factors are divided into two main groups, non-modifiable and modifiable risk factors respectively. Non-modifiable risk factors commonly consist of age, gender and ethnicity, while modifiable risk factors consist of hypertension, diabetes mellitus, dyslipidemia, and atrial fibrillation.²⁵ In the pathophysiology of post-ischemic stroke cognitive decline, vascular risk factors mediate the occurrence of neurovascular unit dysfunction and cerebral blood flow dysregulation in certain brain regions serving particular cognitive functions, either independently or by their interaction with stroke event.⁹ Recent studies show that age is the main predictor for the occurrence of post-ischemic stroke cognitive decline compared to the overall existing vascular risk factors.²⁶ Among modifiable vascular risk factors, hypertension is the main predictor for the occurrence of post-ischemic stroke cognitive decline.²⁷ In this study, among various vascular risk factors identified, only hypertension had a significant relationship with the decline of post-ischemic stroke cognitive status. This result indicated a

significant consistency¹⁹ the relationship between hypertension and the prevalence of post-ischemic stroke cognitive decline among recent studies. The proposed mechanisms underlying the role of hypertension in the development of post-ischemic stroke cognitive decline is the occurrence of hypertension²-related small vessel disease and arterial stiffness resulting in both white matter and gray matter changes, especially in the prefrontal cortex, hippocampus, and inferior temporal cortex and inferior parietal lobules, brain structures important in carrying out cognitive functions.²⁸ The association between hypertension and post-ischemic stroke cognitive decline described above is strongly influenced by advancing age.²⁹ Antihypertensive treatment is beneficial in reducing the risk of stroke and post-stroke cognitive decline.³⁰

In addition to the existing vascular risk factors, the vulnerability of ischemic stroke survivors to be suffered from cognitive decline is also determined by their cognitive reserve status. The concept of cognitive reserve explains how an ischemic stroke survivor is able to perform a certain cognitive task properly and in flexible manner even though there is some degree of stroke-related brain dysfunction in certain areas that carry out this cognitive function.³¹ In this case, the concept of cognitive reserve can be proposed to explain the incompatibility of the relationship between the degree of ischemic stroke-related neuronal damage and the decline in cognitive status among ischemic stroke survivors.³² In such condition there is a neural compensation in which ischemic stroke-related neuronal damage in certain brain areas that serve specific cognitive functions will lead to recruitment as well as activation of compensatory networks that normally play an important role as a secondary area carrying out those specific cognitive functions.^{33,34} This neural compensation can explain the differences in clinical outcomes of post-ischemic stroke cognitive decline as well as the effectiveness in its recovery process among ischemic stroke survivors. The complexity of neural compensation itself, which describes the concept of cognitive reserve, is determined by the life experiences of a subject in working on a particular cognitive task which was obtained mainly through formal education, occupation, and daily physical as well as social activities prior ischemic stroke event.⁷

Recent studies have shown that level of education is a representation of the concept of cognitive reserve which has significant relationship with cognitive status in elderly population and Alzheimer's disease patients.^{7,35,36,37} The study aimed to investigate the impact of low level of education to the decline of cognitive status among stroke survivors regardless to the stroke subtype are still very limited and have not been conducted specifically in ischemic stroke subtype patients. A cohort study conducted by Levine et al. have shown showed that low level of education was significantly decrease the cognitive status of both ischemic and hemorrhagic stroke survivors, prominently on executive domain.¹² This was the first study focused on investigating the relationship between level of education and cognitive status in ischemic stroke outpatients representing hospital-based population. The results of this study showed that there was a significant relationship between the level of education and cognitive status in those population. These results indicated that level of education, beside hypertension, was an important risk factor for the decline in the post-ischemic stroke cognitive status in those subjects. As noted earlier, level of education as one of the determinants of cognitive reserve also played an important role in determining the successful of neural compensation mechanism which favor either in maintaining the cognitive status to remain intact or in accelerating the healing process of cognitive status among ischemic stroke survivors.^{33,34} Therefore, level of education is a better indicator of brain capacity to compensate for pathology efficiently either using the existing cognitive networks or recruiting the alternative networks.³⁸ Alternatively, stroke survivors with higher level of education generally have better lifestyle and adherence to treatment and this might explain the better cognitive status of stroke survivors in this group compared to group of stroke survivors with lower level of education.³⁹

The limitation of this study was that it did not identify the relationship between cognitive reserve determinants other than level of education, including recent occupation as well as the intensity of daily physical and social activities. Therefore, the significance relationship of level of education and cognitive status in this study could only represent the role of cognitive reserve partially. The heterogeneity of the subjects and their recall of detailed process of the occupation as well as the intensity of actual daily physical and social activities prior ischemic stroke event which were poorly documented. So, it made an obstacle in the analysis of the relationship between cognitive reserve status as a whole and cognitive status among ischemic stroke survivors. Therefore, a cohort study with systematic recording of the intensity of the occupation as well as the intensity of actual daily physical and social activities in those subjects in order to generalize their relationship to cognitive status is mandatory.

Conclusion

The conclusion of this of this study, the level of education as a representation of cognitive reserve had a significant relationship with the decline of post-ischemic stroke cognitive status in the hospital-based population of ischemic stroke survivors. A cohort study is mandatory to identify the role of education, occupation and the intensity of routine physical and social activities on cognitive status among ischemic stroke survivors so that the role of cognitive reserve as a whole can be generalized in compensating the decline of cognitive status in those population. In addition, hypertension was consistently still having significant relationship with the decline of cognitive status among those population, regarding to previous studies.

Acknowledgement

We would like to acknowledge the contribution of Made Rika Anastasia Pratiwi and Hardinata.

RELATIONSHIP BETWEEN LEVEL OF EDUCATION AND POST-STROKE COGNITIVE STATUS IN HOSPITAL-BASED ISCHEMIC STROKE SURVIVORS

ORIGINALITY REPORT

11%

SIMILARITY INDEX

8%

INTERNET SOURCES

5%

PUBLICATIONS

%

STUDENT PAPERS

PRIMARY SOURCES

1	mnj.ub.ac.id Internet Source	2%
2	Tu, Qiuyun, Binrong Ding, Xia Yang, Song Bai, Junshi Tu, Xiao Liu, Ranran Wang, Jinghua Tao, Hui Jin, Yiqun Wang, and Xiangqi Tang. "The current situation on vascular cognitive impairment after ischemic stroke in Changsha", Archives of Gerontology and Geriatrics, 2013. Publication	1%
3	bjo.bmj.com Internet Source	1%
4	www.ijdr.in Internet Source	1%
5	"Original Research", European Heart Journal Supplements, 2017 Publication	1%
6	repository.unair.ac.id Internet Source	1%

7	bmccardiovascdisord.biomedcentral.com Internet Source	1 %
8	"Poster Sessions", Pediatric Pulmonology, 2019 Publication	<1 %
9	global.oup.com Internet Source	<1 %
10	link.springer.com Internet Source	<1 %
11	repository.unam.na Internet Source	<1 %
12	mafiadoc.com Internet Source	<1 %
13	Hee-Joon Bae, Dilip K. Pandey. "Treatment of vascular dementia: Evidence from epidemiologic studies", Current Treatment Options in Cardiovascular Medicine, 2005 Publication	<1 %
14	jisponline.com Internet Source	<1 %
15	"Posters", Pediatric Pulmonology, 2018 Publication	<1 %
16	Buja, P.. "Moderate-to-severe ischemic mitral regurgitation and multivessel coronary artery disease: Impact of different treatment on	<1 %

survival and rehospitalization", International
Journal of Cardiology, 20060728

Publication

17

Mohammad Abufaraj, Zaid Eyadat,
Mohammed Qussay Al-sabbagh, Abdullah
Nimer et al. "Gender-based disparities on
health indices during COVID-19 crisis: a
nationwide cross-sectional study in Jordan",
International Journal for Equity in Health, 2021

Publication

<1 %

18

scielo.isciii.es

Internet Source

<1 %

19

www.ncbi.nlm.nih.gov

Internet Source

<1 %

20

eprints.whiterose.ac.uk

Internet Source

<1 %

21

"Abstracts Presented at the Thirty-Fourth
Annual International Neuropsychological
Society Conference, February 1-4, 2006,
Boston, Massachusetts, USA", Journal of the
International Neuropsychological Society,
2006

Publication

<1 %

Exclude quotes Off

Exclude matches Off

Exclude bibliography Off

