




# JURNAL FARMASI SAINS DAN KOMUNITAS

Journal of Pharmaceutical Sciences & Community



Accredited by RISTEKDIKTI  
 No. 30/E/KPT/2018





HOME ABOUT LOGIN REGISTER SEARCH CURRENT ARCHIVES  
ANNOUNCEMENTS

Home > Archives > Vol 16, No 2 (2019)

## Vol 16, No 2 (2019)

### Table of Contents

#### Articles

<a href="#">ANTI-DIARRHEAL ACTIVITY OF AQUEOUS EXTRACT OF NAGASARI FLOWERS (Mesua ferrea L.) IN BALB/c MICE INDUCED BY Escherichia coli</a>	<a href="#">PDF</a>	50-55
Putu Monik Ananta Puspitarini, Iman Surya Pratama, Bambang Fajar Suryadi		
<a href="#">LIPID AND SILVER NANOPARTICLES GELS FORMULATION OF TEMPEH EXTRACT</a>	<a href="#">PDF</a>	56-62
Felicia Satya Christania, Rini Dwiastuti, Sri Hartati Yuliani		
<a href="#">ANTIOXIDANT AND ANTICANCER ACTIVITIES OF MURBEI (Morus alba L.) STEM EXTRACT ON IN VITRO WIDR CANCER CELLS</a>	<a href="#">PDF</a>	63-67
Asril Burhan, Akbar Awaluddin, Zulham Zulham, Burhanuddin Taebe, Abdul Gafur		
<a href="#">PROFILE OF ANTICANCER ACTIVITIES OF BROTOWALI (Tinospora crispa L.) PLANTS OF VARIOUS REGIONS IN EAST JAWA</a>	<a href="#">PDF</a>	68-77
Roihatul Mutiah, Laily Nurul Azizah, Rahmi Annisa, Anik Listyana		
<a href="#">ANTIOXIDANT ACTIVITY, TOTAL PHENOLIC, AND FLAVONOID CONTENTS OF THE EXTRACT OF ENDOPHYTIC FUNGI DERIVED FROM TURMERIC (Curcuma longa) LEAVES</a>	<a href="#">PDF</a>	78-85
Eris Septiana, Siti Irma Rahmawati, Fauzia Nurul Izzati, Partomuan Simanjuntak		
<a href="#">THE SUB-CHRONIC TOXICITY TEST OF MENIRAN (Phyllanthus niruri L.) AND PEGAGAN (Centella asiatica) EXTRACT IN WISTAR STRAIN RATS ON LIVER AND KIDNEY FUNCTION</a>	<a href="#">PDF</a>	86-95
Endang Darmawan, Iin Narwanti, Siti Fatmawati Fatimah, Ira Aprilia Wulandari, Ria Putri Salma, Dzulhaifa Dzulhaifa		
<a href="#">THE CORRELATION OF TOTAL FLAVONOID AND TOTAL PHENOLIC WITH ANTIOXIDANT ACTIVITY OF SINGLE BULB GARLIC (Allium Sativum) FROM TAWANGMANGU AND MAGETAN</a>	<a href="#">PDF</a>	96-103
Ika Buana Januarti, Hudan Taufiq, Sulistyaningsih Sulistyaningsih		

Focus and Scope

Author Guidelines

Online Submission

Editorial Team

Publication Ethics

Journal History

Order Journal

Visitor Statistics

#### USER

Username

Password

Remember me

#### NOTIFICATIONS

- [View](#)
- [Subscribe](#)

#### JOURNAL CONTENT

Search

Search Scope

All

#### Browse

- [By Issue](#)
- [By Author](#)
- [By Title](#)
- [Other Journals](#)

#### TOOLS

grammarly

MENDELEY

turnitin

Article Template



Jurnal Farmasi Sains dan Komunitas (Journal of Pharmaceutical Sciences and Community)

Published by [Faculty of Pharmacy, Universitas Sanata Dharma Yogyakarta](#)



This work is licensed under a [Creative Commons Attribution 4.0 International License](#).

00053917 [JPSC Stats](#)



### Visitors

 48,629	 145
 5,403	 107
 553	 106
 213	 106
 149	 96

 FLAG counter

### KEYWORDS

[Eurycoma longifolia](#)  
[adherence](#) [analgesic](#)  
[analysis](#)  
[antioxidant](#)  
[aqueous extract](#)  
[asthma](#) [breast cancer](#)  
[carbon tetrachloride](#) [co-](#)  
[processed excipient](#) [cream](#)  
[flavonoid gel](#)  
[hypertension](#)  
[liposome](#) [partide size](#)  
[physical activity](#) [self-](#)  
[medication](#) [sonication](#)  
[tempeh extract](#) [virtual](#)  
[screening.](#)



HOME ABOUT LOGIN REGISTER SEARCH CURRENT ARCHIVES  
ANNOUNCEMENTS

Home > About the Journal > **Editorial Team**

## Editorial Team

### Editor in Chief

[Enade Perdana Istyastono](#), Faculty of Pharmacy, Sanata Dharma University, Yogyakarta

### Vice Editor in Chief

[Phebe Hendra](#), Faculty of Pharmacy, Sanata Dharma University, Yogyakarta, Indonesia  
[Maywan Hariono](#), Faculty of Pharmacy, Sanata Dharma University, Yogyakarta, Indonesia  
[Yosef Wijoyo](#), Faculty of Pharmacy, Sanata Dharma University, Yogyakarta, Indonesia

### Managing Editor

[Michael Raharja Gani](#), Faculty of Pharmacy, Universitas Sanata Dharma, Indonesia

### Editorial Board

[Aris Widayati](#), Faculty of Pharmacy, Universitas Sanata Dharma, Indonesia  
[Aty Widawaruyanti](#), Faculty of Pharmacy, Universitas Airlangga, Indonesia  
[Auliya A. Suwantika](#), Faculty of Pharmacy, Universitas Padjadjaran, Indonesia  
[Bandana Saini](#), Faculty of Pharmacy, University of Sydney, Australia  
[Christine Patramurti](#), Faculty of Pharmacy, Sanata Dharma University, Yogyakarta, Indonesia  
[Erna Tri Wulandari](#), Faculty of Pharmacy, Sanata Dharma University, Yogyakarta, Indonesia  
[Florentinus Dika Octa Riswanto](#), Faculty of Pharmacy, Sanata Dharma University, Yogyakarta, Indonesia  
[Irma Melyani Puspitasari](#), Faculty of Pharmacy, Universitas Padjadjaran, Indonesia  
[Monet M. Loquias](#), College of Pharmacy, University of the Philippines, Manila, Philippines  
[Phayom Sookaneknun](#), Faculty of Pharmacy, Mahasarakham University, Thailand  
[Phebe Hendra](#), Faculty of Pharmacy, Sanata Dharma University, Yogyakarta, Indonesia  
[P.T. Thomas](#), Faculty of Health and Medical Sciences, Taylor's University, Malaysia  
[Rano Kurnia Sinuraya](#), Faculty of Pharmacy, Universitas Padjadjaran, Indonesia  
[Rizky Abdulah](#), Faculty of Pharmacy, Universitas Padjadjaran, Indonesia  
[Sri Hartati Yuliani](#), Faculty of Pharmacy, Sanata Dharma University, Yogyakarta, Indonesia  
[Titien Siwi Hartayu](#), Faculty of Pharmacy, Sanata Dharma University, Yogyakarta, Indonesia  
[Yashwant Pathak](#), Faculty Affairs, College of Pharmacy, University of South Florida, United States  
[Yustina Sri Hartini](#), Faculty of Pharmacy, Sanata Dharma University, Yogyakarta, Indonesia

### Administrative Editor

[Christianus Heru Setiawan](#), Faculty of Pharmacy, Sanata Dharma University, Yogyakarta, Indonesia  
[Dina Christin Ayuning Putri](#), Faculty of Pharmacy, Sanata Dharma University, Yogyakarta, Indonesia  
[Putu Dyana Christasani](#), Faculty of Pharmacy, Sanata Dharma University, Yogyakarta, Indonesia

Focus and Scope

Author Guidelines

Online Submission

Editorial Team

Publication Ethics

Journal History

Order Journal

Visitor Statistics

#### USER

Username

Password

Remember me

#### NOTIFICATIONS

- [View](#)
- [Subscribe](#)

#### JOURNAL CONTENT

Search

Search Scope

All

#### Browse

- [By Issue](#)
- [By Author](#)
- [By Title](#)
- [Other Journals](#)

#### TOOLS

grammarly

MENDELEY

turnitin

Article  
DOC Template



Jurnal Farmasi Sains dan Komunitas (Journal of Pharmaceutical Sciences and Community)

Published by [Faculty of Pharmacy, Universitas Sanata Dharma Yogyakarta](#)



This work is licensed under a [Creative Commons Attribution 4.0 International License](https://creativecommons.org/licenses/by/4.0/).

00053917 [JPSC Stats](#)



**KEYWORDS**

[Eurycoma longifolia](#)  
[adherence](#) [analgesic](#)  
[analysis](#)  
[antioxidant](#)  
[aqueous extract](#)  
[asthma](#) [breast cancer](#)  
[carbon tetrachloride](#) [co-](#)  
[processed excipient](#) [cream](#)  
[flavonoid gel](#)  
[hypertension](#)  
[liposome](#) [particle size](#)  
[physical activity](#) [self-](#)  
[medication](#) [sonication](#)  
[tempeh extract](#) [virtual](#)  
[screening](#).

## ANTI-DIARRHEAL ACTIVITY OF AQUEOUS EXTRACT OF NAGASARI FLOWERS (*Mesua ferrea* L.) IN BALB/c MICE INDUCED BY *Escherichia coli*

Putu Monik Ananta Puspitarini<sup>1\*</sup>, Iman Surya Pratama<sup>1</sup>, Bambang Fajar Suryadi<sup>2</sup>

<sup>1</sup>Pharmacy Programme, Faculty of Medicine, Mataram University, Majapahit Street No. 62, Mataram 83115, Indonesia

<sup>2</sup>Biology Programme, Faculty of Mathematics and Natural Science, Mataram University, Majapahit Street No. 62, Mataram 83115, Indonesia

Received April 1, 2019; Accepted September 29, 2019

### ABSTRACT

Nagasari (*Mesua ferrea* L.) is empirically used for management for diarrhea in Nusa Tenggara Barat Province. However, the safety and efficacy of the nagasari have not been scientifically validated in an animal model. This study evaluated the activity of aqueous extract of the nagasari flowers in BALB/c mice induced by *Escherichia coli* to determine the effective dosage for anti-diarrheal effect. The aqueous extract of nagasari flowers (0,4% (w/v); 0,8% (w/v); and 1,6% (w/v)) was administered orally to 3 groups of mice, whereas negative and positive control received aquadestilate and ciprofloxacin 130 mg/kgBW respectively orally once daily for 3 days. The frequency and consistency of feces were observed then the total colony of *E. coli* in feces was calculated using colony counter. Aqueous extract of nagasari flowers at concentration 1,6% showed significant inhibitory activity against *Escherichia coli* comparable with positive controls (diarrhea index 0,425;  $p > 0,05$  and total colony of *E. coli* in feces 100,2;  $p > 0,05$ ). This study provides scientific support for the traditional use of aqueous extract of nagasari flowers for the treatment of diarrheal diseases.

**Keywords:** anti-diarrheal; Aqueous extract; diarrhea index; *Escherichia coli*; nagasari.

### INTRODUCTION

Diarrhea is an abnormal event that characterized by an increased in the frequency of defecate more than 3 times in 24 hours accompanied by the consistency of liquid stool (Kotloff *et al.*, 2013).

Based on the Health Profile in Nusa Tenggara Barat (NTB) Province, the number of diarrhea patients had not experienced a significant decrease that is 158.993 and 175.361 in the year of 2015 and 2017 (Dinkes RI, 2015; 2018). Diarrhea can be caused by infectious agents such as bacteria, viruses and parasites, including *Escherichia coli* which includes Gram-negative bacteria.

Ciprofloxacin as the first-line in the treatment of diarrhea but has limitations such as side effects and resistance if used over a long period of time. In addition, this drug is a second-line treatment for tuberculosis

XDR/MDR (Extensively Drug Resistance/Multi-Drug Resistance) so it must be used wisely (Raini, 2016). Herbal medicine is an option to be developed as anti-diarrhea.

Nagasari (*Mesua ferrea* L.) is one of the plant used traditionally as anti-diarrhea. In Dharma Usada Kuranta Bolong, states that part of the nagasari plant can be used as a diarrhea medicine (Listiwati *et al.*, 2016). *Mesua ferrea* L. flowers is illustrated in Figure 1.

In vitro, the methanol extract of all parts of the nagasari flowers at concentration of 10 µg/mL, 50 µg/mL and >200 µg/mL showed the minimum inhibitory concentration values of *E. coli* of 2, 6 and 14 strains. Flavonoids mesuaferone-A and mesuaferone-B are through to acts in the antibacterial effect (Mazumder *et al.*, 2004).

\*Corresponding author: Putu Monik Ananta Puspitarini  
Email: [monikananta1@gmail.com](mailto:monikananta1@gmail.com)

In vivo scientific studies of nagasari flower extract as anti-diarrhea are still limited. Therefore, this study evaluated the activity of aqueous extract of the nagasari flowers in Balb/c mice induced diarrhea by *E. coli* in decreasing diarrhea index and total of *E. coli* colony in feces.

## METHODS

### Collection and Identification of Flowers

The fresh flowers of *Mesua ferrea* were collected from Lingsar, West Lombok in January 2018 and identified by Herbarium Bandungense SITH, Bandung Institute of Technology, Bandung No. 1493/11.CO2.2/PL/2018.

### Drying and Grinding

The whole parts of flowers were washed, dried under shade for five days. The powdered of dry flowers were prepared with a mechanical grinder and passed through a 35-mesh sieves. The powder was stored in a well-closed container until the analysis was commenced.

### Extraction

The powder of *M. ferrea* (nagasari) flowers were boiled in aquadestilate (0,8% w/v) during 15 minutes. The aqueous extract was stored properly (Hamad *et al*, 2017).

### Bacterial Preparation

*Escherichia coli* ATCC 25922 strain was isolated from urine samples. This strain is part of the pathogen from the Yogyakarta Hall Health's Laboratory. The bacteria were grown on Nutrient Agar (NA) (Oxoid) (14 gram dehydrated NA in 500 mL aquadest) then transferred to the *Eosin Methylene Blue Agar* (EMBA) (Oxoid) (18,75 gram dehydrated EMBA in 500 mL aquadest (from Health Research and Development Installation of NTB Provincial Hospital) as selective media and Gram staining was carried out for the identification process accordance with the Bergey's Manual of Systematic Bacteriology (Whitman, 2010).



Figure 1. *Mesua ferrea* L. flowers

The bacteria were suspended in sterile physiological saline (0,9 gram NaCl (Merck) in 100 mL aquadest) until a concentration of approximately  $10^6$  CFU/mL (Arief *et al*, 2010). The concentration of bacteria was calculated using Neubauer methods.

### Experimental Animals

All procedures involving animals were approved by the Ethics Committee Faculty of Medicine of The Mataram University No. 91/UN18.8/ETIK/2018.

A total of 30 BALB/c mice weight between 25-40 g obtained from the Immunology Laboratory of Mataram University were used in this study. The mice were acclimatized under the standard environmental condition for 7 days. All mice had free access to food and destilate water *adlibitum* throughout the experimental period. The mice were divided into 6 groups randomly with 6 mice per group as follow: normal group (normal mice received destilate water 0,5 mL), negative control group (diarrhea mice received destilate water 0,5 mL), positive control group (diarrhea mice received ciprofloxacin 130 mg/kg BW (13 mg ciprofloxacin (Hexpharm Jaya) powder in 5 mL aquadest)), and three group of treatment (0,4% w/v; 0,8% w/v; 1,6% w/v).

### Assessment of In-vivo Anti-diarrheal Activity *Escherichia coli* induced diarrhea

Each mice was administrated orally with 100  $\mu$ L of  $10^6$  CFU of *E. coli* except the normal group to induced diarrhea (Astawan *et*

al, 2009). After 30 minutes, each mouse is treated with the extract and ciprofloxacin.

All mice were treated every morning for 3 days. After the treatment, all mice were placed in individual cages, the floor of which was covered with filter paper. The number and consistency of stool were recorded for 6 hours.

The severity quantitatively of diarrhea was defined using three indexes (1) *Loose Stool Incidence Rate / LSIR*, (2) *Average Loose Stool Grade / ALSG*, (3) *Diarrhea Index / DI* and reduction in the total of colony of *E. coli* in feces (Hui *et al*, 2012; Han *et al*, 2014; Yu *et al*, 2017).

**Table I.** Effect of aqueous extract of *M. ferrea* (nagasari) flowers in decreasing total colony of *Escherichia coli* in feces

Group		Before treatment (CFU)	After treatment (CFU)
I	<i>E. coli</i> + AEONF 0,4%	142,2 ± 97,9	155,6 ± 96,9
II	<i>E. coli</i> + AEONF 0,8%	97,6 ± 42,4	41,4 ± 31,2
III	<i>E. coli</i> + AEONF 1,6%	122,2 ± 51,5	22 <sup>a)</sup> ± 15,9
IV	<i>E. coli</i> + Cip	149 ± 34,7	29 ± 8,8
V	<i>E. coli</i> + Destilated water	128,2 ± 51,4	136,4 ± 48,6
VI	Destilated water	0 ± 0	0 <sup>a)</sup> ± 0

Note. AEONF = Aqueous extract of Nagasari flowers, Cip = Ciprofloxacin, <sup>a)</sup> p > 0,05 comparable with positive controls.

**Table II.** Effect of aqueous extract of *M. ferrea* (nagasari) flowers in decreasing LSIR

Group		LSIR day 1	LSIR day 3
I	<i>E. coli</i> + AEONF 0,4%	0,161 ± 0,06	0,144 ± 0,033
II	<i>E. coli</i> + AEONF 0,8%	0,273 ± 0,202	0,036 <sup>a)</sup> ± 0,05
III	<i>E. coli</i> + AEONF 1,6%	0,355 ± 0,25	0 <sup>a)</sup> ± 0
IV	<i>E. coli</i> + Cip	0,194 ± 0,169	0,040 ± 0,089
V	<i>E. coli</i> + Destilate water	0,367 ± 0,031	0,315 ± 0,159
VI	Destilate water	0 ± 0	0 <sup>a)</sup> ± 0

Note. AEONF = Aqueous extract of Nagasari flowers, Cip = Ciprofloxacin, <sup>a)</sup> p > 0,05 comparable with positive controls.

**Table III.** Effect of aqueous extract of *M. ferrea* (nagasari) flowers in decreasing ALSG


Group		ALSG day 1	ALSG day 3
I	<i>E. coli</i> + AEONF 0,4%	1,04 ± 0,089	1,2 ± 0,447
II	<i>E. coli</i> + AEONF 0,8%	1 ± 1	0,8 <sup>a)</sup> ± 1,095
III	<i>E. coli</i> + AEONF 1,6%	1,131 ± 0,144	0 <sup>a)</sup> ± 0
IV	<i>E. coli</i> + Cip	1,167 ± 0,745	0,267 ± 0,596
V	<i>E. coli</i> + Destilate water	1,218 ± 0,287	1,506 ± 0,389
VI	Destilate water	0 ± 0	0 <sup>a)</sup> ± 0

Note. AEONF = Aqueous extract of Nagasari flowers, Cip = Ciprofloxacin, <sup>a)</sup> p > 0,05 comparable with positive controls.


**Table IV.** Effect of aqueous extract of *M. ferrea* (nagasari) flowers in decreasing DI

Group	DI day 1	DI day 3
I <i>E. coli</i> + AEONF 0,4%	0,170 ± 0,074	0,164 ± 0,03
II <i>E. coli</i> + AEONF 0,8%	0,273 ± 0,202	0,362 ± 0,099
III <i>E. coli</i> + AEONF 1,6%	0,425 ± 0,33	0,000 <sup>a)</sup> ± 0
IV <i>E. coli</i> + Cip	0,306 ± 0,251	0,053 ± 0
V <i>E. coli</i> + Destilate water	0,458 ± 0,108	0,638 ± 0,713
VI Destilate water	0,000 ± 0	0,000 <sup>a)</sup> ± 0

Note. AEONF = Aqueous extract of Nagasari flowers, Cip = Ciprofloxacin, <sup>a)</sup> p > 0,05 comparable with positive controls.



**Figure 2:** Effect of aqueous extract of *Mesua ferrea* in decreasing total colony of *Escherichia coli* in feces. AEONF = Aqueous extract of Nagasari flowers, Cip = Ciprofloxacin, <sup>a)</sup> p > 0,05 comparable with positive controls.



**Figure 3:** Effect of aqueous extract of *Mesua ferrea* in decreasing LSIR. AEONF = Aqueous extract of Nagasari flowers, Cip = Ciprofloxacin, <sup>a)</sup> p > 0,05 comparable with positive controls.

**Data Collection and Analysis Samples**

**Total colony of *Escherichia coli* in feces**

The fecal sample as much 50 mg was placed in 2 mL Eppendorf tube, to which distilled water was added. The sample was diluted 4 times dilution. The fourth dilution (100 µL) was fertilized by spread plate method on EMBA, then incubated at 37°C for 24 hours. Metallic green color on media

showed a positive result of *Escherichia coli* in feces (Zhang *et al*, 2010; Analytic Jena, 2011).

Calculation of the total colony of *Escherichia coli* was carried out using a colony counter (Stuart™ Scientific colony counter Model SC5, AC/DC input 220 V AC). Mice infected if the total colony of bacteria in feces are 10<sup>5</sup> CFU (Newsome *et al*, 1987). Calculation formula for the total colony of bacteria in feces:

$$N = \frac{EC}{((1 \times n) \times d)} \dots \dots \dots (1)$$


N : total of colonies (mL/g)  
 n : total of plates in the first dilution;  
 d : the level of dilution obtained from the first cup calculated.

**Three indices of severity diarrhea**


The LSIR the ratio of the number of loose stools to the total stools within an animal. Loose stool grade describes the degree of loose stools, based on the diameter of the stool on the filter papers. The LSG was classified into four grades according to the diameter of loose stools; Grade 1 (< 1 cm), Grade 2 (1~< 2 cm), Grade 3 (2~3 cm) and Grade 4 (>3cm). The ALSG is the ratio of the sum of the LSG of each loose stool to the total number of loose stools within each mice. The DI is the result of LSR multiplying ALSG (Hui *et al*, 2012; Han *et al*, 2014; Yu *et al*, 2017).

**Statistical analysis**

The data were analyzed by using Kruskal Wallis procedures, followed by Mann Whitney. Values of p < 0,05 were showed a significant difference between the effect of aqueous extract with positive controls statistically using SPSS 16 software.



**Figure 4:** Effect of aqueous extract of *Mesua ferrea* in decreasing ALSG. AEONF = Aqueous extract of Nagasari flowers, Cip = Ciprofloxacin, <sup>a)</sup>  $p > 0,05$  comparable with positive controls.



**Figure 5:** Effect of aqueous extract of *Mesua ferrea* in decreasing diarrhea index. AEONF = Aqueous extract of Nagasari flowers, Cip = Ciprofloxacin, <sup>a)</sup>  $p > 0,05$  comparable with positive controls.

## RESULTS AND DISCUSSION

The AEONF (Table I and Figure 2) at concentration 1,6% showed the highest decreasing in the total colony of *Escherichia coli* compared to other concentration and the antibacterial effect compared to positive controls ( $p > 0,05$ ).

The LSIR (Table II and Figure 3) describes the level of frequency of defecation. The LSIR did not differ between positive control and treatment group. The AEONF at concentration 1,6% showed the highest decreasing in the frequency of defecation compared to other concentration and the effect comparable with positive controls ( $p > 0,05$ ).

The ALSG (Table III and Figure 4) describes the level of consistency of feces, which is viewed from the size of the feces diameter on filter paper. The ALSG did not differ between positive control and treatment group (AEONF at concentration 0,8% and

1,6%). AEONF at concentration 1,6% showed the highest decreasing in the consistency of feces compared to AEONF at concentration 0,8% and the effect comparable with positive controls ( $p > 0,05$ ).

Diarrhea index (Table IV and Figure 5) is an accumulation of LSIR and ALSG. The DI did not differ between positive control and treatment group. The AEONF at concentration 1,6% the highest decreasing in the values of DI significantly different with negative control. The AEONF at concentration 1,6% showed the highest decreasing in the DI compared to other concentration and the effect comparable with positive controls ( $p > 0,05$ ).

In this study, the AEONF showed anti-diarrhea activity as evident from the inhibition of the increased diarrhea index induced by *Escherichia coli*.

Based on these four parameters, the AEONF at concentration 1,6% showed the highest effectiveness as specific anti-diarrheals. Mazumder *et al* (2004) have reported, nagasari flowers contain several flavonoid compounds. Allegedly, the content of flavonoids and tannins causes a decreasing in all four parameters.

Flavonoids can inhibit Gram-negative or Gram-positive bacteria. Alcohol groups in flavonoids can bind to peptidoglycan and lipopolysaccharide on the cell wall so that it will damage bacteria cell membranes. The Flavonoids have mechanisms such as ciprofloxacin by inhibiting the function of gyrase DNA by which interferes with the process of bacterial replication obtained (Jawetz *et al*, 2013; Mufti *et al*, 2017).

The role of tannin in adhesion cell and enzyme inactivating and disturb with protein transport in the inner layer of the cell. In addition, tannin bind to cell wall polypeptides which cause the formation of cell walls to be less than perfect. This cause osmotic pressure in cells so that bacterial cells die and lysis (Okuda, 2005; Smullen *et al*, 2007).

In this study, no signs of toxicity and mortality were found in mice so it was safe to use. Toxicity data on ethanol extracts of nagasari flowers doses of 500-2000 mg/kgBW also did not cause signs of acute toxicity in

mice (Tiwari *et al*, 2012). However, further toxicity tests on the AEONF need to be carried out.

## CONCLUSION

Aqueous extract of nagasari flowers at concentration 1,6% has the most effective specific anti-diarrheal activity indicated by the highest decreasing in the total colony of *Escherichia coli* and diarrhea index.

## REFERENCES

- Analytic Jena. 2012. innuPREP Stool DNA Kit. Analytic Jena AG. URL <https://www.analytikjena.com/products/kitsassaysreagents/kitsfordnarna-extraction/innuprep-stooldnakit/> (accessed 17.9.2019)
- Arief, I.I., Jenie, B.S.L., Astawan, M., Witarto, A.B., 2010. Efektivitas Probiotik *Lactobacillus plantarum* 2 C12 dan *Lactobacillus acidophilus* 2B4 sebagai Pencegah Diare pada Tikus Percobaan. *Media Peternakan*. 137-43.
- Astawan, M., Wresdiyati, T., Arief, I. I., Febiyanti, D., 2011. Gambaran Hematologi Tikus Putih (*Rattus norvegicus*) yang Diinfeksi *Escherichia coli* Enteropatogenik dan diberikan Probiotik. *Media Peternakan*. 7-13.
- Dinas Kesehatan Provinsi NTB. 2015. Profil Kesehatan Provinsi Nusa Tenggara Barat.
- Hamad, A., Anggraeni, W., Hartanti, D., 2017. Potensi Infusa Jahe (*Zingiber officinale* R) sebagai Bahan Pengawet alami pada Tahu dan Daging Ayam Segar. *Jurnal Aplikasi Teknologi Pangan*. 6(4), 177-183.
- Han, X., Pang, Y., Liu, S., Tan, Z., Tang, S., Zhou, C., Wang, M., Xiao, W., 2014. Antidiarrhea and Antioxidant Activities of Honokiol Extract from *Magnolia officinalis* Cortex in Mice. *Tropical Journal of Pharmaceutical Research*. 13(10), 1643-1651.
- Hui, H., Chen, M., Li, G., Feng, H., Zang, L., Xi, Y., Dong, T., Zhou, X., 2013. Andrographolide Attenuates Senna- and Castor Oil-induced Diarrhea in Mice. *Latin American Journal of Pharmacy*. 32(8), 1113-1117.
- Jawetz, E., Melnick, J., Adelberg. 2013. *Medical Microbiology*. Edisi ke-26. Jakarta: EGC.
- Kementerian Kesehatan Republik Indonesia. 2018. Data dan Informasi Kesehatan Indonesia 2017. NTB: Kementerian Kesehatan Republik Indonesia. URL [http://www.depkes.go.id/resources/download/pusdatin/profilkesehatanindonesia/DatadanInformasi\\_P-rofilKesehatanIndonesia2018.p-df](http://www.depkes.go.id/resources/download/pusdatin/profilkesehatanindonesia/DatadanInformasi_P-rofilKesehatanIndonesia2018.p-df) (accessed 18.11.2018).
- Kotloff, K.L., Narato, J.P., Blackwelder, W.C., Nasrin, D., Farag, T.H., 2013. Burden and Aetiology of Diarrhoeal Disease in Infants and Young Children in Developing Countries (the Global Enteric Multicenter Study, GEMS): a Prospective, Case-Control Study. *CrossMark*. 382(9888), 209-222.
- Listiawati, N.P., Juniawan, I.B.H., Anom, A.A.I.A., Sudarsana, I.W.G., Wibawa, I.G.J.S., 2016. Dharma Usadha Kuranta Bolong Klimosadha Sang Hyang Klimosadha.
- Mazumder, R., Dasridar, S.G., Basu, S.P., Mazumder, A., Singh, S.K., 2004. Antibacterial Potentiality of *Mesua ferrea* Linn. Flowers. *Phytotherapy Research*. 18, 824-26.
- Mufti, N., Bahar, E., Arisanti, D., 2017. Uji Daya Hambat Ekstrak Daun Sawo terhadap Bakteri *Escherichia coli* secara *In Vitro*. *Jurnal Kesehatan Andalas*. 6(2), 289-294.
- Newsome, P.M., Burgess, M.N., Burgess, M.R., Coney, K.A., Goddard, J.A.E., Morris, J.A., 1987. A Model of Acute Infectious Neonatal Diarrhoea. *Journal of Medical Microbiology*. 23(1), 19-28.