

Understanding the Uniqueness of Artocarpus Flavonoids: Isolation and Structure Elucidation of Cycloartocarpin from the Roots of Artocarpus altilis

by Aliefman Hakim Dkk

Submission date: 02-Aug-2022 07:42AM (UTC+0700)

Submission ID: 1877896314

File name: acs.jchemed.0c00199.pdf (634.87K)

Word count: 2658

Character count: 15090

Understanding the Uniqueness of Artocarpus Flavonoids: Isolation and Structure Elucidation of Cycloartocarpin from the Roots of *Artocarpus altilis*

Aliefman Hakim,* A. Wahab Jufri, Jamaluddin, Supriadi, and Putri Ayu Mutmainnah



Cite This: *J. Chem. Educ.* 2020, 97, 4133–4136



Read Online

ACCESS |



Metrics & More



Article Recommendations



Supporting Information

ABSTRACT: Secondary metabolite compounds contained in each plant genus have certain characteristics, which is true also of the flavonoid in the genus *Artocarpus*. In the experiment described here, third-year undergraduate students performed the isolation and characterization of cycloartocarpin from the roots of *Artocarpus altilis* (commonly known as breadfruit), giving them experience in skills such as extraction, fractionation, purification, and the structural elucidation of cycloartocarpin. Manipulation of the liquor by thin-layer and gravitation column chromatographic techniques proved to be a simple method to isolate cycloartocarpin. The flavonoid structure, cycloartocarpin, was treated with structure elucidation 1- and 2-dimensional NMR and infrared spectrometry. Students then compared the structure of the cycloartocarpin obtained with that of other flavonoids from the genus *Artocarpus* and other plant genera. From these comparisons, they were able to formulate structural characteristics of the flavonoid compounds of the genus *Artocarpus*. The laboratory method, accomplished in four, 3 h laboratory sessions, allowed students to readily understand the process of isolating flavonoids and their peculiarities.

KEYWORDS: Upper-Division Undergraduate, Organic Chemistry, Laboratory Instruction, Hands-On Learning/Manipulatives, Problem Solving/Decision Making, Natural Products

INTRODUCTION

The purpose of this laboratory was first to provide students with the essential skills required to perform the extraction, fractionation, purification, and structural elucidation of a flavonoid. Second, it provided them with knowledge of how to formulate the structural characteristics of secondary metabolites from a plant genus, in this case, the flavonoids from the genus *Artocarpus*. Many of the activities in this laboratory may be appropriate for other courses, such as ones involving the structure elucidation of organic compounds, organic chemistry, and chemotaxonomy.

Secondary metabolites, including flavonoids from the genus *Artocarpus* are isolated by extraction, fractionation, purification, and structural elucidation.^{1,2} Furthermore, students can formulate the structural characteristics of the flavonoids from the genus *Artocarpus* by comparing their structure with those of other flavonoids from the genus *Artocarpus* and from other plant genera. These activities were carried out in this natural product chemistry laboratory. The natural product chemistry laboratory had been previously described in several studies.^{2–10}

Several studies on the genus *Artocarpus* have shown that plants from this genus contain phenol derivative compounds, especially in flavonoid groups that display certain peculiarities.^{11,12} The peculiarities involve the prenylated group in position C-3 and ring B oxygenated at position C-4' or C-2', C-4' or C-2', C-4', C-5'. In addition, prenylation can also occur at positions C-6 and C-8; such a pattern is not found in other genera (Figure 1). The the flavonoid structure in the genus *Artocarpus* is unique, and the flavonoid displays

interesting biological activities, such as antimalarial,¹³ anticancer,¹⁴ and cytotoxic¹⁵ ones.

The structure of flavonoids from other plant genera follows the shikimic acid and acetic malonate biogenesis pathways. Comparison between the flavonoid prenylation and oxidation of the genus *Artocarpus* and other plant genera can be seen in Figure 1.

METHODOLOGY

This study uses quasi-experimental research with one group pretest–posttest design. Participants in this study consisted of 32 third-year students (teacher preparation) from chemistry education department at one of the state universities in West Nusa Tenggara, Indonesia. These laboratory activities took place during the second semester of the 2018–2019 academic years. Participants were divided into eight groups with four participants per group. All of the groups worked on the same plant sample, roots of *Artocarpus altilis* (commonly known as breadfruit).

A questionnaire containing 15 questions was used to assess the student knowledge of isolation processes and the ability of students to analyze the uniqueness of *Artocarpus* flavonoids.

Received: February 26, 2020

Revised: August 24, 2020

Published: September 18, 2020




ACS Publications

© 2020 American Chemical Society and
Division of Chemical Education, Inc.

4133

<https://dx.doi.org/10.1021/acs.jchemeduc.0c00199>
J. Chem. Educ. 2020, 97, 4133–4136



A. Prenylation and oxidation pattern of genus *Artocarpus* flavonoids.
B. Prenylation and oxidation patterns of flavonoid in other genera.

Figure 1. Possibility of prenylation and possibility of oxidation pattern of flavonoids.

The validity of the test was determined by using content validity experts. Qualitative analysis was done by interpretive descriptive analysis. Quantitative analysis of data was done by calculating the percentage of normalized gain scores (%) using the formula in eq 1.¹⁶

$$\%g = \frac{S_{\text{post}} - S_{\text{pre}}}{S_{\text{max}} - S_{\text{pre}}} 100 \quad (1)$$

where S_{post} and S_{pre} are the post-test and pretest scores, respectively, and S_{max} is the maximum possible score. Values of %g were then characterized as high for %g > 70%, medium for 30% ≤ %g ≤ 70%, and poor for %g < 30%.

Laboratory reports were also used at the end of the laboratory activity that was arranged in groups. Student surveys were used to demonstrate student engagement with the experiment.

EXPERIMENTAL PROCEDURE

The experiment, conducted over four, 3 h laboratory sessions, was useful for the third-year undergraduate students. Throughout the experiment, they gained first-hand experience of various techniques involved in the extraction, isolation, and purification of cycloartocarpin, a flavonoid compound, from *Artocarpus altilis* roots. The students were able to formulate the unique structure of the genus *Artocarpus* flavonoid by comparing the structure of cycloartocarpin obtained with that of other flavonoids from the genus *Artocarpus* and other plant genera. Procedures of the student activity are described below; detailed procedures can be found in the Supporting Information.

Isolation of Cycloartocarpin from *Artocarpus altilis* Roots

The cycloartocarpin isolation procedure conducted in this laboratory activity differs from previous descriptions in the literature.^{17,18} Students followed these steps for the first week:

- The groups prepared air-dried root stems of *A. altilis* by macerating these using methanol.
- The methanol extract was filtered using cellulose filter paper.
- The solvent was evaporated by a rotary evaporator to dryness to obtain a crude extract.

During the second week of the experiment, students undertook these steps:

- The crude extract was fractionated using gravity column chromatography.
- A single compound was characterized by the presence of a single spot in a TLC chromatogram.

This isolation procedure was also carried out by students in the previous semester, the second semester of 2017–2018. The isolation procedures were used by the students two times. Further details are provided in the student handout of the Supporting Information.

Structure Elucidation Studies

In the third week of the experiment, students performed IR and detailed NMR spectroscopic analyses and confirmed that the substances isolated were cycloartocarpin. The IR (KBr) spectrum of V_{max} (cm^{-1}) showed the presence of a conjugated carbonyl group ($\text{C}=\text{O}$), typical for flavone at an uptake of 1651 cm^{-1} and $\text{C}=\text{C}$ aromatic groups at an uptake of $1620\text{--}1450 \text{ cm}^{-1}$. The IR spectrum also showed the presence of absorption for the hydroxyl (OH) group at $V_{\text{max}} = 3390 \text{ cm}^{-1}$ and aliphatic C-H at $V_{\text{max}} = 2924\text{--}2854 \text{ cm}^{-1}$. The IR data of a single compound showed that all functional groups were found in the cycloartocarpin compounds, as seen in Figure 2.




Figure 2. Structure of cycloartocarpin.

Further determination of the structure of a single compound was obtained from the ^{13}C NMR, which indicated 26 signals of carbon atoms. There was a chemical shift at δ_{c} 178.4 ppm, which is a typical sign of carbonyl ($\text{C}=\text{O}$), and a sign of carbon oxyaryl at δ_{c} (139–162) ppm. In addition, there were also signs of aromatic carbon at δ_{c} 104–125 ppm, including four methyl carbons and carbon methoxy. ^1H NMR (500 MHz, CDCl_3) showed a typical sign in the region of 13.42 ppm for a chelated $-\text{OH}$, which was evidence of flavonoid derivative. The pirano group γ,γ -dimethyl allyl was indicated by the presence of proton signals in chemical shifts at 1.69 (3H, s), 1.96 ppm (3H, s); 5.43 ppm (1H, d, $J = 9.4 \text{ Hz}$); 6.25 ppm (1H, d, $J = 9.4 \text{ Hz}$). Proton signals for ring B were shown at chemical shifts of 6.42 ppm (1H, d, $J = 2.3 \text{ Hz}$); 6.52 (1H, dd, $J = 2.3$; 8.5 Hz); 7.66 (1H, d, $J = 8.5 \text{ Hz}$). These signals indicate the existence of the ABX system on ring B, which was substituted at positions 2' and 4'. The signal for the proton in the methoxy group was in the 3.93 (3H, s, OCH_3) chemical shift. HSQC showed five C-oxyaryl signals, six C-quaternary signals, and nine signals, including four methyl, methoxyl, and

carbonyl. HMBC showed that ring A was substituted by a prenyl group substituted at C-6 in the presence of proton correlation of chelated OH with carbon from the prenyl group. More details can be found in the [Instructor Notes of the Supporting Information](#).

Formulation of the Uniqueness of the Flavonoid Structure of the Genus *Artocarpus*

In the fourth week, students conducted a literature study through the internet and university library in order to find the structure of flavonoid compounds from the genus *Artocarpus* and other plant genera. Furthermore, they compared the patterns of prenylation and oxidation of the structure of isolated compounds with the structure of the flavonoid compounds obtained from the literature. From this activity, the students were able to deduce the unique flavonoid structure of the genus *Artocarpus*. The complete assignment is provided in the [Supporting Information](#). Following completion of the experiment, each group of students submitted a detailed report.

Students Result

Of the eight groups who participated in this laboratory, only three groups succeeded in getting a single compound (based on the purity of TLC chromatograms compared with standard cycloartocarpin). However, all groups were still given the IR and NMR spectra provided by the instructor for interpretation by all groups. From the reports collected, all groups could determine all functional groups found in the cycloartocarpin compounds based on IR data. However, they had difficulty interpreting the NMR spectrum. Two groups could show an almost-correct interpretation of the NMR data. The other groups demonstrated poor NMR interpretation abilities. Six groups succeeded in formulating the specificity of *Artocarpus* flavonoids in the conclusions of their laboratory report. In the quantitative analysis of pretest and post-test scores, the average of students' n-gain percentage was 63% (medium category). See the student gain score spreadsheet in the [Supporting Information](#).

HAZARDS

This experiment uses some potentially hazardous substances and flammable solvents, including methanol, dichloromethane, *n*-pentane, and petroleum ether (highly flammable). The use of flammable solvents should not involve direct flames. In addition, dichloromethane has limited evidence of a carcinogenic effect.⁷ The use of carcinogenic substances must be carried out in a fume hood. Ultraviolet (UV) radiation can cause severe damage to the eyes, so one should not look directly into the light source. Students should use personal protective equipment, including gloves and safety goggles.

SUMMARY

This experiment, with its straightforward method, allows students to isolate and characterize cycloartocarpin, an *Artocarpus altilis* flavonoid compound, and understand its uniqueness over four, 3 h laboratory sessions. Students work in small groups. If one group fails to isolate the pure material, that group can join a successful group for subsequent characterization and analysis. Isolated compounds can be replaced and come from different genera.

ASSOCIATED CONTENT

Supporting Information

The Supporting Information is available at <https://pubs.acs.org/doi/10.1021/acs.jchemeduc.0c00199>.

Instructor notes (PDF, DOCX)

Student handout (PDF, DOCX)

Student gain score spreadsheet (XLSX)

AUTHOR INFORMATION

Corresponding Author

Aliefman Hakim — Study Program of Chemistry Education,
University of Mataram, Mataram-Lombok 83125, Indonesia;
orcid.org/0000-0002-5670-853X;
Email: Aliefmanhakim27@gmail.com

Authors

A. Wahab Jufri — Study Program of Biology Education,
University of Mataram, Mataram-Lombok 83125, Indonesia
Jamaluddin — Study Program of Biology Education, University of
Mataram, Mataram-Lombok 83125, Indonesia
Supriadi — Study Program of Chemistry Education, University of
Mataram, Mataram-Lombok 83125, Indonesia
Putri Ayu Mutmainnah — Study Program of Chemistry
Education, University of Mataram, Mataram-Lombok 83125,
Indonesia

Complete contact information is available at:
<https://pubs.acs.org/doi/10.1021/acs.jchemeduc.0c00199>

Notes

The authors declare no competing financial interest.

ACKNOWLEDGMENTS

We wish to thank the Ministry of Research, Technology and Higher Education, Indonesia, for funding the research.

REFERENCES

- (1) Nguyen, M. T. T.; Le, T. H.; Nguyen, H. X.; Dang, P. H.; Do, T. N. V.; Abe, M.; Takagi, R.; Nguyen, N. T. Artocarmins G–M, Prenylated 4-Chromenones from the Steams of *Artocarpus rigida* and Their Tyrosinase Inhibitory Activities. *J. Nat. Prod.* **2017**, *80* (12), 3172–3178.
- (2) Hakim, A.; Liliarsari; Kadarohman, A.; Syah, Y. M. Making a Natural Product Chemistry Course Meaningful with a Mini Project Laboratory. *J. Chem. Educ.* **2016**, *93* (1), 193–196.
- (3) Griffin, R. W. Natural Products. An Independent Study Project. *J. Chem. Educ.* **1974**, *51* (9), 601–602.
- (4) Cannon, J.; Li, D.; Wood, S. G.; Owen, N. L.; Gromova, A.; Lutsky, V. Investigation of Secondary Metabolites in Plants—A General Protocol for Undergraduate Research in Natural Products. *J. Chem. Educ.* **2001**, *78* (9), 1234–1237.
- (5) Doyle, A. M.; Reilly, J.; Murphy, N.; Kavanagh, P. V.; O'Brien, J. E.; Walsh, M. S.; Walsh, J. J. Nature's Sedative: Isolation and Structural Elucidation of Valtrate from *Centranthus ruber*. *J. Chem. Educ.* **2004**, *81* (10), 1486–1487.
- (6) Kavanagh, P. V.; Douglas, R. H.; Muldowney, C. A.; Mohamed, R.; Keohane, F.; Shanahan, C.; Walsh, J. J. Detection and Quantification of Valerenic Acid in Commercially Available Valerian Products. *J. Chem. Educ.* **2007**, *84* (5), 829–831.
- (7) Halpin, C. M.; Reilly, C.; Walsh, J. J. Nature's Anti-Alzheimer's Drug: Isolation and Structure Elucidation of Galantamine from *Leucojum aestivum*. *J. Chem. Educ.* **2010**, *87* (11), 1242–1243.

- (8) Walsh, E. L.; Ashe, S.; Walsh, J. J. Nature's Migraine Treatment: Isolation and Structure Elucidation of Parthenolide from *Tanacetum parthenium*. *J. Chem. Educ.* **2012**, *89* (1), 134–137.
- (9) Nazri, M. M.; Samat, F. D.; Kavanagh, P. V.; Walsh, J. J. Nature's Cholesterol-Lowering Drug: Isolation and Structure Elucidation of Lovastatin from Red Yeast Rice-Containing Dietary Supplements. *J. Chem. Educ.* **2012**, *89* (1), 138–140.
- (10) Carroll, A. M.; Kavanagh, D. J.; McGovern, F. P.; Reilly, J. W.; Walsh, J. J. Nature's Chiral Catalyst and Anti-Malarial Agent: Isolation and Structure Elucidation of Cinchonine and Quinine from *Cinchona calisaya*. *J. Chem. Educ.* **2012**, *89* (12), 1578–1581.
- (11) Hakim, A. A Prenylated Flavone from The Heartwood of *Artocarpus scorteichinii* King (Moraceae). *Indones. J. Chem.* **2009**, *9* (1), 146–150.
- (12) Nomura, T.; Hano, S.; Aida, M. Isoprenoid Substituted Flavanoid from *Artocarpus* Plants (Moraceae). *Heterocycles* **1998**, *47* (2), 1179–1205.
- (13) Boonlaksiri, C.; Oonant, W.; Kongsaree, P.; Kittakoo, P.; Tanticharoen, M.; Thebtaranonth, Y. An Antimalarial Stilbene from *Artocarpus integer*. *Phytochemistry* **2000**, *54* (4), 415–417.
- (14) Ren, Y.; de Blanco, E. J. C.; Fuchs, J. R.; Soejarto, D. D.; Burdette, J. E.; Swanson, S. M.; Kinghorn, A. D. Potential Anticancer Agents Characterized from Selected Tropical Plants. *J. Nat. Prod.* **2019**, *82* (3), 657–679.
- (15) Musthapa, I.; Hakim, E. H.; Syah, Y. M.; Juliawaty, L. D. Cytotoxic Activities of Prenylated Flavonoids from *Artocarpus heterophyllus*. *ARNP J. Eng. Appl. Sci.* **2016**, *11* (16), 9754–9758.
- (16) Hake, R. Interactive Engagement vs. Traditional Methods: A Six Thousand-Student Survey of Mechanics Test Data for Introductory Physics Courses. *Am. J. Phys.* **1998**, *66* (1), 64–74.
- (17) Boonphong, S.; Baramee, A.; Kittakoo, P.; Puangsombat, P. Antitubercular and Antiplasmodial Prenylated Flavones from the Roots of *Artocarpus altiss.* *Chiang Mai J. Sci.* **2007**, *34* (3), 339–344.
- (18) Nair, P. M.; Rama Rao, A. V.; Venkataraman, K. Cyclo-artocarpin. *Tetrahedron Lett.* **1964**, *5* (2), 125–128.

Understanding the Uniqueness of Artocarpus Flavonoids: Isolation and Structure Elucidation of Cycloartocarpin from the Roots of Artocarpus altilis

ORIGINALITY REPORT

19%

SIMILARITY INDEX

12%

INTERNET SOURCES

14%

PUBLICATIONS

5%

STUDENT PAPERS

PRIMARY SOURCES

1	digilib.unimed.ac.id Internet Source	2%
2	mafiadoc.com Internet Source	1%
3	Paige J. Monsen, Frederick A. Luzzio. "Isolation and Derivatization of Sucralose from an Artificial Sweetener to Provide a Hands-On Laboratory Experiment Emphasizing Synthesis and Purification", Journal of Chemical Education, 2019 Publication	1%
4	iopscience.iop.org Internet Source	1%
5	scholar.google.com Internet Source	1%
6	Aliefman Hakim, A. Wahab Jufri, Jamaluddin. "Innovative natural product chemistry	1%

laboratory: Isolation of artelastin from Artocarpus scortechinii", AIP Publishing, 2019

Publication

7	www.scirp.org Internet Source	1 %
8	pubs.acs.org Internet Source	1 %
9	Maisarah Mohd Nazri, Farah D. Samat, Pierce V. Kavanagh, John J. Walsh. "Nature's Cholesterol-Lowering Drug: Isolation and Structure Elucidation of Lovastatin from Red Yeast Rice-Containing Dietary Supplements", Journal of Chemical Education, 2012 Publication	1 %
10	patentscope.wipo.int Internet Source	1 %
11	Submitted to Kennesaw State University Student Paper	1 %
12	www.freepatentsonline.com Internet Source	1 %
13	N Sirajudin, J Suratno, Pamuti. "Developing creativity through STEM education", Journal of Physics: Conference Series, 2021 Publication	1 %
14	link.springer.com Internet Source	1 %

15	Graham R. Parslow ‡. "Websites of note", Biochemistry and Molecular Biology Education, 01/2011 <small>Publication</small>	1 %
----	--	-----

16	Pablo Martín-Ramos, Maria Susano, Francisco P. S. C. Gil, Pedro S. Pereira da Silva, Jesús Martín-Gil, Manuela Ramos Silva. "Facile Synthesis of Three Kobolds: Introducing Students to the Structure of Pigments and Their Characterization", Journal of Chemical Education, 2018 <small>Publication</small>	<1 %
----	--	------

17	Remko T. Winter, Hugo L. van Beek, Marco W. Fraaije. "The Nose Knows: Biotechnological Production of Vanillin", Journal of Chemical Education, 2012 <small>Publication</small>	<1 %
----	--	------

18	The Flavonoids, 1975. <small>Publication</small>	<1 %
----	---	------

19	pendidikankimia.walisongo.ac.id <small>Internet Source</small>	<1 %
----	---	------

20	www.science.gov <small>Internet Source</small>	<1 %
----	---	------

21	Ewa Iwanek (née Wilczkowska), Marek Gliński. "Application of Thermal Analysis in Determining Properties of Herbaceous	<1 %
----	---	------

22

Lisa T. Alty, Marcia B. France, Isaac G. Alty,
Christine A. Saber, Donna M. Smith.

"Synthesis of 1,1-Diphenylethylene (DPE): The
Marriage of a Grignard Reaction and a
Column Chromatography Experiment",
Journal of Chemical Education, 2016

Publication

<1 %

23

Submitted to Macquarie University

Student Paper

<1 %

24

Maeve Egan, Éilis Margaret Connors, Zeeshan
Anwar, John J. Walsh. "Nature's Treatment for
Irritable Bowel Syndrome: Studies on the
Isolation of (–)-Menthol from Peppermint Oil
and Its Conversion to (–)-Menthyl Acetate",
Journal of Chemical Education, 2015

Publication

<1 %

25

Mutai, C.. "Cytotoxic lupane-type triterpenoids
from *Acacia mellifera*", *Phytochemistry*,
200404

Publication

<1 %

26

uwm.edu

Internet Source

<1 %

27

worldwidescience.org

Internet Source

<1 %

Exclude quotes On

Exclude bibliography On

Exclude matches < 5 words

Understanding the Uniqueness of Artocarpus Flavonoids: Isolation and Structure Elucidation of Cycloartocarpin from the Roots of Artocarpus altilis

GRADEMARK REPORT

FINAL GRADE

/0

GENERAL COMMENTS

Instructor

PAGE 1

PAGE 2

PAGE 3

PAGE 4