

## Document Viewer

## Turnitin Originality Report

Processed on: 25-Jun-2022 06:49 WIB

ID: 1862507568

Word Count: 4296

Submitted: 1

Seed Coating with Organomineral Fertilizer,  
a... By Joko Priyono

Similarity Index

24%

## Similarity by Source

|                   |     |
|-------------------|-----|
| Internet Sources: | 22% |
| Publications:     | 13% |
| Student Papers:   | 10% |

[exclude quoted](#) [exclude bibliography](#) [exclude small matches](#)  
[print](#) [refresh](#) [download](#)

mode: quickview (classic) report

Change mode

4% match (Internet from 26-Oct-2019)

<https://journalgazett.com/journals/asian-journal-of-advances-in-agricultural-research/>

1% match (Internet from 11-Jun-2022)

<https://journalarrb.com/index.php/ARRB/article/download/30142/56577>

1% match (Internet from 04-Oct-2019)

<https://pinpdf.com/557184fcd883e8f0aaa38f556db6cca745730.html>

1% match (publications)

[K. N. N. Ezike, E. N. Okeowata, V. A. Chukwu, Ogonna O. Osuafor. "Funding of Agricultural Public Extension Service in a Deregulated Economy: A Study of Ebonyi State, Nigeria", Asian Research Journal of Agriculture, 2020](#)

1% match (Internet from 09-Dec-2017)

<http://sciencedomain.org>

1% match (publications)

[Teshome Gizachew Belay, Tamirat Tefera Negash. "Perceptions of Flower Farm Workers on the Health Effects of Chemicals Exposures around Bahir Dar Town, Ethiopia", Asian Research Journal of Agriculture, 2020](#)

1% match (student papers from 05-Nov-2020)

[Submitted to University of Western Australia on 2020-11-05](#)

|   |   |
|---|---|
| 1% match (Internet from 30-Oct-2020)<br><a href="https://arpgweb.com/pdf-files/jac6(5)68-72.pdf">https://arpgweb.com/pdf-files/jac6(5)68-72.pdf</a>   | ✕ |
| 1% match (Internet from 13-Apr-2022)<br><a href="http://journal.ummat.ac.id">http://journal.ummat.ac.id</a>   | ✕ |
| 1% match (Internet from 20-Mar-2022)<br><a href="http://repository.maranatha.edu">http://repository.maranatha.edu</a>   | ✕ |
| 1% match (student papers from 07-Jul-2020)<br><a href="#">Submitted to Wageningen University on 2020-07-07</a>  | ✕ |
| 1% match (publications)<br><a href="#">Benita Silva, Remigio Paradelo, Nuria Vázquez, Eduardo García-Rodeja, María Teresa Barral. "Effect of the addition of granitic powder to an acidic soil from Galicia (NW Spain) in comparison with lime", Environmental Earth Sciences, 2012</a> | ✕ |
| 1% match (publications)<br><a href="#">Tobias Pfeiffer, Astrid von Galen, Petra Zink, Sebastian Hübner et al. "Selection of bacteria and fungi for control of soilborne seedling diseases of maize", Journal of Plant Diseases and Protection, 2021</a>                                 | ✕ |
| 1% match (Internet from 11-Oct-2020)<br><a href="https://res.mdpi.com/d_attachment/land/land-09-00198/article_deploy/land-09-00198-v2.pdf">https://res.mdpi.com/d_attachment/land/land-09-00198/article_deploy/land-09-00198-v2.pdf</a>   | ✕ |
| 1% match (Internet from 01-Jun-2022)<br><a href="https://www.ikpress.org/index.php/PCBMB/article/view/7207">https://www.ikpress.org/index.php/PCBMB/article/view/7207</a>   | ✕ |
| 1% match (Internet from 20-Dec-2021)<br><a href="https://3lata.pl/12/Oct/2009-9203.html">https://3lata.pl/12/Oct/2009-9203.html</a>   | ✕ |
| 1% match (Internet from 20-Oct-2020)<br><a href="https://www.intechopen.com/profiles/15636/ruth-wanyera">https://www.intechopen.com/profiles/15636/ruth-wanyera</a>   | ✕ |
| <1% match (Internet from 27-Sep-2020)<br><a href="https://download.atlantis-press.com/proceedings/piceeba-18/25902656">https://download.atlantis-press.com/proceedings/piceeba-18/25902656</a>  | ✕ |
| <1% match (Internet from 05-Aug-2021)<br><a href="http://journalrepository.org">http://journalrepository.org</a>  | ✕ |

|   |   |
|---|---|
| <1% match (Internet from 03-Aug-2021)<br><a href="http://journalrepository.org">http://journalrepository.org</a>  | ✕ |
| <1% match (Internet from 16-Feb-2022)<br><a href="http://repository.unida.ac.id">http://repository.unida.ac.id</a>  | ✕ |
| <1% match (Internet from 28-Jan-2022)<br><a href="http://cyxb.magtech.com.cn">http://cyxb.magtech.com.cn</a>  | ✕ |
| <1% match (Internet from 22-Nov-2016)<br><a href="http://oar.icrisat.org">http://oar.icrisat.org</a>  | ✕ |
| <1% match (student papers from 27-May-2022)<br><a href="#">Submitted to The Federal Polytechnic Ede on 2022-05-27</a>   | ✕ |
| <1% match (student papers from 15-Feb-2016)<br><a href="#">Submitted to Universitas Brawijaya on 2016-02-15</a>   | ✕ |
| <1% match (Internet from 14-Nov-2021)<br><a href="https://journals.ansfoundation.org/index.php/jans/article/view/641">https://journals.ansfoundation.org/index.php/jans/article/view/641</a>  | ✕ |
| <1% match (Internet from 03-Jan-2022)<br><a href="https://www.degruyter.com/document/doi/10.1515/revce-2020-0044/html">https://www.degruyter.com/document/doi/10.1515/revce-2020-0044/html</a>  | ✕ |
| <1% match (Internet from 04-Jun-2022)<br><a href="https://dergipark.Org.Tr/en/pub/akuiibfd/indexes">https://dergipark.Org.Tr/en/pub/akuiibfd/indexes</a>  | ✕ |
| <1% match (Internet from 06-Oct-2021)<br><a href="https://www.grocentre.is/static/gro/publication/719/document/abdul-rahman2019.pdf">https://www.grocentre.is/static/gro/publication/719/document/abdul-rahman2019.pdf</a>                                      | ✕ |
| <1% match (publications)<br><a href="#">M. Shahinul, M. Rahman, M. M. R. Salim, B. Ahmed, M. H. Rashid. "Effect of Planting Method and Seeding Date on Leaf and Seed Production of Lettuce", Asian Journal of Agricultural and Horticultural Research, 2019</a> | ✕ |
| <1% match (Internet from 26-Feb-2022)<br><a href="https://cis01.ucv.ro/revistadestiintepolitice/linkuriBDI.php">https://cis01.ucv.ro/revistadestiintepolitice/linkuriBDI.php</a>  | ✕ |
| <1% match (Internet from 12-Oct-2018)<br><a href="http://www.sciencedomain.org">http://www.sciencedomain.org</a>  | ✕ |

|  |   |
|--|---|
| <1% match (publications)<br><a href="#">Md. Nur Alam Miah, Md. Ramiz Uddin Miah, Md. Zinnatul Alam, Md. Mofazzal Hossain. "Influence of Weather on Sugarcane Mealybugs and Their Natural Enemies", Asian Research Journal of Agriculture, 2019</a>   | ✕ |
| <1% match (Internet from 10-Dec-2016)<br><a href="http://unhas.ac.id">http://unhas.ac.id</a>   | ✕ |
| <1% match (Internet from 12-Aug-2016)<br><a href="https://www.linknovate.com/affiliation/indian-institute-of-vegetable-research-106767/all/?query=lime">https://www.linknovate.com/affiliation/indian-institute-of-vegetable-research-106767/all/?query=lime</a>   | ✕ |
| <1% match (Internet from 07-Oct-2021)<br><a href="https://www.akademisains.gov.my/asmsj/?mdocs-file=4999">https://www.akademisains.gov.my/asmsj/?mdocs-file=4999</a>   | ✕ |
| <1% match (Internet from 21-Apr-2020)<br><a href="https://www.neliti.com/publications/132407/adding-organic-matter-enhanced-the-effectiveness-of-silicate-rock-fertilizer-for">https://www.neliti.com/publications/132407/adding-organic-matter-enhanced-the-effectiveness-of-silicate-rock-fertilizer-for</a>   | ✕ |
| <1% match (publications)<br><a href="#">Yohannes Gebremichael, Gebremedhen Gebretsadikan. "Effect of Mineral Nitrogen and Phosphorus Fertilizer Rates on Marketable Yield and Economic Return of Tomato (Lycopersicon esculentum Mil) at Northwestern Zone of Tigray", Asian Research Journal of Agriculture, 2019</a>   | ✕ |
| <1% match (Internet from 24-Mar-2022)<br><a href="https://jabonline.a2hosted.com/jab_php/abstract.php?article_id=664">https://jabonline.a2hosted.com/jab_php/abstract.php?article_id=664</a>   | ✕ |
| <1% match (publications)<br><a href="#">S K Sebayang, M J Dibley, P Kelly, A V Shankar, A H Shankar. "Modifying effect of maternal nutritional status on the impact of maternal multiple micronutrient supplementation on birthweight in Indonesia", European Journal of Clinical Nutrition, 2011</a>  | ✕ |
| <p><a href="#">Asian Research Journal of Agriculture 12(1): 12-17, 2020; Article no.ARJA.52771 ISSN: 2456-561X Seed Coating with Organomineral Fertilizer, an Alternative Method to Improve the Efficiency of Farming</a> Joko Priyono1* and Anak Agung Ketut Sudharmawan2 1Department of Soil Science, <a href="#">University of Mataram, Jalan Pendidikan 37, Mataram, Lombok Island, Indonesia</a>. 2Department of Agroecotechnology, <a href="#">University of Mataram, Jalan Pendidikan 37, Mataram, Lombok Island, Indonesia</a>. <a href="#">Authors' contributions</a> This work was carried out in collaboration between both <a href="#">authors</a>. <a href="#">Author</a> JP designed <a href="#">the</a> study, technical aspects <a href="#">of</a> seed coating, constructs <a href="#">the</a> outline and wrote <a href="#">the</a> first draft of <a href="#">the</a> manuscripts. <a href="#">Author</a> AAKS <a href="#">performed the statistical analysis, managed the literature searches</a> for seed treatments. <a href="#">Both authors read and approved the final manuscript</a>. <a href="#">Article Information DOI: 10.9734/ARJA/2020/v12i130073 Editor(s): (1) Dr. Rusu Teodor, Department of Technical and Soil Sciences,</a></p> |   |

University of Agricultural Sciences and Veterinary Medicine Cluj-Napoca, Romania. Reviewers: (1) [Maria Lúcia Pereira da Silva, University of São Paulo, Brazil](#). (2) Mohammed, Dahiru Toungos, Adamawa State University, Nigeria. (3) [Ajoy Kumer, European University of Bangladesh, Bangladesh](#). [Complete Peer review History: http://www.sdiarticle4.com/review-history/52771](#) Received 12 September 2019 Original Research Article Accepted 18 November 2019 Published 06 January 2020 ABSTRACT Seed and fertilizer are two important farming inputs, which are commonly available and used separately. Combining both materials into a unit of fertilizer-coated seed may improve farming efficiency. However, the appropriate seed coating method must be found out, and this research was the first effort of finding the method. [A glasshouse experiment was carried out to identify the growth and yield of the coated seeds of rice and groundnut with organomineral fertilizer in three different sizes, i.e., small \(SS\), medium \(MS\), and big sizes \(BS\)](#). Four sets of experiments were prepared, two of those were for testing two varieties of rice and the others were for testing two varieties of groundnut. Each [experiment was laid out in a complete randomized design; the treatment was the size of coated seeds \(SS, MS, BS, and a control - uncoated seeds\)](#) in triplicates. Results reveal that the seed coating delayed the germination of rice seeds for 2 – 3 days and groundnut seeds for 7 –

\*Corresponding author: E-mail: [joko\\_priyono@unram.ac.id](mailto:joko_priyono@unram.ac.id); 16 days, suppressed the growth and yield of rice but improved the growth and yield of groundnut. The highest yield of groundnut was the grown groundnut from the small and medium sizes of coated seeds (weight ratios of 1:4 and 1:9). The reduces of growth and yield of rice were most probably due to the direct contact of the high concentration of nutrients, especially nitrogen, with the seeds. In conclusion, the seed coating with organomineral fertilizer was a potentially developed method to improve farming efficiency. Further efforts were needed to fix the composition of organomineral fertilizer, especially the type N substances used and the steps of applying the materials onto the seeds. Keywords: Organomineral fertilizer; coating materials; seed coating; silicate rock fertilizer. 1. INTRODUCTION Seed and fertilizer are two vital inputs in the farming of crops grown from seed, such as [rice \(\*Oryza sativa\* L.\)](#) and [groundnut \(\*Arachis hypogaea\* L.\)](#). The current issue of farming technology relating to the seeds is seed coating, and of that for fertilizer is the utilization of the natural sources of plant nutrients, including silicate rocks. Combining those farming inputs into a unit material, which is coating the seeds with the fertilizer, may be an appropriate method to improve the productivity and profitability of the farming. Seed coating is an important technology that has been applied by seed industries for decades. The industries use various coating materials, and the most common one is pesticides [1]. The main aims of using pesticides are [to suppress, control, or repel the soil-borne pathogens, insects, and other pests attacking seeds, seedlings, or plants](#) [2,3]. However, exposing pesticides into farming land may potentially develop pest resistance and create various negative-ecological effects [4]. Thus, it is a challenge for the seed industries to find out and use the more environmentally sound coating materials. Fertilizer is another important farming input, used to fulfil the optimum requirement of crops for essential nutrients. The appropriate fertilizer application, referring to the ideal concept of proportional fertilization, however, is often difficult to be applied on the farm level, especially by small farmers in developing countries, due to technical or/and economic constraints. The high variability of soil properties, for example, may cause the difficulties to define the appropriate rate of fertilizer application; and the lack of farming capital often limits the small farmers to access the appropriate quantities of synthetic fertilizers. Responding to the cases, many researchers [5,6,7,8] proposed the utilization of natural sources of plant-nutrients, e.g., silicate rocks, as cheap and environmentally sound fertilizer [11]. However, the use of silicate rock fertilizers (SRFs) was limited in farming practices due to the very slow release of nutrients

from the SRFs. Consequently, the farming required very high doses (> 20 t.ha-) of SRFs [7]. However, by employing ball milling in producing SRFs [9,10], or/and mixing SRFs with organic matter (termed as organomineral fertilizer), significantly improved the effectiveness of SRFs [11]. The limitations of the seed coating technology, especially in selecting the type of coating materials, and fertilizer application described above inspired us to propose combining the seed with organomineral fertilizer into the organomineral-coated seeds as an alternative solution for the farming constraints. The organomineral fertilizer consisted mainly of SRFs, rock phosphate (RP), and organic matters (OM), and those materials were coated onto the seeds, and the coating materials were functioned as the major nutrient source sufficient for fulfilling the optimum requirement of essential nutrients for the plant grown from the coated seeds without additional fertilizers. Besides reducing the farming costs (for fertilizers and labor) with less negative-environmental effects, planting the coated seeds may also improve soil quality due to the residual effects of the coating materials (organomineral fertilizer) remaining in the soil. This research was the first attempt to evaluate if the proposed farming method described above was realistic. The [main objective of this research was to identify the effects of seed coating with organomineral fertilizer on the growth and yield components of rice \(\*Oryza sativa\* L\) and groundnut \(\*Arachis hypogaea\* L\).](#)

## 2. MATERIALS AND METHODS

### 2.1 Seeds and Coating Materials

The seeds of two varieties of rice, i.e., Impari 32 and Bunda Sri Madrin (BSM), and two varieties of groundnut, i.e., Lombok and Kelinci, were prepared. The coating materials for rice seeds were SRF (87.5% w/w), PR (4% w/w), urea (4% w/w), OM (2% w/w), and liquid-foliar fertilizer (trademarked as Orrin) (2.5% v/w). The SRF, RP, and OM were mixed, while urea was dissolved into Orrin that was functioned as the binding substance. The coating materials for groundnut seeds were the same as those for rice seeds, but the urea was substituted with CaCO<sub>3</sub> (4% w/w) added to the mixture of SRFs + RP + OM, and the binding substance for the coated seeds was only Orrin. The SRF was the finely ground basaltic rocks from Rinjani Mt. in Lombok Island – Indonesia. The rocks were mechanically broken to  $\varnothing < 1$  cm and then ball-milled for 40 minutes producing the SRF-powder with the median diameter (D<sub>50</sub>) < 5.4  $\mu$ m. The chemical composition of the SRF-powder was 54.22% SiO<sub>2</sub>, 25.01% Al<sub>2</sub>O<sub>3</sub>, 5.00% CaO, 1.90% 4.44% K<sub>2</sub>O, 2.39% Na<sub>2</sub>O, 6.47% FeO, 0.16% MnO, 0.01% ZnO, 0.31 % CuO, and < 0.10% others. The RP ( $\varnothing < 0.5$  cm) was ball-milled for 20 minutes producing RP-powder with the median diameter (D<sub>50</sub>) < 22  $\mu$ m, and it contained a total of 16.8% P<sub>2</sub>O<sub>5</sub>. The OM was the composted cattle wastes, screened to pass the 1-mm stainless steel screen. Urea and CaCO<sub>3</sub> respectively contained 44.2% N and 38% CaO. The liquid-foliar fertilizer (Orrin) contained of 4.04% N, 3.22% P<sub>2</sub>O<sub>5</sub>, 3.36% K<sub>2</sub>O, 0.32% Ca, 0.40% Mg, 0.12% S, and 40, 122, 260, 10, 3, 0.1, and 1.2 mg.L<sup>-1</sup> respectively of Fe, Mn, Zn, Cu, B, Co, and Mo.

### 2.2 Seed Coating Process

The seeds were coated using a rotating drum (350-L plastic drum), rotated with a 0.5-HP power machine at the speed of about 30 rpm. For coating the seeds of rice, 250-g seeds were coated with 2,500 g of the mixture of coating materials described above. In the first step, half of the mixed coating materials (SFR + RP + OM) were applied step-by-step onto the seed surface in the rotating drum, followed by spraying fresh water onto the seeds, forming the capsule-like coated seeds. Next, the other half of the coating materials were applied in the same way but using the binding substance of the urea + Orrin solution. The coated seeds were dried with a hairdryer (< 40°C) in the rotating drum to avoid cracking of the coated seeds. Finally, the coated seeds were taken out from the drum, dried completely under sunray, and stored in a plastic container. The coating procedure for the seeds of groundnut was the same as that for rice seeds, but the coating material was the mixture of SRF + RP + OM + CaCO<sub>3</sub> and the binding material was Orrin applied in the second step of the seed coating process. Due to the size variability of seeds within and between crop varieties, the coating processes produced various sizes and weight ratios of the seed to the coating material. Thus, the coated seeds were grouped

into three size categories, i.e., small (SS), medium (MS), and big (BS) coated seeds for each variety, and the corresponding weight ratio (see Table 1).

**2.3 Experimental Setting** Four sets of experiments were prepared in a glasshouse; two sets were for testing the coated seeds of two varieties rice (Impari 32 and BSM varieties), and the other two sets were for testing the coated seeds of two varieties of groundnut (Lombok and Kelinci varieties). A completely randomized experimental design was laid out for each set of experiments with a treatment consisting of uncoated seed (NC), small (SS), medium (MS), and big (BS) sizes of coated seeds, and those were triplicated. Nitrogen and phosphate fertilizers (2.0 g.pot<sup>-</sup> of N and 0.4.pot<sup>-</sup> of P<sub>2</sub>O<sub>5</sub>) were applied to the NC of rice; and phosphate fertilizer (0.4 g pot<sup>-</sup> of P<sub>2</sub>O<sub>5</sub>) was applied to the NC of groundnut. The soil used in this experiment was the 20-cm top of Ustipsamments of Akar-Akar, Northern Lombok, Indonesia, developed from pumice stones. The soil was [air-dried and](#) screened [to pass a 2-mm](#) stainless steel screen. [The](#) characteristics of the [soil](#) were sandy textured (4.2% clay, 23.3% silt, 74.5% sand), neutral (pH<sub>H2O</sub> 7.4), low N-total (0.02% N), low phosphate (116 mg.kg<sup>-</sup> of Bray II- extractable P), low organic matter (1.9% total C), low cation exchangeable capacity (4.8 cmol.kg<sup>-</sup>), Table 1. The size category and the weight ratio of seed to coating material

| Size Category | Weight Ratio |
|---------------|--------------|
| Rice          | 1:10         |
| Groundnut     | 1:8          |
| Impari 32     | 1:20         |
| BSM           | 1:15         |
| Lombok        | 1:30         |
| Kelinci       | 1:20         |
| Small (SS)    | 1:4          |
| Medium (MS)   | 1:6          |
| Big (BS)      | 1:6          |

lack of exchangeable cations (0.69, 0.22, 1.6, and 1.5 cmol.kg<sup>-</sup> of K, Na, Ca, and Mg, respectively) and) .01N-acetic acid-extractable micronutrients (7.4, 1.2, and 1.3 mg.kg<sup>-</sup> of Fe, Cu, and Zn, respectively). Sowing rice seeds: Each pot was filled with 10- kg soil, saturated with deionized water (2 L.pot<sup>-</sup>), mudded, and allowed to equilibrate overnight. Three seeds of rice were sown in each pot accordingly to the treatment at about 0.5-cm depth. After the young plants grew about 5-cm height, the water level in each pot was maintained at 1.5 – 5 cm accordingly to the growing stage of the plant. Rice was harvested at 105 days after sowing. Sowing groundnut seeds: Each pot was filled with 6-kg soil, moistened with deionized water to about field capacity (120 mL.pot<sup>-</sup>), and allowed to equilibrate overnight. One seed was sown in each pot, accordingly to the treatment, at about 0.5-cm depth. Soil moisture was maintained to about field capacity. Groundnut was harvested at 95 days after sowing.

**2.4 Statistical Analyses** The main collected data (parameters) were the germinating period of seeds and several components of growth and yield for each plant. The observed growth components for rice were the germinating period of seeds, number of tillers, and weight of dry biomass, and those for groundnut were the germinating period of seeds and weight dry biomass. The yield component for rice was dry grain, and for groundnut were the number of pods, the weight of total pods, and weight of total seeds (kernels). The analysis of variance (ANOVA) was carried out, and the test of the [least significant difference \(LSD\)](#) at  [\$\alpha = 0.05\$](#)  was applied to [the](#) appropriate [parameters](#) (which [were significantly affected by the](#) treatment).

**3. RESULTS AND DISCUSSION**

**3.1 The Growth and Yield of Rice** Statistically, [the](#) seed coating with organomineral fertilizer significantly increased the days required by the coated seeds to germinate, reduced the growth and yield components of both rice varieties (Table 2). It was also noticeable (data were not shown) that the plants of the coated seeds were most probably deficient of N, indicated by the yellowish-green of most leaves, starting in the third weeks after sowing. As shown in Table 2, the non-coated seeds (NC) germinated in six to seven days, while the coated seeds were in about teen to thirteen days after sowing; and the bigger the size, the more days required by the coated seeds to germinate. Inversely, for each rice variety, the bigger the size of coated seeds, the lower the number of tillers and the weights of biomass and grain yield. The delay of germination for the coated seeds was most probably due to the coated seeds required extra time for cracking the coating materials surrounding the seed surface. Our visual observation indicated that in the water-saturated or mudded soil, the coating materials on the surface of rice seeds started to crack in one to three days after sowing.

**3.2 The Growth and Yield of Groundnut** The effects of the

seed coating with organo- mineral [fertilizer on the growth and yield](#) components of groundnut are presented in Table 3. The seed coating depressed the germination of both varieties of groundnut, and the larger the coating size, the more time required by groundnut seeds to germinate. Inversely, the coating treatment tended to decrease the other growth and yield components. As shown in Table 3, the uncoated seeds (NC) of groundnut germinated in 7 - 8 days, whereas the coated seeds germinated in 13 - 23 days after planting, and the bigger (bigger) the coated seeds, the more time required by the coated seeds to germinate. There were different responses between the two varieties of groundnut to the seed coating treatment. For Lombok variety, the seed coating did not affect the number of pods but increased the weights of dry pods and seeds of SS and MS. For Kelinci variety, the seed treatment quadratically increased all of those components, and the maximum values were reached by the medium size of coated seeds (MS). These results suggested that each variety of groundnut having a specific size or weight of seed required a certain optimum amount of coating material. For Lombok and Kelinci varieties, the appropriate range of weight ratios for the coated seeds was 1:6 to 1:9, or 1:7.5 on average.

**3.3 General Discussion** This research was the first attempt to find out an appropriate method of seed coating for food crops with organomineral fertilizer. So, knowing the causes of unsatisfied results of the coating was very important, enabling us to fix the seed coating method with organomineral fertilizer in the further development of the seed coating. Base on available data from this research, however, it was too difficult, if not impossible, to provide a simple explanation about the primary causes of the decreases of growths and yields of both crops grown from the coated seeds. The most noticeable cause was that the coated seeds required extra time for cracking the coating material before the seeds can germinate. Specific for the coated seeds of rice, an additional cause of decreasing the growth and grain yield was N-deficiency. However, those facts were most probably not the primary causes of decreasing the growth and yield of both crops. The coating materials contained a high concentration of plantessential nutrients, and urea and Orrin were very soluble substances. This condition had brought to the thought that the coated seeds were toxified by direct contact of the high concentration of nutrients dissolved from the coating materials to the seed surface. In this research, anticipation for the seed - nutrient direct contact was made which was by applying urea or/and Orrin at the latest step in the application of the coating materials onto the seeds. But that effort seemed to be insufficient to avoid the toxifying the seeds by over- concentration of nutrients from the Table 2.

[The effects of seed coating on the growth and yield components of rice](#) Variety and parameter Seed treatment\* NC SS MS BS Impari 32: a. Germination (days) b. Number of tellers (.pot-) c. Dry biomass (g.pot-) d. Dry grains (g.pot-) 6.3a 15.0b 29.94c 39.07c 10.0b 17.3b 29.12c 39.27c 11.3b 10.7a 15.71b 34.93b 11.7b 11.0a 6.77a 29.77a Bunda Sri Madrin (BSM): a. Germination (days) 7.0a 10.3b 11.0b 12.7c b. Number of tellers (.pot-) 18.7c 13.7b 11.0a 9.3a c. Dry biomass (g.pot-) 25.43c 16.54ab 19.90b 15.16a d. Dry grains (g.pot-) 37.57b 35.87b 31.50a 29.17a \* NC, SS, MS, and BS respectively are the non-coated seeds, small coated seeds, medium coated seeds, and big coated seeds. The numbers [in the same](#) raw, [followed by the same](#) letters indicate [not significantly different based on](#) its critical value of the [LSD \$\alpha\$ =0.05](#) for each parameter [Table 3](#).

[The effects of seed coating on the growth and yield components of groundnut](#) Variety and parameter Seed treatment\* NC SS MS BS Lombok: Germination (days) Biomass (g.pot-) Number of pods (.pot-) Dry pods (g.pot-) Dry seeds (g.pot-) 7.7a 27.20b 18.3 16.00b 12.60ab 13.3a 17.47a 16.0 18.10b 15.43b 14.7bc 20.17a 18.0 18.30b 13.10ab 17.7c 18.53a 14.7 11.67a 9.30a Kelinci: Germination (days) 7.3a 16.3b 18.7bc 22.3d Biomass (g.pot-) 26.67b 21.03a 21.03a 18.53a Number of pods (.pot-) 11.7a 13.0a 13.0a 9.0a Dry pods (g.pot-) 13.33b 13.10b 13.10b 9.33a Dry seeds (g.pot-) 9.23b 10.07b 10.07b 6.97a \* NC, SS, MS, and BS respectively are the non-coated seeds, small coated seeds, medium coated seeds, and big coated seeds. The numbers [in the same](#) raw



followed by the same letters indicate not significantly different based on its critical value of the LSD $\alpha$ =0.05 for each parameter coating materials. Thus, the seed coating method described in this research needed some modification, especially for the composition of seed coating or/and binding materials, and the procedure of applying the coating materials onto the seed surface.

4. CONCLUSION AND RECOMMENDATION The seed coating with the organomineral fertilizer consisting mainly of silicate rock fertilizer, rock phosphate, and organic matter delayed the seedling and decreased most of the growth and yield components of rice but improved of the yield of groundnut. The seed coating with organo-mineral fertilizer described in this research was potentially developed as an appropriate technology to improve farming efficiency but needed further development. Some modifications must be made, especially for the composition of seed coating or/and binding materials and the procedure of applying the coating materials onto the seed surface.

ACKNOWLEDGEMENT The authors thank The Directorate of Higher Education, The Ministry of Research, Technology and Higher Education, The Republic of Indonesia for its financial support for the research.

COMPETING INTERESTS Authors have declared that no competing interests exist.

REFERENCES

1. Sherma KK, Singh US, Sharma P, Kumar A, Sharma L. Seed treatments for sustainable agriculture – A review. *J. Applied and Nat. Sci.* 2015;7(1):521-539.
2. Arias-Rivas B. Evaluation of seed coating treatment on maize (*Zea mays*) stands establishment and seed rot caused by *Pythium* spp. at early planting season. (Ph.D. thesis: Iowa State University, Ames, IA, U.S.A); 1994.
3. Castañeda LMF, Genroa C, Roggia I, Bendera SS, Benderb RJ, Pereiraa CN. Innovative rice seed coating (*Oryza sativa*) with polymer nanofibres and microparticles using the electrospinning method. *J. Res. Updates in Polymer Sci.* 2014;3(1):33-39.
4. Rahman MME, Ali ME, Ali MS, Rahman MM, Islam MN. Hot water thermal treatment for controlling seed-borne mycoflora of maize. *Int. J. Sustain. Crop Prod.* 2008;3(5):5-9.
5. Leonardos OH, Fyfe WS, Kronberg BI. The use of ground rocks in laterite systems: An improvement to the use of conventional fertilizers? *Chem. Geol.* 1987;361-370.
6. Coroneos C, Hinsinger P, Gilkes RJ. Granite powder as a source of potassium for plants: A glasshouse bioassay comparing two pasture species. *Fert. Res.* 1996;45:143-152.
7. Hinsinger P, Bolland MDA, Gilkes RJ. Silicate rock powder: Effect on selected chemical properties of a range of soils from Western Australia and on plant growth as assessed in a glasshouse experiment. *Fert. Res.* 1996;45:69-79.
8. Coventry RJ, Gillman GP, Burton ME, McSkimming D, Burkett DC, Horner NLR. Rejuvenating soils with Minplus<sup>TM</sup>, a rock dust and soil conditioner to improve the productivity of acidic, highly weathered soils. (A Report for RIRDC: Townsville, Qld); 2001.
9. Harley AD. Evaluation and improvement of silicate mineral fertilizers. (Ph.D. thesis: The University of Western Australia); 2002.
10. Priyono J, Gilkes RJ. High-energy milling improves the effectiveness of silicate rock fertilizers: A glasshouse assessment. *Comm. Soil Sci. and Plant Anal.* 2008;39: 358-369.
11. Priyono J, Arifin Z. Adding organic matter enhanced the effectiveness of silicate rock fertilizer for food crops grown on nutritionally disorder S: A glasshouse assessment. *Journal of Tropical Soils.* 2008;7(2):97-104.

© 2020

Priyono and Sudharmawan; This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited. Peer-review history: The peer review history for this paper can be accessed here: <http://www.sdiarticle4.com/review-history/52771> Asian Research Journal of Agriculture | ICI Journals Master List <https://journals.indexcopernicus.com/search/details?id=49759> ICI World of Journals (/search/form) / Asian Research Journal of Agriculture ? Back Asian Research Journal of Agriculture English title: Asian Research Journal of Agriculture ISSN: 2456-561X (online) GICID: n/d DOI: 10.9734/ARJA Website: <http://journalarja.com/index.php/ARJA/> (<http://journalarja.com/index.php/ARJA/>)

Publisher: SCIENCEDOMAIN international Country: IN Language of publication: MaiEnNpage  
 (http://jml.indexcopernicus.com) © Index Copernicus 2017 [Deposited publications](#): 268 ? [Full text](#): 49% |  
[Abstract](#): 100% | [Keywords](#): 100% | [References](#): 59% [Issues and contents](#) [Journal description](#) () [Details](#) ()  
[Scientific profile](#) () (http://indexcopernicus.com) [Editorial office](#) () 1 of 2 2/27/2022, 10:03 PM Asian Research  
 Journal of Agriculture | [ICI Journals Master List](#) <https://journals.indexcopernicus.com/search/details?id=49759>  
 Publisher () Metrics () [ISSN: 2456-561X](#) [Frequency: Monthly](#) [Journal DOI: http://dx.doi.org/10.9734/arja](#) [Peer-review model: Advanced OPEN peer review](#) [Digital Archiving: Journal Repository \(JR\)](#), LOCKSS, CLOCKSS

---

[Asian Research Journal of Agriculture \(ISSN: 2456-561X\)](#) aims to publish high quality papers in field of agricultural science. This journal facilitates the research and wishes to publish papers as long as they are technically correct, scientifically motivated. The journal also encourages the submission of useful reports of negative results. This is a quality controlled, OPEN peer reviewed, open access INTERNATIONAL journal. Subject matters include all areas of agricultural science viz., agronomy, agricultural chemistry, entomology, plant pathology, agricultural genomics, biology, chemistry, earth science, genetics & plant breeding, stored products research, post harvest biology and technology, seed science, irrigation and drainage, plant physiology, soil classification, soil fertility, soil conservation, insect and pest control, animal husbandry, aquaculture, agricultural engineering, agricultural economics, agroecology, biotechnology, organic agriculture, biodynamic agriculture, agribusiness and other allied areas. Every volume of this journal will consist of 4 issues. Every issue will consist of minimum 5 papers. Each issue will be running issue and all officially accepted manuscripts will be immediately published online. State-of-the-art running issue concept gives authors the benefit of 'Zero Waiting Time' for the officially accepted manuscripts to be published. This journal is an international journal and scope is not confined by boundary of any country or region. [Indexed in the ICI Journals Master List](#)  
[2020 ICV 2020: 94.38](#) [Archival ratings ? Citations: Coming soon](#) [MSHE points: n/d](#) [Archival ratings ? 2 of 2](#)  
 2/27/2022, 10:03 PM Priyono and Sudharmawan; [ARJA, 12\(1\)](#): 12-17, 2020; [Article no.ARJA.52771](#) Priyono and Sudharmawan; [ARJA, 12\(1\)](#): 12-17, 2020; [Article no.ARJA.52771](#) Priyono and Sudharmawan; [ARJA, 12\(1\)](#): 12-17, 2020; [Article no.ARJA.52771](#) Priyono and Sudharmawan; [ARJA, 12\(1\)](#): 12-17, 2020; [Article no.ARJA.52771](#) Priyono and Sudharmawan; [ARJA, 12\(1\)](#): 12-17, 2020; [Article no.ARJA.52771](#) 13 14 15 16 17