

PAPER • OPEN ACCESS

Mercury pollution in water and its effect on renal function of school age children in gold mining area Sekotong West Lombok

To cite this article: A Ekawanti *et al* 2021 *IOP Conf. Ser.: Earth Environ. Sci.* **637** 012055

View the [article online](#) for updates and enhancements.

Mercury pollution in water and its effect on renal function of school age children in gold mining area Sekotong West Lombok

A Ekawanti^{1*}, S Priyambodo¹, H Kadriyan², A Syamsun³, I A Lestarini⁴, G Wirasaka⁵ and A R Ardianti⁵

¹ Departement Biomedic, Medical Faculty, Mataram University, Mataram, Indonesia

² Departement Otorhinolaryngology, Medical Faculty, Mataram University, Mataram, Indonesia

³ Department Forensic, Medical Faculty, Mataram University, Mataram, Indonesia

⁴ Departement Paraclinic, Medical Faculty, Mataram University, Mataram, Indonesia

⁵ Student of Medical Faculty, Mataram University, Mataram, Indonesia.

Corresponding author: ardiana.ekawanti@unram.ac.id

Abstract. Artisanal small-scale gold mining (ASGM) has affected environment, since gold processing released mercury into the environment. Contaminated water was used as drinking water and affected human body. Kidney is one of affected organ by mercury, where children population is vulnerable for poisoned. The objective of this research was to find out mercury pollution in drinking water and its effect on renal function of children in ASGM area Sekotong in West Lombok. This preliminary study design was cross sectional. Sample of drinking water collected from various sources in Taman Sari and Telaga Lebur village. Mercury was assessed by using AAS. Renal function was evaluated from urine sample by using dipstick kit to perform proteinuria. Correlation between contaminated water consumption and renal function was statistically analyzed by Spearman's test. Mercury level found in water samples taken from well was 4.009 ppb, while from wellspring was 0.724 ppb and 0 ppb from pipe. From 15s samples, 12s samples higher than mercury threshold for drinking water. Two of 30s school age children had positive proteinuria. Spearman's test showed P-value of 0.000. It concluded that drinking water in ASGM Sekotong West Lombok has been contaminated by mercury and affected kidney function of school-age children.

1. Introduction

Mercury detected in human body is a heavy metal that comes from the environment. Mercury pollution could be from soil containing cinnabar ore, volcanoes, artisanal small-scale gold mining (ASGM), the use of cosmetics, equipment used such as battery, lamp etc. and consumption of fish [1]. Mercury contamination in West Nusa Tenggara Province (WNT) predominantly from Artisanal small-scale gold mining (ASGM) and also from nature, since rock containing cinnabar ore was found in West Nusa Tenggara (WNT) [2].

Activities of Artisanal small-scale gold mining (ASGM) in province of West Nusa Tenggara (WNT) has started in 2009 [3]. ASGM is a gold mining activity which conducted simple technology. Technology applied hazardous chemicals including mercury (amalgamation) and cyanide

(cyanidation) [4]. Mercury amalgamation has been the most common method used to recapture gold, because of the availability of mercury, it is considered to be efficient and effective [3]. The amalgamation method results in mercury emissions that get into the environment. Mercury was spilled onto the ground or agriculture land. Atmospheric transport and deposition at normal temperature is another common way for Hg to enter many water systems. In addition, Hg is often discharged together with other wastes into inadequate tailings ponds, or directly disposed into rivers and water systems [5].

Mercury and its conjugate chemicals will be absorbed by the body and cause intoxication at certain levels [1]. Once mercury enter the body, it will be distributed to almost entirely organ, the most potent affinity with neuron cell at central nervous system and kidney [6]. Many studies have demonstrated that mercury emissions affect human health [7–11]. Organ disorder depend on several factors, such as dose, route, duration of exposure, development phase (age), species of mercury and organ function [1, 12]. Study by Holmes, et al, observed that there was an increased level of low molecular weight proteins in urine of ASGM workers, although this level was only slightly above those found in the general population, but the significance of such changes in renal excretory profile is of toxicological importance [13]. Study by Ekawanti, et.al, found that 92.6 % of miner in Sekotong had proteinuria [5], so that, it is important to evaluate renal function of children in ASGM area. Krisnayanti reported that soil in Sekotong has already contaminated by mercury [3]. So that, it is important to identify the route and source of mercury from the environment to provide data for promotive health care program.

2. Materials and methods

This was a preliminary study and applied cross sectional study design. The research was conducted at ASGM area Sekotong, West Lombok Regency, West Nusa Tenggara Province. The subjects were school age children from two villages; Sekotong Tengah and Taman Sari. The source of contamination assessed by collecting sample sources of drinking water from those villages. Before conducting the study, ethical clearance was obtained from the Ethical Code Board of the Medical Faculty, Mataram University, School age children which fulfilled inclusion and exclusion criteria were selected as subjects in this study. Inclusion criteria were 6 until 12-year-old children, inhabit in ASGM area for more than one year, did not have history of kidney disorders, had permission to participate from parents by signing informed consent form.

The subjects were recruited by using a non-probability sampling technique (purposive sampling). 20 mL of urine was collected and proteinuria was assessed immediately upon collection by using dipstick urinalysis and the results were stated as g L⁻¹. 200 mL water sample collected from well, pipe and spring water which were used as drinking water in Sekotong Tengah and Taman Sari villages which the location was not more than 500 m from gelondong (ore grinder which used mercury in its process) or tong (ore grinder which used cyanide in its process). Mercury in water sample was measured using Atomic Absorption Spectrophotometry in Analytical Laboratory Mataram University and concentration stated as ppb. Water and urine data were analyzed using descriptive statistical methods and Spearman's correlation test.

3. Results and discussion

3.1. Mercury concentration in drinking water sample

Mercury concentration from drinking water samples collected demonstrated as Table 1.

Table 1. Mercury concentration in drinking water of ASGM area.

Water source	Mean of mercury concentration (ppb)	Below threshold level		Higher than threshold level	
		n	%	n	%
Well	4.009 ppb	1	6.67%	12	80%
Pipe (PDAM)	0 ppb	1	6.67%	0	0%
Spring water	0.724 ppb	1	6.67%	0	0%
Total	3.500 ppb	3	20%	12	80%

Table 1. showed that the higher mercury concentration found in well water sample namely 4.009 ppb, while in pipe water sample was found no mercury contamination.

Table 2. Comparison of mercury concentration in well water sample and reference value.

Water source	Mean of mercury concentration (ppb)	Deviation standard	T value	P value
Well	4.009 ppb	2.03		
Government standard	1 ppb	-	1.432	0.178

Table 2. demonstrated that mercury concentration in well water sample (4.009 ppb) was four times higher than of government standard value (1.00 ppb). Comparison tested by using t-test resulted p value 0.178, that was meant no significantly different with government standard.

3.2. Characteristic of research participants

This research involved thirty school-age children from two villages in ASGM area, which has characteristic as below.

Table 3. Characteristics of research participants.

Participant characteristics	Percentage (%) (n= 32)
Age	
8 year old	11 (34.4%)
9 year old	9 (28.1%)
10 year old	12 (37.5%)
Gender	
Male	13 (40.6%)
Female	19 (59.4%)
Duration of resided	
8 year old	11 (34.4%)
9 year old	9 (28.1%)
10 year old	12 (37.5%)
Parent occupation	
Government officer	4 (12.5%)
Entrepreneur	1 (3.1%)
Farmer	20 (62.1%)
Construction worker	6 (18.8%)
Others	1 (3.1%)

Table 3. presented that most participants were female (75%), and 12% were 10-year-old children. The children participated were not from miner's family but they lived in the vicinity of the smelting process within a radius of less than 500 m and around amalgamation and cyanidation process. Their duration of exposure was more than five years (mean = 9.03 years).

Table 4. Characteristics participant with proteinuria.

Duration of exposure (year)	Proteinuria	Percentage (n, %)
8	Negative	10 (31.3 %)
	Positive	1 (3.1 %)
9	Negative	9 (28.1 %)
	Positive	0
10	Negative	11 (34.4 %)
	Positive	1 (3.1 %)
Total		30 (100 %)

Table 4. displayed that of 30s samples two of them were had proteinuria at duration of exposure 8 years and 10 years. Protein was assessed qualitatively using the dip-stick method and was categorized as negative and positive. This was considered to indicate proteinuria since this level was higher than 300 mg L^{-1} (0.3 g L^{-1}). The effect of consumption contaminated water to renal function using Spearman's test, p value was 0.000. There was an impact of mercury on renal function.

All participants in the present study were directly exposed to mercury, since they has been living in the contaminated atmosphere, water, food stuff and soil in the vicinity of the mining activities. Study by Krisnayanti showed that soil and rice in Sekotong has higher mercury level than threshold [5]. Since, gold mining started at 2009 in Sekotong, so that the mercury exposure happened when the children in utero. Fetus was more vulnerable period than other period of children, immaturity of kidney affected excretion of mercury from the body and it was a mechanism of nephrotoxicity [14, 15].

Some study subjects were children that had direct contact with mercury from mercury panning after school and standing in close proximity to burning processes. This type of child activity has also been reported in other gold mining areas [7]. The duration of exposure to mercury contaminants was an average of 9.03 years. This period was much shorter than the 14.8 years needed to show specific clinical manifestations in previous reports. [9]. Other study by Ekawanti, et al, found that in miners and their families need shorter period (5.4 years) to had manifestations including proteinuria and anemia [5]. This manifestation is affected by the type and dose of mercury, the age or developmental stage, duration of exposure and route of exposure [1, 16]. Subjects in this research did not evaluated children of miners, so that they did not had an intensive contact to mercury. Although, some studies found that low level of inorganic mercury exposure did not affected renal function [17, 18]. Weakness of this study that did not identified the dose and species of mercury exposure.

Protein levels in urine were obtained by converting the qualitative proteinuria dip-stick results categorized as negative, trace, +1, +2, +3, +4 to quantitative values of 0 g L^{-1} , $<0.3 \text{ g L}^{-1}$, 0.3 g L^{-1} , 1 g L^{-1} and $\geq 20 \text{ g L}^{-1}$, respectively. Proteinuria was identified by results of +2 and $\geq 0.3 \text{ g L}^{-1}$ [19, 20]. This result was similar to findings of proteinuria in patients exposed to products containing mercury [21]. In addition, proteinuria is a clinical manifestation of mercury intoxication with elemental, inorganic and ethyl-mercury [22]. This study found that two children have proteinuria, also the same as found by same studies [5, 9, 21, 23–26]. Nephrotoxicity happened when mercury exposure level more the level for neurotoxicity. Common causes of urinary mercury excretion were elemental [27] and organic mercury exposure [8]. The source of mercury was from minerals containing cinnabar (mercury containing ore) as source of mercury from nature [2]. These children lived around small scale mining, that activity in processing gold was conducted at home, this process would directly exposed the children to mercury vapor during the smelting process. When inhaled, mercury would be dispersed rapidly into the blood and might deposit in other organs e.g. brain, kidney, placenta, thyroid and others [6]. In ASGM area Sekotong, most of the water sources for amalgamation processes and daily life come from the same well, and often the tailing ponds are nearby the well [5]. In this study found that mercury concentration in well water samples were higher than Indonesia Ministry of Health standard [28, 29].

Persistent proteinuria indicates kidney disease and the most common impacts of this condition are diabetes, hypertension, obesity and medicine or chemical substances [12, 24]. One of the chemical substances that affects proteinuria is mercury [27], and to confirm this, repeated tests are required. People with consistent positive results in repetitive measurement could be categorized as having persistent proteinuria and considered to have kidney disease. A weakness of this preliminary study was that there were no repeated tests to confirm persistent proteinuria. Evaluation of blood urea nitrogen and serum creatinine is needed in future studies.

The distance from gold processing location determined the level of mercury concentration that accumulates in well water, where the closer the distance from the gold processing location, the higher mercury concentration level [30]. In addition, according to Hanafiah study, soil nature affected mercury uptake into groundwater, where sand-dominated soil had large macro pores, soil dominated

by dust (sift) had moderate pores (meso), and those that are predominantly mud had a lot of small pores (micro). The slope of the soil affected speed of water flow which in turn will affect the deposition of mercury in water [31]. The construction of a watertight well wall is a construction that meets the requirements according to Chandra study that watertight well can reduce mercury pollution from the ground [32]. Most of well were watertight well, and all of wells were closer to gold processing location (gelondong)

4. Conclusion

It concluded that drinking water in ASGM Sekotong West Lombok has been contaminated by mercury and affected kidney function of school-age children. This result need further investigation to evaluate kidney function test.

Acknowledgments

We thank to Mataram University for funding this research. We thank to the headmaster of SDN 4 Sekotong Tengah for supporting sample collection. We thank to H. Idris for helping sample preparation and sample measurement.

References

- [1] WHO 2008 *Guidance for Identifying Population at Risk from Mercury Exposure* (Geneva: UNEP)
- [2] Gunradi R 2005 *Evaluasi Sumber Daya Cadangan Bahan Galian untuk Pertambangan Skala Kecil Daerah Pulau Lombok Provinsi Nusa Tenggara Barat (Evaluation of Mineral Reserve Resources for Small-Scale Mining in Lombok Island, West Nusa Tenggara Province)* (Lombok: Pemaparan Hasil Kegiatan Subdit Konservasi) Available from: psdg.bgl.esdm.go.id
- [3] Krisnayanti B D, Anderson C W N, Utomo W H, Feng X, Handayanto E, Mudarisna N, Ikram H and Khususiah 2012 *J. Environ. Monit.* **14** 2598
- [4] Veiga M M, Angeloci S G and Meech J A 2014 *The Extractive Industry and Society* **1** 351–361
- [5] Ekawanti A and Krisnayanti B D 2015 *J. Health Pollut.* **5** 25–32
- [6] Fornazier, Mariana Z G, Nascimento, Tatiani B, Marques, Vinicius B, Saint’Pierre, Tatiana D, Vassalo, Dalton, Brandão, Geisamanda P, Santos, Leonardo D, Carneiro and Maria T W D 2018 *Journal of the Brazilian Chemical Society* **29** 1579–1584
- [7] Bose O S 2008 *Environmental Research* **107** 89–97
- [8] Claire D B, Buchet J P, Leroyer A, Nisse C, Haguenoer J M, Mutti A, Smerhovsky Z, Cikrt M, Ochocka M T, Razniewska G, Jakubowski M and Bernard A 2005 *Environ Health Perspect.* **114** 584–590
- [9] Franko A, Budihna M V and Fikfak M D 2005 *Ann. Occup. Hyg.* **49** 521–7
- [10] Kobal A B, Flisar Z, Miklavcic V, Dizdarevic T and Briski A S 2000 *Arh. Hig. Rada Toxicol.* **51** 369–380
- [11] Lentini P, Zanolli L, Granata A, Signorelli S S, Castellino P and Dell A R 2017 *Mol. Med. Rep.* **15** 3413–3419
- [12] Kadriyan H, Sulaksana M A, Nurhidayati and Suprihartini B E 2019 *International Journal Of Nasopharyngeal Carcinoma* **1** 1–2
- [13] Holmes P, James K A F and Levy L S 2009 *Science of The Total Environment* **408** 171–182
- [14] Weidemann D K, Weaver V M and Fadrowski J J 2016 *Pediatr. Nephrol.* **31** 2043–2054
- [15] Roels H A, Hoet P and Lison D 1999 *Ren. Fail.* **21** 251–262
- [16] Limbong D, Kumampung J, Rimper J, Arai T and Miyazaki N 2003 *Science of The Total Environment* **302** 227–236
- [17] Zhang C, Gan C, Ding L, Xiong M, Zhang A and Li P 2020 *Ecotoxicology and Environmental Safety* **189** 109987
- [18] Kim N H, Hyun Y Y, Lee K B, Chang Y, Rhu S, Oh K H and Ahn C 2015 *J. Korean Med. Sci.*

30 272–277

- [19] De S D A, Halsted A C, Cote A M, Sabr Y, Dadelszen P and Magee A L 2014 *J. Obstet. Gynaecol. Can.* **36** 605–612
- [20] BPAC N Z 2013 *Interpreting Urine Dipstick Test in Adult (A Reference Guide for Primary Care)* (New Zealand: BPAC)
- [21] Li P, Du B, Chan H M and Feng X 2015 *Environmental Research* **140** 198–204
- [22] Clarkson T W 2002 *Environmental Health Perspective* **110** 11–23
- [23] Dhanapriya J, Gopalakrishnan N, Arun V, Dineshkumar T, Sakthirajan R, Balasubramaniyan T and Haris M 2016 *Indian J. Nephrol.* **26** 206–8
- [24] Sommar J N, Svensson M K, Björ B M, Elmståhl S I, Hallmans G, Lundh T, Schön S M I, Skerfving S and Bergdahl I A 2013 *Environ. Health.* **12** 9
- [25] Li S J, Zhang S H, Chen H P, Zeng C H, Zheng C X, Li S H and Liu Z H 2010 *Clin. J. Am. Soc. Nephrol.* **5** 439–444
- [26] Rana M N, Tangpong J and Rahman M M 2018 *Toxicol Rep.* **5** 704–713
- [27] Abdennour C, Khelili K, Boulakoud MS, Nezzal A, Boubsil S and Slim S 2002 *Environmental Research* **89** 245–249
- [28] Menkes RI 1990 *Peraturan Menteri Kesehatan Nomor: 416/MEN.KES/PER/IX/2009 (Minister of Health Regulation Number: 416/MEN.KES/PER/IX/2009)* (Indonesia: Minister of Health of the Republic of Indonesia)
- [29] Menkes RI 2010 *Peraturan Menteri Kesehatan Nomor: 492/MEN.KES/PER/IV/2010 (Minister of Health Regulation Number: 492/MEN.KES/PER/IV/2010)* (Indonesia: Minister of Health of the Republic of Indonesia)
- [30] Boky H, Umboh J and Ratag B 2015 *Jurnal Ilmu Kesehatan Masyarakat Unsrat* **5** 63–70
- [31] Hanafiah K A 2005 *Dasar-Dasar Ilmu Tanah (Basics of Soil Science)* (Jakarta: Raja Grafindo Persada)
- [32] Chandra B 2007 *Pengantar Kesehatan Lingkungan (Introduction to Environmental Health)* (Jakarta: Penerbit Buku Kedokteran)

IOP Mercury pollut

_2021_IOP_Conf_Ser_Earth_Environ_Sci_637_011001.pdf - Adobe Acrobat Reader DC (64-bit)

File Edit View Sign Window Help

Home Tools _2020_IOP_Conf... _2021_IOP_Conf.... x _2021_IOP_Conf.... _2021_IO

Save Star Cloud Print Search Previous Next Page Zoom

The 7th International Conference on Sustainable Agriculture and Environment IOP Publishing
IOP Conf. Series: Earth and Environmental Science 637 (2021) 011001 doi:10.1088/1755-1315/637/1/011001

Preface

Contemporary events show us that we humans are very vulnerable to economic and environmental fluctuations. As humanity progress, produce and consume like it will exist until infinity. Sustainability has always been the case, is accepted as life guidance but until then we; researchers, companies and students should try to work on this concept so that next generation can benefit from our efforts.

Sustainable way of dealing with environment and agriculture is not easy. The benefits from sustainable agriculture vary and sustainable agriculture doesn't have harmful effects on environment or sources. In agricultural activities have been performed for chaining the environment or utilizations of environment as sources with maximum benefits. If we do not success that, it is impossible for us to talk about modern agriculture. Under poor management of agriculture, biodiversity and soil properties can be damage, contaminate and some problems limiting efficient agricultural activities may be occurred.

The main objective of farming is to produce food for human being, but unfortunately some parts of the farmlands have used also as settlements areas. By this way, some natural resources have been deteriorated; consequently, size of farmland has reduced day by day. In that regard, it is possible to say that we are in going in a circle. More than one-third of food supplies is as form of wastes due to the over buying than required amounts of food by costumers.

In the lights of the information mentioned above, there is no doubt that continues sustainable agricultural activities are practical solution to prevent the human being from the starvation. The International Conference on Sustainable Agriculture and Environment (ICSAE) series were aimed to provide a platform for researchers and academics as well as practicing professionals from all over the world, to present their research and professional development activities in agriculture, environment, food and other relevant subjects. This conference series was an effort to identify the ideas, practices and policies that constitute our concept of sustainable agriculture. The concept of sustainable agriculture itself is still evolving and thus, we published not as a definitive or final statement, but as an invitation to continue our dialogue. Moreover, as we experienced lately during COVID-19 pandemic situation, agriculture is a sector which stay struggle and can be source of many materials for fight the diseases.

The current ICSAE-7 at 2020 was held during global pandemic situation, for that the committee decided to shift from offline to online meeting. It's not an easy task by the way, there were travel banned, different time zone, and internet connection are factors need to be overcome during the organization of this event. Question may be arising, why the ICSAE is not postponed. Well, it is the commitment of scientist to communicate the agriculture issues during pandemic situation, moreover the event also has been planned well since last year. After some adjustments, and considering available resources, the conference then held virtually at 25-27 August 2020.

Committee use online meeting platform for two main agendas in this event. First, the parallel session that was held for 3 days in row using website as the media for the researcher to deliver their talk. Each author(s) has 15 minutes to talk about their paper, recorded then uploaded both in Youtube and in the virtual venue (conference website). All the documentation of author(s) talk can be accessed in the conference website (<http://icsae.id>). Committee set 3 days online meeting, since realizing the time zone different and internet connection differences among the participant(s). Using this strategy, the talks can be enjoyed by the participants during their stay at home, or can be accessed anywhere and anytime. Participants can raise question in every single uploaded talk, by typing their question in the conference website. The system will notify the author(s), then question can be answer it accordingly.

Content from this work may be used under the terms of the Creative Commons Attribution 3.0 license. Any further distribution of this work must maintain attribution to the author(s) and the title of the work, journal citation and DOI.
Published under license by IOP Publishing Ltd 1

This site uses

The 7th International Conference on Sustainable Agriculture and Environment
IOP Publishing
IOP Conf. Series: Earth and Environmental Science 637 (2021) 011001 doi:10.1088/1755-1315/637/1/011001

The second program is the General Assembly meeting which held at 27 August 2020. This program was held by online via Zoom meeting, and there were 6 keynote speakers invited and delivering talk about agriculture and its challenge during pandemic situation. Both meeting program, parallel session and general assembly, were organized virtually by committee which stay in Universitas Sebelas Maret Surakarta Indonesia, and in Turkey by the help of Dr. Mithat Direk.

As can be listed, committee receive participation around the globe such as from Europe, Turkey, Egypt, Pakistan, China, India, Indonesia, Japan, Taiwan and etc. Officially, committee receive online registration to participate this virtual conference around 320 participants from around the world. As the commitment to open access all the conference talk, we let the virtual venue open and can be visited online, until now (15 November 2020) the virtual venue has been visited by more than 1,850 viewers from around the world.

Overall, the major obstacles were internet connection stability which vary between participant(s) in his/her region. A condition of can't join the live meeting, breaking up voice during talk, can't see screen share during presentation at the participant(s) side, were associated with the connection. However, according to the participant(s) feedback, ICSAE-7 conference was successfully held by virtually and this was a new experience in participation the international conference.

Lastly, committee would like to thank whole people being at Scientific Board, Managerial Board, authors or participants by supporting their valuable works. Particularly Prof. Ahmad Yunus from Universitas Sebelas Maret Indonesia and his team, for all their contributions on continuity of these conference series since it was held, for the 1st, at 2013 in Surakarta Indonesia, then move around the globe and at 7th ICSAE its back again in Surakarta. I do appreciate all the efforts given to prepare this event. This is the first digital conference of ICSAE series which could be reach wider and bigger participant around the world.

ICSAE-7 2020
Chair

The 7th International Conference on Sustainable Agriculture and Environment
IOP Conf. Series: Earth and Environmental Science 637 (2021) 011002 doi:10.1088/1755-1315/637/1/011002

Committee of ICSAE 2020

Honorary Planning Committee (in alphabetical order)

1. Prof. Dr. Ahmad Yumuz, Vice Rector 1, Sebelas Maret University, Indonesia
2. Dr. Mustafa Şahin, President, Selcuk University, Turkey
3. Dr. Muhammad Sarjan, Mataram University, Indonesia
4. Prof. Dr. Samanhudi, Dean Faculty of Agriculture Sebelas Maret University, Indonesia

Conference Chairs (in alphabetical order)

1. Dr. Mithat Durek, Agricultural Economy, Selcuk University, Konya, Turkey
2. Dr. Sigit Prastowo, Animal Science, Sebelas Maret University, Indonesia

Technical Program Committee (in alphabetical order)

1. Dr. Ahmed Mohammed Ahmed, Agribusiness & Applied Economic, Tanta University, Egypt
2. Dr. Azam Kakar, Biotechnology, Quetta, Pakistan

Conference Secretary (in alphabetical order)

1. Dr. Adi Rathiyanto, Animal Science, Sebelas Maret University, Indonesia
2. Dr. Amalia Tetrani Sakya, MSc. Department of Agrotechnology, Sebelas Maret University, Indonesia

Treasurer

1. Prof. Dr. Sri Hartati, Agrotechnology, Sebelas Maret University, Indonesia

Organizing Committee Members (in alphabetical order)

1. Ms. Desyya Putri
2. Ms. Rina Puji Lestari
3. Ms. Rissa Rahmadhiyati
4. Mr. Rohmad Setiaji
5. Ms. Wafa Nur Hanifah
6. Ms. Winna Candra M.

Scientific Committee Members (in alphabetical order)

1. Prof. Dr. Ahmed Mohamed Ahmed, Tanta University, Egypt
2. Dr. Aluh Nikmatullah, Agroecotechnology, Faculty of Agriculture, Mataram University, Indonesia
3. Dr. Anissa Gara, Institut national agronomique de Tunisie and Institut national de recherches agronomiques de Tunisie, Tunisia
4. Dr. Agha Ali, Institute of Agricultural and Resource Economics, Faculty of Social Sciences, University of Agriculture, Pakistan
5. Prof. Dr. Bambang Hari Kusumo, Postgraduated Program, Mataram University, Indonesia
6. Prof. Dr. Bambang Pujiasmanto, College of Agriculture, Sebelas Maret University, Indonesia
7. Dr. Bilal Acar, Irrigation and Agricultural Building Department, Selcuk University, Turkey
8. Dr. Bilal Cemek, Agricultural & Biosystems Engineering, Ondokuz Mayis University, Turkey

Content from this work may be used under the terms of the [Creative Commons Attribution 3.0 license](#). Any further distribution of this work must maintain attribution to the author(s) and the title of the work, journal citation and DOI.
Published under licence by IOP Publishing Ltd

This si

Document

_2021_IOP_Conf._Ser._Earth_Environ._Sci._637_011003.pdf - Adobe Acrobat Reader DC (64-bit)

File Edit View Sign Window Help

Home Tools _2020_IOP_Conf.... _2021_IOP_Conf.... _2021_IOP_Conf....

1 / 3 64,4% 1 2 3 4 5 6 7 8 9 10 11 12 13 14 15 16 17 18 19 20 21 22 23

IOP Conference Series: Earth and Environmental Science

PAPER • OPEN ACCESS

Peer review declaration

To cite this article: 2021 IOP Conf. Ser.: Earth Environ. Sci. 637 011003

View the [article online](#) for updates and enhancements.

You may also like

- Peer review declaration
- Peer review declaration
- Peer review declaration

Page

The Electrochemical Society Advancing science & electrochemical science & technology

242nd ECS Meeting Oct 9 – 13, 2022 • Atlanta, GA, US

Early hotel & registration pricing ends September 12

Presenting more than 2,400 technical abstracts in 50 symposia

The meeting for industry & researchers in

BATTERIES ENERGY TECHNOLOGY SENSORS AND MORE!

Register now!

ECS Plenary Lecture featuring M. Stanley Whittingham, Binghamton University, Nobel Laureate – 2019 Nobel Prize in Chemistry

This content was downloaded from IP address 103.133.160.14 on 02/09/2022 at 08:08

This s

Document2

_2021_IOP_Conf_Ser_Earth_Environ_Sci_637_011003.pdf - Adobe Acrobat Reader DC (64-bit)

File Edit View Sign Window Help

Home Tools _2020_IOP_Conf.... _2021_IOP_Conf.... _2021_IOP_Conf.... _2021_IOP_Conf....

Save Star Cloud Print Search Up Download 2 / 3 Hand Zoom Out Zoom In 64,5%

The 7th International Conference on Sustainable Agriculture and Environment IOP Publishing
IOP Conf. Series: Earth and Environmental Science 637 (2021) 011003 doi:10.1088/1755-1315/637/1/011003

Peer review declaration

All papers published in this volume of IOP Conference Series: Earth and Environmental Science have been peer reviewed through processes administered by the Editors. Reviews were conducted by expert referees to the professional and scientific standards expected of a proceedings journal published by IOP Publishing.

- Type of peer review: Single-blind
 - All the full papers submitted and presented in ICSAE-7 were reviewed by single blind review method. Firstly, along with registration process, author(s) are requested to submit their short abstract. Following registration, abstract then screened by conference committee. Screening was done by the conference committee by considering its relevancy with conference topic. Once its fit, then committee send notification to the author for acceptance to be presented and author(s) are requested to send the full paper and preparing their conference talk.
 - The full paper then sent to the expert reviewer for comments and criticism, returning back to the author(s) for revision, one or two round. In here, papers which not meet the criteria or can't be revised according to the reviewer comments were rejected for further process. The revised paper then submitted to the conference committee for proof read, layout, camera ready paper production, and similarity test. Finally approved paper then submitted to IOP.
- Conference submission management system:
The ICSAE-7 2020 use Conference Management Toolkit as submission management system at the following link: <https://cmt3.research.microsoft.com/User/Login?ReturnUrl=%2FICSAE2020%2Fhome>.
- Authors can track their paper progress using submission system, and also the committee inform the update by e-mail.
- Number of submissions received: 143 abstracts
- Number of submissions sent for review: 114 papers
- Number of submissions accepted: 98 papers
- Acceptance Rate [Number of Submissions Accepted / Number of Submissions Received X 100]: 85,96%
- Average number of reviews per paper: -
- Total number of reviewers involved: 7 reviewers
- Any additional info on review process:
Along with the revised paper submission, author(s) also send to the committee the signed author agreement which declare the following:
 1. The manuscript contains original, unpublished results which are not currently being considered for publication elsewhere
 2. The manuscript has been read and approved by all the authors
 3. The corresponding author or primary contact is responsible for communicating with the other authors about progress, submissions of revisions, signed the agreement form and final approval of the manuscript

Content from this work may be used under the terms of the Creative Commons Attribution 3.0 licence. Any further distribution of this work must maintain attribution to the author(s) and the title of the work, journal citation and DOI.
Published under licence by IOP Publishing Ltd. 1 This site

IOP Mercury pollution in water and X IOP IOP Conference Series: Earth an X +

Peer review declaration

+ Open abstract View article PDF

Papers

OPEN ACCESS 012001
Strategy for reducing environmental disaster based on social capital (A case study in Indonesia)
A Wibowo, R Karsidi, B Sudardi and M Wijaya
+ Open abstract View article PDF

OPEN ACCESS 012002
Semar Sandy-app: monitoring system for sandy soil irrigation based on android application
I Arifin, A Herawati, Sumani Mujijo and H Widjianto
+ Open abstract View article PDF

OPEN ACCESS 012003
The importance of tropical edible fruit plants for tribal communities in East Aceh region, Indonesia
Z I Navia, A B Suwardi and Nuraini
+ Open abstract View article PDF

OPEN ACCESS 012004
Seasonal variation in hybrid seed adoption: The case of chilli in Indonesia
A L Sayekti, R A Nugrahapsari and A M Hasibuan
+ Open abstract View article PDF

OPEN ACCESS 012005
The effect of row proportion of maize and soybean intercropping on growth and yield of component crops in sandy soil North Lombok, Indonesia
W Astiko, N M L Ernawati and I P Silawibawa
+ Open abstract View article PDF

OPEN ACCESS 012006
Mung bean cover crop improved soil organic carbon and maize yield in a semiarid area
I K D Jaya, Sudirman and I W Sudika
+ Open abstract View article PDF

This site uses cookies. By continuing to use this site you agree to our use of cookies. To find out more, see our [Privacy and Cookies](#) policy.



[IOP Mercury pollution in water and](#) [IOP Conference Series: Earth an](#) +

← → ⌂ https://iopscience.iop.org/issue/1755-1315/637/1 ⌂

+ Open abstract [View article](#) [PDF](#)

OPEN ACCESS 012055
Mercury pollution in water and its effect on renal function of school age children in gold mining area Sekotong West Lombok
A Ekawanti, S Priyambodo, H Kadriyan, A Syamsun, I A Lestarini, G Wirasaka and A R Ardianti
+ Open abstract [View article](#) [PDF](#)

OPEN ACCESS 012056
Carbon stock estimation on some land cover: secondary forest, agroforestry, palm oil plantation and paddy fields
S S Karuru, B Rasyid and S Millang
+ Open abstract [View article](#) [PDF](#)

OPEN ACCESS 012057
The application of amendments for growth and nutrient absorption of medicinal plant *Petiveria aleacea* on entisol soils type
Sulandjari, AT Sakya, J Syamsiah and Viviana
+ Open abstract [View article](#) [PDF](#)

OPEN ACCESS 012058
The effects of catfish oil supplementation as unsaturated fatty acid source on Bali cow gas production kinetics, dry matter digestibility, and organic matter digestibility in vitro
D N Cahyo, L M Yusiatyi, A Kurniawati, C Hanim and Muhiisin
+ Open abstract [View article](#) [PDF](#)

OPEN ACCESS 012059
Economic feasibility analysis of service business of agricultural equipment and machinery in Kubu Raya Regency, Indonesia
Ekawati, Ellyta and S Sugardi
+ Open abstract [View article](#) [PDF](#)

OPEN ACCESS 012060
Increasing nutrient quality and in vitro digestibility of mixed feed oil palm waste for ruminant feed by solid-state fermentation technology
M N Rofiq, S Martono and R A Gopar
+ Open abstract [View article](#) [PDF](#)

This site uses cookies. By continuing to use this site you agree to our use of cookies. To find out more, see our [Privacy and Cookies](#) policy.

