

Jurnal 2

by andre scabra

Submission date: 10-Oct-2022 09:13AM (UTC+0700)

Submission ID: 1901230126

File name: JURNAL_20H_26P_20_2019.pdf (1.67M)

Word count: 4711

Character count: 25575

JOURNAL OF HEALTH & POLLUTION

**PURE
EARTH**
BLACKSMITH INSTITUTE

Volume 8 - Number 20 | December 2018

ISSN 2156-9614



Ponmo vending in Ketu Market, Lagos, Nigeria. Ponmo is a low cost food source for many people across Africa. Questions have been raised as to it's safety. Photo by Hadijat Oluwatobi Osilagun, 2017

Journal of Health and Pollution Editorial Board

Tom Achoki, MD, PhD, Sloan Fellow, MIT Sloan School of Management, USA. Research Fellow, Utrecht Institute for Pharmaceutical Sciences, Utrecht University, The Netherlands

Nicholas Albergo, PE, DEE, MSCE, Environmental Professor of Practice, Civil and Environmental Engineering Department, College of Engineering, University of South Florida, Tampa, Florida, USA

Andy Beke, MMed, FCPHM, FGCP. Professor, School of Health Systems and Public Health, University of Pretoria, Pretoria, South Africa.

Nil Basu, PhD, Associate Professor, Canada Research Chair in Environmental Health Sciences, School of Human Nutrition, McGill University, Montreal, Canada

Ana Boischio, PhD, MSc, Regional Advisor, Toxicology, Sustainable Development and Health Equity, Pan American Health Organization, Washington DC, USA

Stephan Bose-O'Reilly, MD, MPH, Workgroup leader Paediatric Environmental Epidemiology, WHO Collaborating Center for Occupational Health, Munich, Germany

Anne Riederer, ScD, Affiliate Assistant Professor, Environmental and Occupational Health Sciences, School of Public Health, University of Washington, USA, Professorial Lecturer, Environmental and Occupational Health, George Washington University, USA

Inoka Suraweera, MBBS, MSc, MD, Specialist in Community Medicine, Consultant Community Physician and Unit Head (Environmental and Occupational Health), Environmental and Occupational Health Directorate, Ministry of Health, Colombo, Sri Lanka

Mark Patrick Taylor, PhD, Professor, Department of Environmental Sciences, Macquarie University, Sydney, Australia

Mitko Voutchkov, PhD, Senior Lecturer, Department of Physics, The University of the West Indies, Kingston, Jamaica

Editor in Chief

Jack Caravanos, DrPH, CIH, Clinical Professor, Environmental Public Health, College of Global Public Health, New York University, New York, USA

Managing Editor

Sandy Page-Cook, MPH, sandy@journalhealthpollution.org

Director of Manuscript Editing

LeAnn Moore, MA, MPH

Editorial Assistant

This site uses cookies. By continuing to use our website, you are agreeing to [our privacy policy](#). [Accept](#)

Volume 9, Issue 21

1 March 2019

ISSN 2156-9614

In this Issue

Editorial

RESEARCH

Conflicting Conclusions or Competing Methodologies? Documenting Soil Lead Pollution in Owino Uhuru, Kenya

Jack Caravanos

Extract

View Article

PDF

RESEARCH

Risk of Mercury Exposure from Fish Consumption at Artisanal Small-Scale Gold Mining Areas in West Nusa Tenggara, Indonesia

Muhammad Junaldi; Baiq Dewi Krisnayanti; Juharfa; Christopher Anderson

Abstract

View Article


PDF

Bioaccumulation of Heavy Metals by *Clarias gariepinus* (African Catfish) in Asa River, Ilorin, Kwara State

Olaniyi Afolabi Opasola; Adedotun Timothy Adeolu; Ayodeji Yusuf Iyanda; Solomon Olayinka Adewoye; Sunday Asabi Olawale

13

Risk of Mercury Exposure from Fish Consumption at Artisanal Small-Scale Gold Mining Areas in West Nusa Tenggara, Indonesia

Muhammad Junaidi,¹ Baiq Dewi Krisnayanti,² Juharfa,¹ Christopher Anderson³ 

1 Department of Fish and Aquaculture, University of Mataram, Mataram, Indonesia

2 International Research Centre for the Management of Degraded and Mining Lands, Indonesia

3 Soil and Earth Science, Institute of Natural Resources, Massey University, Palmerston North, New Zealand

Corresponding author:
Baiq Dewi Krisnayanti
bdewi.krisnayanti@gmail.com

Introduction

The use of mercury (Hg) to recover gold in artisanal small-scale gold mining (ASGM) activities has contributed to widespread mercury contamination of aquatic systems in many areas worldwide.¹⁻⁵ Across ASGM areas in Indonesia, it is common to see waste from ASGM activities discharged directly into agricultural lands and water bodies that empty into the sea.⁶

Mercury species are present in aqueous media, including methylmercury. This species can be absorbed by aquatic biota and accumulate in aquatic organisms. As a result, mercury concentrations can be magnified at higher levels

Background. The primary environmental risk associated with artisanal small-scale gold mining (ASGM) activities in Sekotong and Taliwang is waste discharged directly into the environment. This waste contains variable concentrations of heavy metals and a high level of mercury. When these elements are released into the environment, plants and animals can be contaminated. If mercury is methylated to methylmercury, levels can increase in concentration at each level of the food chain (biomagnify). Fish are a primary risk vector for methylmercury poisoning in humans, and represent a significant source of protein for the Sekotong and Taliwang communities.

Objectives. The present study aimed to identify the concentration of mercury in fish from ASGM sites in Sekotong and Taliwang.

Methods. Descriptive research was used to describe the mercury concentrations of fish in the present study. The fish species collected for the samples represented commercially available fish most commonly consumed by the community on a daily basis.

Results. In Sekotong's ASGM area, the mercury concentration in *Pilsbryconcha exilis* tissue was 596 ppb, 721 ppb for *Sephia officinalis* and 50% of the *Euthynnus affinis* samples had a high level of mercury, above the World Health Organization (WHO) maximum permissible limit for the sale of fish for human consumption of 0.5 ppb.

Conclusions. Some fish species from the studied ASGM sites had high mercury concentrations above the maximum permissible mercury concentration in edible fish tissue. The risks associated with mercury exposure from fish consumption threaten community health.

Ethics Approval. All experiments were performed in accordance with relevant local guidelines and regulations.

Competing Interests. The authors declare no competing financial interests.

Keywords. ASGM, fish, mercury

Received October 17, 2018. Accepted January 25, 2019

J Health Pollution 21: (190302) 2019

© Pure Earth

of the food chain. In this system, mercury is chemically transformed, bioaccumulated and biomagnified. Thus, fish may accumulate high levels of mercury and can act as a pathway for the mercury to be consumed by people or wildlife.^{2,7} Concentrations of total mercury in human hair are significantly correlated with frequency of fish consumption.⁸

Human exposure to methylmercury

(MeHg) occurs most frequently from consumption of marine and freshwater fish. This has become a public health challenge as fish are an important source of nutrition globally. In addition, fish are culturally important for many communities and are a profitable global commodity. The chemical form of MeHg in fish tissue has been detected in fish protein, and is not removed or destroyed by cooking or cleaning.⁹

23

16

Fish from the ASGM site in Tatelu, South Sulawesi, Indonesia have accumulated high concentrations of mercury in fish tissue due to excessive losses of mercury from mining activities to the aquatic system, contributing to the high bioavailability of mercury in the Talawan River.⁴ Similarly, concentrations of mercury were significantly high in freshwater, estuarine and marine fishes around the area of ASGM activities in Suriname.¹⁰

Signs of mercury contamination have been found in seafood, molluscs, and crustaceans downstream of ASGM sites at Buru Island, Indonesia due to the geochemistry of sediments in which mercury methylation leads to highly bioavailable mercury that threatens the aquatic food chain and marine resources.¹

High mercury concentrations from amalgamation and cyanidation tailings have been analyzed from Sekotong and Taliwang, with mean mercury concentration of the amalgamation tailings of about 3000 µg/g and greater than 1600 µg/g for the cyanidation tailings, exceeding the maximum permissible concentration for mercury in soil set by the Indonesian government of 2 µg/g.⁶ Mining in Sekotong began in mid-2009, and by 2012, 70% of Sekotong's miners had been exposed to mercury, as indicated by mean concentrations of mercury in hair samples between 7.72 – 11.61 µg/g, exceeding the recommended limit of 1 µg/g set by the World Health Organization (WHO).¹¹ In 2015, Ekawanti and Krisnayanti reported that 61% of urine samples and 81% of hair samples of miners in Sekotong contained alarming levels of Hg.¹² The participants in the study were directly exposed to mercury, either as workers (miners) or as family members (non-miners) living in the contaminated area, including children that were exposed to mercury through

Abbreviations

ASGM	Artisanal small-scale gold mining	WHO	World Health Organization
------	-----------------------------------	-----	---------------------------

direct inhalation in close proximity to burning processes. The duration of exposure to mercury contaminants in Sekotong's ASGM area was an average 12.4 years. This period was much shorter than the 14.8 years needed to show specific clinical manifestations in previous reports.¹³ Krisnayanti found similar evidence of mercury exposure around Taliwang ASGM activities where the mean concentrations of mercury in hair of exposed, indirectly exposed and non-exposed groups were 13, 1.3 and 0.56 µg/g, respectively.¹⁴ The ASGM sector in Taliwang has a large migrant worker population, which is an important economic support to the local community, but ASGM activities involve high mercury use and illegal mercury trading. These activities have affected the health of miners in a relatively short time, as evidenced by the high mercury residue on the bodies of miners. This indicates that mercury exposure has occurred through inhalation, however a second exposure pathway through consumption of methylmercury-contaminated food (rice) and fish could occur in the future.^{6,15} The primary environmental risk associated with ASGM activities in Sekotong and Taliwang is waste discharged directly into the environment. This waste contains variable concentrations of heavy metals, such as soluble complexed mercury, including mercury cyanide.

Since fish are a primary risk vector of methylmercury poisoning in humans and represent a significant source of protein for ASGM communities,

the consumption of fish caught around the ASGM area represents a possible human health risk. Fish is an important food for the Sekotong community as Sekotong is located along the Lombok Strait. Marine fish/seafood and freshwater fish are also an important staple food in the Taliwang diet. The aim of the present study was to identify the concentration of mercury in fish from Sekotong and Taliwang's ASGM sites.

Methods

Fish sampling sites were located in two ASGM sites in West Nusa Tenggara Barat, in Sekotong District, West Lombok Regency at Lombok island and Taliwang District, West Sumbawa Regency at Sumbawa island. The characteristics of Sekotong and Taliwang sea waters are influenced by both the Pacific and Indian Ocean where the mass of water originating from the northern Pacific enters the Sulawesi Sea to the south of Mindanao, then enters the heart of Indonesian waters through the Makassar Strait. At the end of the Makassar Strait, this line branches into two, with water either flowing directly to the Indian Ocean through the Lombok Strait (West Nusa Tenggara) or east through the Flores Sea to the Banda Sea.¹⁶

The fish species collected for the samples from these areas represented commercially available fish species most commonly consumed by the community on a daily basis for the study season and the most abundant catch of the season.




Figure 1 — Research location: Sekotong District, West Lombok Regency, West Nusa Tenggara Province




Figure 2 — Fish sampling sites: Taliwang District, West Sumbawa Regency, West Nusa Tenggara Province

ASGM activities at Sekotong began in mid-2009, and were gradually reduced in 2014, when only 15% of miners were still mining. Krisnayanti calculated that there were around 4630 of trommel (using 250-500 gram of mercury/trommel) and 150 cyanidation tanks in Sekotong, involving around 14,500 miners.⁶ In

contrast, ASGM activities in Taliwang began in 2010, and mining activity has gradually increased without signs of stopping compared to Sekotong. In 2013, there were around 5000 trommel (using 250-500 gram of mercury/trommel) and 95 cyanidation tanks involving around 15,380 miners.¹⁶

Sekotong sampling site

Fish sampling at Sekotong was conducted in November 2016. There are three main traditional markets in this area, and therefore fish were sampled from all of these markets. In addition, fish were also collected from spots where fishermen most commonly sell their daily catch to local markets or local personal buyers. However, fish are also sold to markets in Mataram City and personal buyers who own restaurants in Mataram. Fish were collected from three main markets in Sekotong District: Sekotong Tengah (1), Tawun (2), and Pelangan (3) and two spots where fisherman sell their catch directly: Batu Putih (4) and Tembowong (5) (Figure 1). Figure 1 also shows the main ASGM spots across Sekotong District from 2009-2014. These spots were identified directly by the co-author in previous research.¹⁷

Taliwang sampling site

Fish sampling was conducted in Taliwang Regency in November 2016. The fish were collected from three main fishing spots: Labuan Lalar (1); Taliwang estuary (2), Lake Taliwang (3) and the main market at Taliwang (4) (Figure 2). Figure 2 also demonstrates ASGM activities across the Taliwang area in 2013. These spots were identified directly by the co-author in previous projects.

During sampling, fish were identified and length, size and fresh weight of fish were recorded. Mercury concentrations can increase with increased length and age. Mercury also biomagnifies and generally increases in concentration with increasing food chain length.^{11,18}

Each fish specimen was filleted, then dried to a constant weight at 40°C. The

fish were thoroughly homogenized in a blender and bagged in plastic zip lock bags for laboratory analysis. The fish samples were analyzed at "Laboratorium Penelitian and Pengujian Terpadu" University of Gadjah Mada Yogyakarta for total mercury using a mercury analyzer (type 254) with measuring range of 0.01-10 ppb. Mercury detection limits were 0.17 µg/g dry weight. The laboratory is accredited by the Indonesian government.

Fish samples were pre-digested overnight at room temperature with aqua regia (analytical grade nitric acid: hydrochloric acid =1:3). The next day, the samples were digested at 120°C for 24 hours. Samples were filtered using Whatman No. 42 filter paper into a 100 ml volumetric flask. Deionized water was used to quantitatively transfer the solution into the volumetric flask and to make up the solution to the 100 ml mark. Five (5) mL of sample solution with 5 mL of nitric acid and 1 mL of freshly prepared tin chloride (hydrochloric acid, stannous chloride) was placed into a measurement flask and prepared for measurement. For analytical precision and quality control, reagent blank solutions were analyzed in parallel with the samples. Linear calibration was prepared from 1000 mg/L of mercury standard no. 16482 as a standard reference material which was issued by TraceCERT™ and accredited by ISO/IEC 17025 and ISO Guide 34.^{19,20}

A number of countries and the WHO have established a threshold for mercury concentration in fish and fish products in order to protect human health.¹¹ The WHO's maximum permissible mercury concentration in edible fish tissue is 0.5 µg Hg g⁻¹, which is equal to 0.5 ppm or 500 µg Hg kg⁻¹, equal to 500 ppb fish tissue (wet mass basis). Mercury concentrations for

Tropic Category	Species	No. of specimens (n)	Weight (g)	Length (cm)
Planktivore	<i>Anadara granosa</i>	50	-	-
	<i>Caesio cuning</i>	3	340±148.49	24±4.24
	<i>Pilsbryconcha exilis</i>	30	-	-
	<i>Pomacanthus imperator</i>	2	632±7.07	21.5±2.12
	<i>Sardinella lemuru</i>	35	33.75±7.41	13.38±1.11
	<i>Siliqua patula</i>	30	-	-
Carnivore	<i>Chromileptes altivelis</i>	1	27	12
	<i>Euthynnus affinis</i>	13	153±46.03	21.67±1.15
	<i>Lethrinus erythracanthus</i>	1	-	-
	<i>Loligo indica</i>	40	43.5±7.55	17.25±6.08
	<i>Nemipterus nematophorus</i>	4	127.25±2.06	16.5±2.52
	<i>Sepia officinalis</i>	6	43.5±7.55	17.25±6.08
Herbivore	<i>Lethrinus erythracanthus</i>	1	168	22
	<i>Ctenochaetus striatus</i>	5	122±1.73	14±1.73
	<i>Naso sp.</i>	5	43.5±7.55	24.75±7.63
	<i>Plectorhincus gaterinoides</i>	2	390±1.41	23.5±0.71
	<i>Parupeneus multifasciatus</i>	3	162.67±1.15	19.67±1.15
	<i>Siganus sp.</i>	46	201.25±79.61	18.92±2.65
	<i>Scaridae sp.</i>	23	375±264.88	21.9±5.36
	<i>Valamugil seheli</i>	5	261±3.54	22.5±3.54
Omnivore	<i>Scolopsis affinis</i>	3	246.33±110.59	22±2
	<i>Acanthurus sp.</i>	11	196.67±47.34	17.67±1.53

Table 1 — Characteristics of Fish Sampled at Sekotong

the results of the present study are presented in ppb.

Results

Some fish species from the studied ASGM sites had high mercury concentrations above the maximum permissible mercury concentration in edible fish tissue.

Sekotong

The fishing area in West Lombok spreads across the Lombok Strait, Sekotong Bay, and the Indian Ocean. Sekotong Bay is an important fishing area especially for pearl shell, grouper fish and crayfish (lobster). In the Sekotong area, 315 marine fish specimens and mollusks were collected, predominately *Loligo indica*, *Sepia officinalis*, *Siganus sp.*, *Caesio*

cuning, *Scolopsis affinis*, *Valamugil seheli*, *Euthynnus affinis*, *Sardinella lemuru*, *Scaridae sp.*, *Acanthurus sp.*, *Naso sp.*, *Lethrinus sp.*, and *Parupeneus multifasciatus*, and the mollusks included *Anadara granosa*, *Siliqua patula*. No freshwater fish were collected from this area.

Previous studies have found that mercury concentrations in fish tissue were correlated with age, length, and weight of fish, but in the present study, there were no significant differences between fish length, weight and Hg concentration at a significance level of $p > 0.05$ (one-way analysis of variance) and Table 1 indicates that there was a substantial variation in fish growth rate.^{11,18}

Among the planktivores collected in Sekotong, only *Pilsbryconcha*




Figure 3 — Mean level of total mercury in planktivorous fish at Sekotong against the maximum permissible mercury concentration in edible fish tissue of 500 ppb




Figure 4 — Mean level of total mercury in carnivorous fish at Sekotong against the maximum permissible mercury concentration in edible fish tissue of 500 ppb

exilis (shells) showed high mercury concentration above the maximum permissible mercury concentration in edible fish (Figure 3).

Of the analyzed carnivore specimens, four exhibited total Hg levels exceeding the acceptable level (Figure 4), and *Lethrinus erythracanthus* had levels 3-times higher than the maximum permissible mercury concentration in edible fish.

In the herbivore category, *Valamugil seheli* tended to accumulate more mercury in their tissue than other species (Figure 5). In addition, no specimen in the omnivore category contained high mercury (Figure 6).

Taliwang

In the Taliwang area, 216 marine and fresh water fish specimen were collected, including *Lutjanus campechanus*, *Lethrinus erythracanthus*, *Cephalopoeis sp.*, *Siganus sp.*, *Loligo indica*, *Sardinella lemuru*, *Scaridae sp.*, *Siganus virgatus*, *Cheilinus trilobatus*, and *Valamugil seheli*. Freshwater fish collected from this area included *Channa striata*, *Oreochromis mossambikus*, *Anabas testudineus*.

There were no significant differences between fish length, weight and mercury concentration ($p > 0.05$, one-way analysis of variance) (Table 2).

Among the carnivorous fish collected in Taliwang, only *Cheilinus trilobatus* showed a high concentration of mercury, twice the maximum permissible mercury concentration for edible fish (Figure 7).

No samples in the omnivore category of marine or freshwater fish (Figure 8) or herbivore (Figure 9) category contained high concentrations of mercury.

Discussion

Fish and seafood are important dietary constituents because they are a high quality source of protein, vitamins, minerals and omega-3 fatty acids. However, fish have become a major source of methylmercury intake in humans. The main concern is that mercury content in fish can be biomagnified rapidly depending on the environmental parameters in water and sediment. Although the mercury concentration in water bodies in these areas was not measured in the present study, ASGM activities have been recognized as a key factor contributing mercury emissions to the aquatic system. In general, more than 90% of the mercury in fish occurs as methylmercury, but methylmercury contents can vary considerably between species. In aquatic systems, mercury is transformed into methylmercury by microorganisms and abiotic reactions. Methylmercury becomes increasingly concentrated in the marine food chain, in a process referred to as biomagnification, and can reach extremely high levels in predatory fish such as swordfish, tuna, king mackerel and shark. As a result, the risk of human exposure to methylmercury through consumption of large predator fish may be elevated. In the sampling sites in the present study, however, large predator fish are not often part of the catch of traditional fisherman. The small size of fish collected in the present study represents the fish most commonly consumed in this area.

Around 10% of the fish samples from Sekotong contained very high methylmercury concentrations. These results indicate that bioaccumulation of mercury had occurred in fish in Sekotong, particularly for fish in the planktivore and carnivore categories. In the present study, the mercury concentration in *Pilsbryoconcha*




Figure 5 — Mean levels of total mercury in herbivorous fish at Sekotong against the maximum permissible mercury concentration in edible fish tissue of 500 ppb




Figure 6 — Mean levels of total mercury in omnivorous fish at Sekotong

Tropic category	Species	No. of specimens (n)	Weight (g)	Length (cm)
Planktivore	<i>Sardinella lemuru</i>	22	32.40±7.09	13.1±1.14
Herbivore	<i>Cephalophoelis mimiata</i>	3	256±0.00	21.5±0.50
	<i>Channa striata</i> *	2	395.50±7.77	30.75±0.35
	<i>Cheilinus trilobatus</i>	3	43.66±0.57	23.16±0.28
	<i>Epinephelus sp.</i>	8	233.33±74.93	20.07±2.61
	<i>Lutjanus sp.</i>	5	375.66±71.06	24.00±1.00
	<i>Lethrinus erythracanthus</i>	3	396.66±32.14	22.00±2.00
	<i>Loligo indica</i>	30	43.5±7.55	17.25±6.08
	<i>Terapon jarbua</i>	4	25.00±1.42	8.70±0.53
Carnivore	<i>Chanos chanos</i>	3	20.66±2.08	10.52±0.85
	<i>Rasbora hengeli</i>	9	12.02±0.36	7.97±0.28
	<i>Siganus sp.</i>	11	180.66±79.95	18.16±3.51
	<i>Scaridae globiceps</i>	6	330.33±95.11	21.00±2.00
	<i>Valamugil seheli</i>	7	22.62±7.59	10.2±1.56
Omnivore	<i>Anabas testudineus</i> *	10	61.20±4.92	11.70±0.97
	<i>Geres punctatus</i>	36	15.98±0.27	7.96±0.17
	<i>Leiognathus equulus</i>	25	10.75±0.46	6.58±0.46
	<i>Oreochromis mossambikus</i> *	11	118.67±63.85	15.5±2.78
	<i>Silago sihama</i>	2	52.50±3.53	16.60±0.85

*freshwater fish

Table 2 — Characteristics of Fish Sampled at Taliwang




Figure 7 — Mean levels of total mercury in fish from the planktivore and carnivore categories at Taliwang against the maximum permissible mercury concentration in edible fish tissue of 500 ppb

exilis (shell, locally *kijing* (596 ppb) (Figure 3) was above the permissible limits for human consumption which is 0.5 µg Hg g.¹¹ The habitat for this organism is mud/sediment that may possibly contain mercury from mine waste discharged to rivers, while biotic demethylation reactions usually occur in sediment. This indicates that mercury contamination may have reached the estuary where this organism is found. Attention should be paid to the consumption of *Euthynnus affinis* (15) small tuna, locally *tongkol*, as 50% of the samples in the present study had high levels of methylmercury in their tissue. In addition, *Sephia officinalis* (squid, locally *cumi*) had a methylmercury concentration of 721 ppb (Figure 4). Even though *Euthynnus affinis* (17) is not a demersal fish and occurs

in open waters, it remains close to the shoreline, and its young may enter bays and harbors for feeding. ¹¹ shown in Table 1, even though there was no statistically significant relationship between mercury concentration, size and length of fish, there was a tendency for higher mercury concentrations in fish tissue in bigger fish (around 20 cm), such as *Scolopis affinis* (24 cm), *Euthynnus affinis* (21.67 cm), and *Lethrinus erythracanthus* (18 cm). Most of the fish samples were small.

Results of the Taliwang samples suggest that marine and freshwater fish in this area are below the permissible limits for consumption by the community. In the Taliwang samples, only one marine fish from the carnivore category had a high mercury concentration, *Cheilinus trilobatus* (1063 ppb) (Figure 7). Attention should be paid to the consumption of this species, as it may pose a health risk. This fish is in the predator category and occasionally feeds in seagrass beds, mainly on benthic invertebrates such as mollusks and crustaceans. Unfortunately, no mollusks and crustaceans were sampled from this area to support this finding. Taliwang residents are concerned about contaminated freshwater fish from Lake Taliwang, due to the many gold mining and processing activities around Lake Taliwang. However, the results showed no evidence of mercury contamination in freshwater fish from Lake Taliwang, even for large fish such as *Channa striata* (Figure 7) or *Oreochromis mossambikus* and *Anabas testudineus* (Figure 8) (30.75, 15.5 and 11.70 cm), respectively. These findings could be due to differences in fish habitat use within the lake and growth rates as a function of lake size.²¹ These three species are the most commonly consumed species in Taliwang.




Figure 8 — Mean levels of total mercury in omnivorous fish at Taliwang




Figure 9 — Mean levels of total mercury in herbivorous fish at Taliwang

Consumption of fish provides excellent nutrition, but fish may contain levels of mercury above the WHO permissible limits for human consumption.¹¹ Sensitive groups in the population can best protect their health by avoiding fish species that may contain high levels of mercury. Vulnerable groups like pregnant women, breast feeding mothers and children up 3 years of

age should be advised not to consume potentially contaminated fish. Frequent consumption of seafood can increase the risk of mercury exposure, which is potentially dangerous to vulnerable groups. Even though rates of fish consumption were not measured in the present study, the authors observed that the average daily fish intake in these areas was high,

as fish are a main dietary component in the communities of Sekotong and Taliwang. It was suggested that the most at risk group are women of childbearing age and children who should keep mercury intake below 0.2 ug/kg body weight per day. They should eat locally caught fish only every second day. It is a limitation of this study that no fish were sampled from control sites, therefore it is not possible to definitively say that the elevated mercury levels were due to ASGM activity in the area, although it is probable. Miners and the surrounding community have become aware of the risks associated with mercury exposure and mining activities, but programs are needed to increase awareness and to address community concern over the risks associated with mercury exposure and fish consumption.

Conclusions

A few fish species from the sites in the present study had mercury concentrations above the maximum WHO permissible mercury concentrations in edible fish tissue. The risks associated with mercury exposure and fish consumption threaten the health of residents of these communities.

Acknowledgements

This research was funded by the European Union in partnership with UNIDO entitled via the Pure Earth "Mitigating Toxic Health Exposures in Low and Middle Income Countries" project. UNIDO Contract No. 3000035403, SAP ID No. 150416. Part of the European Commission's Global and Public Good Challenge. The views expressed herein can in no way be taken to reflect the official opinion of the European Union.

Copyright Policy

This is an Open Access article

distributed in accordance with Creative Commons Attribution License (<http://creativecommons.org/licenses/by/3.0/>).

References

- Reichert-Brushett AJ, Stone J, Howe P, Thomas B, Clark M, Male Y, Nanlohy A, Butcher P. Geochemistry and mercury contamination in receiving environments of artisanal mining wastes and identified concerns for food safety. *Environ Res* [Internet]. 2017 Jan [cited 2019 Jan 30];152:407-18. Available from: <https://doi.org/10.1016/j.envres.2016.07.007> Subscription required to view.
- Alpers CN, Yee JL, Ackerman JT, Orlando JL, Slotton DG, Marvin-DiPasquale MC. Prediction of fish and sediment mercury in streams using landscape variables and historical mining. *Sci Total Environ* [Internet]. 2016 Nov 15 [cited 2019 Jan 30];571:364-79. Available from: <https://doi.org/10.1016/j.scitotenv.2016.05.088> Subscription required to view.
- Faial K, Deus R, Deus S, Neves R, Jesus I, Santos E, Alves CN, Brasil D. Mercury levels assessment in hair of riverside inhabitants of the Tapajós River, Pará State, Amazon, Brazil: fish consumption as a possible route of exposure. *J Trace Elem Med Biol* [Internet]. 2015 Apr [cited 2019 Jan 30];30:66-76. Available from: <https://doi.org/10.1016/j.jtemb.2014.10.009> Subscription required to view.
- Castilhos ZC, Rodrigues-Filho S, Rodrigues AP, Villas-Boas RC, Siegel S, Veiga MM, Beinhoff C. Mercury contamination in fish from gold mining areas in Indonesia and human health risk assessment. *Sci Total Environ* [Internet]. 2006 Sep 1 [cited 2019 Jan 30];368(1):320-5. Available from: <https://doi.org/10.1016/j.scitotenv.2006.01.039> Subscription required to view.
- Kambey JL, Farrell AP, Bendell-Young LI. Influence of illegal gold mining on mercury levels in fish of north Sulawesi's Minahasa Peninsula, (Indonesia). *Environ Pollut* [Internet]. 2001 [cited 2019 Jan 30];114(3):299-302. Available from: [https://doi.org/10.1016/S0269-7491\(01\)00163-4](https://doi.org/10.1016/S0269-7491(01)00163-4) Subscription required to view.
- Krisnayanti BD, Anderson CW, Utomo WH, Feng X, Handayanto E, Mudarisna N, Ikram H, Khususiah. Assessment of environmental mercury discharge at a four-year-old artisanal gold mining area on Lombok Island, Indonesia. *J Environ Monit* [Internet]. 2012 Oct 26 [cited 2019 Jan 30];14(10):2598-607. Available from: <https://doi.org/10.1039/c2em30515a> Subscription required to view.
- Moraes PM, Santos EA, Cavecci B, Padilha CC, Vieira JC, Roldan PS, Padilha PM. GFAAS determination of mercury in muscle samples of fish from Amazon, Brazil. *Food Chem* [Internet]. 2013 Dec 1 [cited 2019 Jan 30];141(3):2614-7. Available from: <https://doi.org/10.1016/j.foodchem.2013.05.008>
- Liu JL, Xu XR, Yu S, Cheng H, Peng JX, Hong YG, Feng XB. Mercury contamination in fish and human hair from Hainan Island, South China Sea: implication for human exposure. *Environ Res* [Internet]. 2014 Nov [cited 2019 Jan 30];135:42-7. Available from: <https://doi.org/10.1016/j.envres.2014.08.023> Subscription required to view.
- Li P, Feng X, Qiu G. Methylmercury exposure and health effects from rice and fish consumption: a review. *Int J Environ Res Public Health* [Internet]. 2010 Jun [cited 2019 Jan 30];7(6):2666-91. Available from: <https://doi.org/10.3390/ijerph7062666>
- Mol JH, Ramlal JS, Lietar C, Verloo M. Mercury contamination in freshwater, estuarine, and marine fishes in relation to small-scale gold mining in Suriname, South America. *Environ Res* [Internet]. 2001 Jun [cited 2019 Jan 31];86(2):183-97. Available from: <https://doi.org/10.1006/enrs.2001.4256> Subscription required to view.
- Environmental health criteria 101: methylmercury [Internet]. Geneva, Switzerland: World Health Organization; 1990 [cited 2019 Jan 31]. 148 p. Available from: http://apps.who.int/iris/bitstream/10665/38082/1/9241571012_eng.pdf
- Ekawanti A, Krisnayanti D. Effect of mercury exposure on renal function and hematological parameters among artisanal and small-scale gold miners at Sekotong, West Lombok, Indonesia. *J Health Pollut* [Internet]. 2015 Dec [cited 2019 Jan 31];5(9):25-32. Available from: <https://doi.org/10.5696/2156-9614-5-9-25>
- Franko A, Budihna MV, Dodic-Fikfak M. Long-term effects of elemental mercury on renal function in miners of the Idrija Mercury Mine. *Ann Occup Hyg* [Internet]. 2005 Aug [cited 2015 Oct 28];49(6):521-7. Available from: <https://doi.org/10.1093/annhyg/mei022>
- Krisnayanti BD. ASGM status in West Nusa Tenggara Province, Indonesia. *J Degraded Min Lands Manag* [Internet]. 2018 [cited 2019 Jan 31];5(2):1077-84. Available from: <https://doi.org/10.15243/jdmlm.2018.052.1077>
- Krisnayanti BD, Vasurra I, Asmara MD, Ekawanti A, Suheri H. Analysis of artisanal small-

- scale gold mining sector in West Sumbawa Regency, Indonesia. *J Health Pollut* [Internet]. 2016 Dec [cited 2019 Jan 31];6(12):26-33. Available from: <https://doi.org/10.5696/2156-9614-6.12.26>
16. **Hasanudin M.** Indonesian through flow. *Oceana*. 1998;23(2):1-9.
17. **Krisnayanti BD, Anderson CW.** Environmental impact assessment: illegal/informal gold mining in Lombok. Jakarta, Indonesia; GIZ-RED; 2012. 69 p.
18. Guideline levels for methylmercury in fish: CAC/GL 7-1991. Geneva, Switzerland: FAO/WHO Joint Publications; 1991 Jan.
19. International Organization for Standardization, ISO/IEC 17025 – Testing and Calibration Laboratories, <https://www.iso.org/home/standards/popular-standards/isoiec-17025-testing-and-calibra.html>. [Accessed 2019, Feb 12]
20. International Organization for Standardization, ISO Guide 34:2009, <https://www.iso.org/standard/50174.html> [Accessed 2019, Feb 12]
21. **Evans MS, Lockhart WL, Doetzel L, Low G, Muir D, Kidd K, Stephens G, Delaronde J.** Elevated mercury concentrations in fish in lakes in the Mackenzie River Basin: The role of physical, chemical, and biological factors. *Sci Total Environ* [Internet]. 2005 Dec 1 [cited 2019 Jan 31];351-352:479-500. Available from: <https://doi.org/10.1016/j.scitotenv.2004.12.086> Subscription required to view

Jurnal 2

ORIGINALITY REPORT

19%

SIMILARITY INDEX

12%

INTERNET SOURCES

13%

PUBLICATIONS

5%

STUDENT PAPERS

PRIMARY SOURCES

- 1 Travis B. Lewis, Pablo A. Garcia-Chevesich, Thomas R. Wildeman, Jonathan O. Sharp. "Changes in surface water quality from small-scale gold mining operations in the Surinamean rainforest", *Toxicological & Environmental Chemistry*, 2020
Publication 2%
- 2 Baiq Dewi Krisnayanti, Christopher W. N. Anderson, Wani Hadi Utomo, Xinbin Feng et al. "Assessment of environmental mercury discharge at a four-year-old artisanal gold mining area on Lombok Island, Indonesia", *Journal of Environmental Monitoring*, 2012
Publication 2%
- 3 www.mysciencework.com
Internet Source 1%
- 4 D Rahmawati, J S Adiansyah. "An overview of remediation technology for mercury-contaminated sediment in Sekotong Sub District, Lombok, Indonesia", *IOP Conference Series: Earth and Environmental Science*, 2020
Publication 1%

5	watermark.silverchair.com Internet Source	1 %
6	www.crawfordscientific.com Internet Source	1 %
7	M.S. Evans, W.L. Lockhart, L. Doetzel, G. Low, D. Muir, K. Kidd, G. Stephens, J. Delaronde. "Elevated mercury concentrations in fish in lakes in the Mackenzie River Basin: The role of physical, chemical, and biological factors", Science of The Total Environment, 2005 Publication	1 %
8	Joice L Kambey, A.P Farrell, L.I Bendell-Young. "Influence of illegal gold mining on mercury levels in fish of North Sulawesi's Minahasa Peninsula, (Indonesia)", Environmental Pollution, 2001 Publication	1 %
9	Submitted to Universitat Internacional de Catalunya Student Paper	1 %
10	www.biomedres.us Internet Source	1 %
11	link.springer.com Internet Source	1 %
12	healthpollutionjournal.org Internet Source	1 %

13	deleerhulpjuf.nl Internet Source	1 %
14	mafiadoc.com Internet Source	1 %
15	Chonghuan Xia, Xiaoguo Wu, James C.W. Lam, Zhouqing Xie, Paul K.S. Lam. "Methylmercury and trace elements in the marine fish from coasts of East China", Journal of Environmental Science and Health, Part A, 2013 Publication	<1 %
16	Submitted to Rutgers University, New Brunswick Student Paper	<1 %
17	www.fishbase.se Internet Source	<1 %
18	D. E. Gaskin, K. I. Stonefield, P. Suda, R. Frank. "Changes in mercury levels in harbor porpoises from the Bay of Fundy, Canada, and adjacent waters during 1969-1977", Archives of Environmental Contamination and Toxicology, 1979 Publication	<1 %
19	Narjes Okati, Abbas Esmaili-sari. "Hair mercury and risk assessment for consumption of contaminated seafood in residents from the coast of the Persian Gulf,	<1 %

Iran", Environmental Science and Pollution Research, 2017

Publication

20	nrronline.org Internet Source	<1 %
21	Berzas Nevado, J.J.. "Mercury in the Tapajos River basin, Brazilian Amazon: A review", Environment International, 201008 Publication	<1 %
22	komw.net Internet Source	<1 %
23	eprints.mdx.ac.uk Internet Source	<1 %
24	univendspace.univen.ac.za Internet Source	<1 %
25	www.mdpi.com Internet Source	<1 %

Exclude quotes On

Exclude matches < 10 words

Exclude bibliography On