

1. Variabilities

by Andre Scabra

Submission date: 31-Jan-2023 08:52PM (UTC-0600)

Submission ID: 2003785707

File name: angroves_in_Gili_Meno_Lake,_North_Lombok_District,_Indonesia.pdf (3.07M)

Word count: 8111

Character count: 41448

ISSN: 1412-033X
E-ISSN: 2085-4722

BIODIVERSITAS

Journal of Biological Diversity

Volume 23 - Number 11 - November 2022



Front cover: Bubalus depressicornis (H. Smith, 1827)
(PHOTO: CHRIS K.)

Published monthly

PRINTED IN INDONESIA

ISSN: 1412-033X

E-ISSN: 2085-4722



9 771412 033931



9 772085 472935

BIODIVERSITAS

Journal of Biological Diversity
Volume 23 - Number 11 - November 2022

ISSN/E-ISSN:

1412-033X (printed edition), 2085-4722 (electronic)

EDITORIAL BOARD:

Abdel Fattah N.A. Rabou (Tunisia), **Agnieszka B. Najda** (Poland), **Ajay Kumar Gautam** (India), **Alan J. Lymbery** (Australia), **Annisa** (Indonesia), **Bambang H. Saharjo** (Indonesia), **Daiane H. Nunes** (Brazil), **Darlina Md. Naim** (Malaysia), **Ghulam Hassan Dar** (India), **Hassan Pourbabaei** (Iran), **Joko R. Witono** (Indonesia), **Kartika Dewi** (Indonesia), **Katsuhiko Kondo** (Japan), **Kusumadewi Sri Yulita** (Indonesia), **Livia Wanntorp** (Sweden), **M. Jayakara Bhandary** (India), **Mahdi Reyahi-Khoram** (Iran), **Mahendra K. Rai** (India), **Mahesh K. Adhikari** (Nepal), **Maria Panitsa** (Greece), **Mochamad A. Hendjoto** (Indonesia), **Mohamed M.M. Najim** (Srilanka), **Mohib Shah** (Pakistan), **Murhasanah** (Indonesia), **Praptiwi** (Indonesia), **Rasool B. Tareen** (Pakistan), **Seyed Aliakbar Hedayati** (Iran), **Seyed Mehdi Talebi** (Iran), **Shahabuddin** (Indonesia), **Shahir Shamsir** (Malaysia), **Shri Kant Tripathi** (India), **Subhash C. Santra** (India), **Sugeng Budiharta** (Indonesia), **Sugiyarto** (Indonesia), **Taufiq Purna Nugraha** (Indonesia), **Yosep S. Mau** (Indonesia)

EDITOR-IN-CHIEF:
Sutarno

EDITORIAL MEMBERS:

English Editors: **Graham Eagleton** (grahameagleton@gmail.com), **Suranto** (surantouns@gmail.com); Technical Editor: **Solichatun** (solichatun_s@yahoo.com), **Artini Pangastuti** (pangastuti_tutut@yahoo.co.id); Distribution & Marketing: **Rita Rakhmawati** (oktia@yahoo.com); Webmaster: **Ari Pitoyo** (aripitoyo@yahoo.com)

MANAGING EDITORS:
Ahmad Dwi Setyawan (unsjournals@gmail.com)

PUBLISHER:
The Society for Indonesian Biodiversity

CO-PUBLISHER:
Department of Biology, Faculty of Mathematics and Natural Sciences, Sebelas Maret University, Surakarta

ADDRESS:
Jl. Ir. Sutami 36A Surakarta 57126. Tel. +62-271-7994097, Tel. & Fax.: +62-271-663375, email: editors@smujo.id

ONLINE:
biodiversitas.mipa.uns.ac.id; smujo.id/biodiv



**Society for Indonesia
Biodiversity**



**Sebelas Maret University
Surakarta**

Published by Smujo International for The Society for Indonesian Biodiversity and Sebelas Maret University Surakarta

GUIDANCE FOR AUTHORS

Aims and Scope *Biodiversitas, Journal of Biological Diversity* or *Biodiversitas* encourage submission of manuscripts dealing with all aspects of biodiversity including plants, animals and microbes at the level of the gene, species, and ecosystem. Ethnobiology papers are also considered.

Article types The journal seeks original full-length: (1) **Research papers**, (2) **Reviews**, and (3) **Short communications**. Original research manuscripts are limited to 8,000 words (including tables and picture), or proportional with articles in this publication number. Review articles are also limited to 8,000 words, while Short communications should be less than 2,000 words, except for pre-study.

Submission The journal only accepts online submission, through open journal system (<https://smujo.id/biodiv/about/submissions>) or email to the editors at unsjournals@gmail.com. Submitted manuscripts should be the original works of the author(s). Please ensure that the manuscript is submitted using *Biodiversitas* template, which can be found at (<https://biodiversitas.mipa.uns.ac.id/D/guidance.htm>). The manuscript must be accompanied by a cover letter containing the article title, the first name and last name of all the authors, a paragraph describing the claimed novelty of the findings versus current knowledge. Please also provide a list of five potential reviewers in your cover letter. Submission of a manuscript implies that the submitted work has not been published before (except as part of a thesis or report, or abstract); and is not being considered for publication elsewhere. When a manuscript written by a group, all authors should read and approve the final version of the submitted manuscript and its revision; and agree the submission of manuscripts for this journal. All authors should have made substantial contributions to the concept and design of the research, acquisition of the data and its analysis; drafting of the manuscript and correcting of the revision. All authors must be responsible for the quality, accuracy, and ethics of the work.

Ethics Author(s) must obedient to the law and/or ethics in treating the object of research and pay attention to the legality of material sources and intellectual property rights.

Copyright If the manuscript is accepted for publication, the author(s) still hold the copyright and retain publishing rights without restrictions. Authors are allowed to reproduce articles as long as they are not used for commercial purposes. For the new invention, authors are suggested to manage its patent before published.

Open access The journal is committed to free-open access that does not charge readers or their institutions for access. Readers are entitled to read, download, copy, distribute, print, search, or link to the full texts of articles, as long as not for commercial purposes. The license type is CC-BY-NC-SA.

Acceptance Only articles written in U.S. English are accepted for publication. Manuscripts will be reviewed by editors and invited reviewers (double blind review) according to their disciplines. Authors will generally be notified of acceptance, rejection, or need for revision within 1 to 2 months of receipt. Manuscripts will be rejected if the content does not in line with the journal scope, does not meet the standard quality, is in an inappropriate format, contains complicated grammar, dishonesty (i.e. plagiarism, duplicate publications, fabrication of data, citations manipulation, etc.), or ignoring correspondence in three months. The primary criteria for publication are scientific quality and biological significance. **Uncorrected proofs** will be sent to the corresponding author by email as .doc or .docx files for checking and correcting of typographical errors. To avoid delay in publication, corrected proofs should be returned in 7 days. The accepted papers will be published online in a chronological order at any time, but printed in January, April, July and October.

A charge Starting on January 1, 2017, publishing costs waiver is granted to foreign (non-Indonesian) authors who first publish the manuscript in this journal, especially for graduate students from developing countries. However, other authors are charged USD 250 (IDR 3,500,000).

Reprints The sample journal reprint is only available by special request. Additional copies may be purchased when ordering by sending back the uncorrected proofs by email. Manuscript preparation Manuscript is typed on A4 (210x297 mm²) paper size, in a single column, single space, 10-point (10 pt) Times New Roman font. The margin text is 3 cm from the top, 2 cm from the bottom, and 1.8 cm from the left and right. Smaller lettering size can be applied in presenting table and figure (9 pt). Word processing program or additional software can be used, however, it must be PC compatible, use the *Biodiversitas* template, and Microsoft Word based (.doc or .rtf; not .docx).

Scientific names of species (incl. subspecies, variety, etc.) should be written in italics, except in italicised sentences. Scientific names (Genus, species, author), and cultivar or strain should be mentioned completely for the first time mentioning it in the body text, especially for taxonomic manuscripts. The Genus name can be shortened after first mention, except where this may generate confusion. Name of the author can be eliminated after first mentioning. For example, *Rhizopus oryzae* L. UICC 524, hereinafter can be written as *R. oryzae* UICC 524. Using trivial names should be avoided. **Biochemical and chemical nomenclature** should follow the order of the IUPAC - IUB. For DNA sequence, it is better used Courier New font. Standard chemical abbreviations can be applied for common and clear used, for example, completely written butilic hydroxyl toluene (BHT) to be BHT hereinafter. **Metric measurements** should use IS denominations, and other system should use equivalent values with the denomination of IS mentioned first. Abbreviations like g, mg, mL, etc. should not be followed by a dot.

Minus index (m-2, L-1, h-1) suggested to be used, except in things like "per-plant" or "per-plot". **Mathematical equations** can be written down in one column with text, in that case can be written separately. **Numbers** one to ten are written in words, except if it relates to measurement, while values above them written in number, except in early sentence. The fraction should be expressed in decimal. In the text, it should be used "%" rather than "percent". Avoid expressing ideas with complicated sentence and verbiage, and used efficient and effective sentence.

The **Title** of the article should be written in compact, clear, and informative sentence, preferably not more than 20 words. Author name(s) should be completely written. Name and institution address should also be completely written with street name and number (location), postal code, telephone number, facsimile number, and email address. Manuscripts written by a group, author for correspondence along with address is required. First page of the manuscript is used for writing above information.

The **Abstract** should not be more than 200 words. Include between five and eight **Keywords**, using both scientific and local names (if any), research themes, and special methods which used; and sorted from A to Z. All important **abbreviations** must be defined at their first mention. Running title is about five words. The **Introduction** is about 400-600 words, covering the background and aims of the research. **Materials and Methods** should emphasize on the procedures and data analysis. **Results and Discussion** should be written as a series of connecting sentences, however, for manuscript with long discussion should be divided into subtitles. Thorough discussion represents the causal effect mainly explains for why and how the results of the research were taken place, and do not only re-express the mentioned results in the form of sentences. A **Conclusion** should be given at the end of the discussion. **Acknowledgments** are expressed in brief; all sources of institutional, private and corporate financial support for the work must be fully acknowledged, and any potential conflicts of interest must be noted.

Figures and Tables of three pages maximum should be clearly presented. Include a label below each figure, and a label above each table (see example). Titled figures can only be accepted if the information in the manuscript can be understood without those images; chart is preferred to use black and white images. Author could consign any picture or photo for the front cover, although it does not print in the manuscript. All images property of others should be mentioned in the text. There is no **Appendix**, all data or data analysis are incorporated into Results and Discussions. For broad data, supplementary information can be provided on the website.

References In the text give the author names followed by the year of publication and arrange from oldest to newest and from A to Z. In citing an article written by two authors, both of them should be mentioned, however, for three and more authors only the first author is mentioned followed by et al., for example: Saharjo and Nurhayati (2006) or (Boonkerd 2003a, b, c; Sugiyarto 2004; El-Bana and Nijs 2005; Balagadde et al. 2008; Webb et al. 2008). Extent citation as shown with word "cit" should be avoided. Reference to unpublished data and personal communication should not appear in the list but should be cited in the text only (e.g., Rifai MA 2007, pers. com. (personal communication); Setyawan AD 2007, unpublished data). In the reference list, the references should be listed in an alphabetical order. Names of journals should be abbreviated. Always use the standard abbreviation of a journal's title according to the ISSN List of Title Word Abbreviations (www.issn.org/2-22661-LTWA-online.php). Please include DOI links for journal papers. The following examples are for guidance.

Journal:

Saharjo BH, Nurhayati AD. 2006. Domination and composition structure change at hemic peat natural regeneration following burning: a case study in Pelalawan, Riau Province. *Biodiversitas* 7: 154-158. DOI: 10.13057/biodiv/d070213

Book:

Rai MK, Carpinella C. 2006. Naturally Occurring Bioactive Compounds. Elsevier, Amsterdam.

Chapter in book:

Webb CO, Cannon CH, Davies SJ. 2008. Ecological organization, biogeography, and the phylogenetic structure of rainforest tree communities. In: Carson W, Schnitzer S (eds) *Tropical Forest Community Ecology*. Wiley-Blackwell, New York.

Abstract:

Assaeed AM. 2007. Seed production and dispersal of *Rhazya stricta*. 50th annual symposium of the International Association for Vegetation Science, Swansea, UK, 23-27 July 2007.

Proceeding:

Alikodra HS. 2000. Biodiversity for development of local autonomous government. In: Setyawan AD, Sutarno (eds.) *Toward Mount Lawu National Park; Proceeding of National Seminary and Workshop on Biodiversity Conservation to Protect and Save Germplasm in Java Island*. Universitas Sebelas Maret, Surakarta, 17-20 July 2000. [Indonesian]

Thesis, Dissertation:

Sugiyarto. 2004. Soil Macro-invertebrates Diversity and Inter-Cropping Plants Productivity in Agroforestry System based on Sengon. [Dissertation]. Universitas Brawijaya, Malang. [Indonesian]

Information from internet: Balagadde FK, Song H, Ozaki J, Collins CH, Bamet M, Arnold FH, Quake SR, You L. 2008. A synthetic *Escherichia coli* predator-prey ecosystem. *Mol Syst Biol* 4:187. www.molecularsystembiology.com

Variabilities of the carbon storage of mangroves in Gili Meno Lake, North Lombok District, Indonesia

SITTI HILYANA^{1,*}, FIRMAN ALI RAHMAN²

¹Departement of Marine Science, Faculty of Agriculture, Universitas Mataram. Jl. Majapahit No.62, Gomong, Selaparang, Mataram 83125, West Nusa Tenggara, Indonesia. Tel./fax.: +62-370-633007, *email: sittihilyana@unram.ac.id

²Department of Biology Education, Faculty of Education and Teacher Training, Universitas Islam Negeri Mataram. Jl. Pendidikan No. 35, Mataram 83127, West Nusa Tenggara, Indonesia

Manuscript received: 5 September 2022. Revision accepted: 10 November 2022.

Abstract. Hilyana S, Rahman FA. 2022. Variabilities of the carbon storage of mangroves in Gili Meno Lake, North Lombok District, Indonesia. *Biodiversitas* 23: 5862-5868. Mangrove is one of the coastal vegetation that can act as carbon mitigation (carbon sink and carbon storage). This study aims to determine the potential for carbon sinks and storage in the leaves and roots of each type of mangrove found in Gili Meno lake, North Lombok, Indonesia. The research includes the identification of species and sampling (leaves and roots) of mangroves in the research quadrant. The organic carbon content of mangrove leaves and roots was tested using the Wakley and Black method. The results showed that there were 5 (five) types of mangroves in Gili Meno lake, namely: *Avicennia marina*, *Lumnitzera racemosa*, *Bruguiera cylindrica*, *Rhizophora apiculata*, and *Excoecaria agallocha*. The highest leaf tissue carbon content value was *R. apiculata* at 45.85%C or equivalent to 3.19 g.C, while in roots, *A. marina* was 50.06%C, equivalent to 4.49 g.C. In addition, the potential carbon stock in the leaves of the entire mangrove ecosystem in an area of 3 ha is 762.81 tons.C \pm 199.257 and at the roots is 659.76 tons.C \pm 394.848, while the largest potential carbon stock in leaf organs is the type of mangrove *R. apiculata*, which is 318.91 tons.C.ha⁻¹. and at the root is the type of mangrove *A. marina*, amounting to 448.54 tons.C.ha⁻¹. The estimated carbon dioxide uptake by the Gili Meno mangrove leaves is in the range of 130.36 g.CO₂-168.27 g.CO₂ or with an average of 154.34 g.CO₂ \pm 14.376, while the species with the highest carbon dioxide absorption capacity is *R. apiculata* (268.27 g.CO₂) and the lowest in the species of *L. racemosa* (130.36 g.CO₂).

Keywords: Carbon dioxide, carbon stores, mangroves

INTRODUCTION

Gili Meno is a small island in North Lombok District, West Nusa Tenggara (NTB) Province. Geographically, Gili Meno is located between Gili Trawangan and Gili Air. One of the characteristics of Gili Meno among small islands in general in NTB or Indonesia is the presence of a saltwater lake located in the middle of the island. Gili Meno saltwater lake has an area of 6.6 ha with a diversity of biota (flora and fauna) and unique physical and chemical characteristics of the lake waters. One of them is extreme salinity conditions with an average of 54.00 \pm 0.82 ppt (Rahman and Hadi 2021), this condition is different in general in Indonesian marine waters, namely in the range of 33-43 ppt by the salinity quality standard based on the Indonesian Minister of Environment Decree No. 51 of 2004.

The uniqueness of extreme environmental parameters in the Gili Meno saltwater lake requires the biota that makes up the lake ecosystem to survive, one of which is the vegetation of various types of mangroves that grow around the lake with an area of \pm 3 ha. Mangrove vegetation that grows around the Gili Meno lake has various environmental services, namely as a buffer for the island ecosystem in its benefits environmental services such as carbon dioxide (CO₂) absorption, disaster mitigation (abrasion, coastal waves, sea breeze barriers, and

tsunamis), availability of clean air (O₂), stability of coastal waters, habitat for biota, mangrove ecotourism and germplasm (Aksornkoae and Kato 2011; Mcleod et al. 2011; Pendleton et al. 2012; Giri et al. 2015; Nordhaus et al. 2019; Rahman et al. 2020; Sadono et al. 2020; Alimbon and Manseguiao 2021a).

One of the important issues related to mangrove ecosystems is the study of mangrove ecology related to environmental services, namely the ability to absorb and store carbon below and above the surface (Estrada and Soares 2017; Taillardat et al. 2018; Widyastuti et al. 2018; Kusumaningtyas et al. 2019; Matatula et al. 2021). Various previous studies have proven that mangrove ecosystems have a greater carbon storage capacity than terrestrial forest and seagrass ecosystems, even though the world's mangrove forests only cover 0.2% of land vegetation cover (Hamilton and Casey 2016). Mangrove forest carbon storage can reach 6-8 tons.C.ha⁻¹.yr compared to land forest carbon storage capacity of 1.8-2.7 tons.C.ha⁻¹. and seagrass ecosystem with a storage capacity of 2-4 ton.C.ha⁻¹.yr⁻¹ (Murray et al. 2011). In addition, according to Murdiyarso et al. (2015) that the total carbon potential of Indonesian mangrove forests is around 3.14 Pg.C or globally of 69 million tonnes of carbon (Worthington and Spalding 2018).

Considering the importance of mangroves as a buffer ecosystem for the Gili Meno lake area, which has a role in

ecological and ecotourism services, it needs attention. The purpose of this study was to determine specifically the carbon content stored in each species found on Gili Meno in the leaves and roots of the mangroves so that they can be used as a reference source for the conservation of certain mangrove species that can absorb and store scattered carbon.

MATERIALS AND METHODS

The research was carried out in the Gili Meno lake ecosystem, Gili Indah Village, Pemenang Sub-district, North Lombok District, West Nusa Tenggara Province, Indonesia in July-August 2021 with a research area of 6.6 ha (Figure 1).

Research procedure

The study began with determining the point of making the quadrant followed by the process of identifying all types of mangroves contained in the quadrant. Next is the process of taking root and leaf samples for each type of mangrove. Samples of mangrove leaves and roots were taken randomly for each different species so that they could represent the same species in the same quadrant. Root samples were taken to a depth of 30 cm and included roots above the soil surface, such as the breath roots of *Avicennia marina*, *Bruguiera cylindrica*, and *Rhizophora apiculata*. All samples of leaves and roots of each type of mangrove were prepared as testing materials for biomass (wet weight

and dry weight) and tissue carbon at the Laboratory for the Study of Agricultural Technology in West Nusa Tenggara and the Soil Laboratory at the University of Mataram.

Identification of mangrove

Type Mangrove identification was carried out based on the morphological characteristics of mangrove species, such as leaf shape and color, fruit shape and color, flower shape and color, and root morphology with reference to reference to the Guide to Introduction to Mangroves in Indonesia.

Data analysis

Biomass of mangroves

The analysis of leaf and root biomass began with the addition of the wet weight of the sample and continued with the oven process at a temperature of 60°C until the dry weight of each sample became stable. Calculation of the analysis of the biomass of mangrove leaves and roots is carried out with the following formula:

$$\% \text{ Water content} = \frac{GW - DW}{GW} \times 100 \%$$

$$\text{Biomass} = \frac{GW}{1 + \frac{\% \text{ Water content}}{100}}$$

Where:

GW: Gross weight (g)

DW: Dry weight (g)




Figure 1. Research site in Gili Meno Lake of Gili Indah Village, Pemenang Sub-district, North Lombok District, West Nusa Tenggara Province, Indonesia

The organic carbon content (%) of mangroves

The organic carbon content of mangrove leaves and roots was carried out using the Wakley and Black method, weigh a minimum of 5 g of sample and put it in a 100 mL volumetric flask. Added 5 mL $K_2Cr_2O_7$ 1 N and then shaken. Added 7.5 mL of concentrated H_2SO_4 , shake, and let stand for 30 minutes. Diluted with ionized water, allow to cool and squeeze. The absorbance of the clear solution was measured by a spectrophotometer at a wavelength of 561 nm. The measurement results are then calculated by the formula:

The organic carbon content (%) = Curva ppm x mL extract 1.000 mL^{-1} x 100 mg sample $^{-1}$ x CF

= Curva ppm x 100 1.000^{-1} x 100 500^{-1} x CF

= Curva ppm x 10 500^{-1} x CF

Where:

Curva ppm : The sample rate obtained from the curve of the relationship between the standard series content and its reading after correcting for blanks

100 : Convert to %

CF/Corr. Factor : $100/(100 - \% \text{ moisture content})$

The carbon stock of mangroves

The carbon stock estimation of mangroves is calculated based on the biomass content and tissue carbon content of each mangrove species with the formula:

The carbon stock of mangroves (ton/ha) = Biomass x The organic carbon content (%)

Mangrove leaf CO_2

Absorption Estimation Carbon dioxide absorption in mangrove leaves is an estimate of the ability of mangroves in the photosynthesis process. This analysis was calculated based on Howard et al. (2014) as follows:

$$CO_2 \text{ absorption} = \frac{Mr\ CO_2}{Ar\ C} \times Cb$$

Where:

CO_2 absorption: Total carbon dioxide absorption (g/g-dry)

Mr CO_2 : Relative molecule

CO_2 : 44 (atomic mass C: 12, O: 16)

Ar C: Relative atoms (C: 12)

Cb: Mangrove carbon content (%)

RESULTS AND DISCUSSION**Results**

Based on the identification results, there are 5 (five) mangrove species found on Gili Meno, namely: *Avicennia*

marina, *Lumnitzera racemosa*, *Bruguiera cylindrica*, *Rhizophora apiculata*, and *Excoecaria agallocha*. Biomass is the result of the photosynthesis process, which is stored in every plant organ, such as leaves, stems, roots, fruits, and flowers. Based on the calculation results, the total leaf biomass content in 5 mangrove species is between 3.66 g-7.08 g with an average of 6.05 ± 1.448 . The highest biomass content in leaves was found in mangrove *E. agallocha* (7.08 g) and the lowest in *B. cylindrica* (3.66 g) (Table 1). In addition, the root biomass content of 5 (five) mangrove species was in the range of 2.82 g-8.96 g with an average of 4.88 ± 2.321 , i.e., the highest root biomass content was found in *A. marina* (8.96 g) and the lowest was at species of *R. apiculata* (2.82 g).

Percentage of carbon content of mangrove leaves and roots

Based on the results of laboratory analysis, it was found that the highest percentage of carbon content in the leaf tissue was *R. apiculata* (45.85% C) and the lowest was *L. racemosa* (35.53% C) (Table 2). Meanwhile, the highest percentage of carbon content in the root tissue was *A. marina* (50.06% C) and the lowest was *L. racemosa* (32.19% C) (Table 3).

Carbon content of mangrove leaves and roots

Carbon content in grams of carbon (g.C) is calculated based on biomass value and the content of % carbon contained in each organ under study. The results showed that the carbon content of mangrove leaves was in the range of 1.55 g C - 3.19 g.C or with an average of 2.54 g C ± 66.419 (Table 4). The highest leaf carbon content was found in the type of mangrove *R. apiculata* (3.19 g.C) and the lowest was in the type of *B. cylindrica* (1.55 g.C). Meanwhile, the carbon content of the roots was in the range of 1.26 g.C-4.49 g.C with an average of 2.20 g.C ± 1.316 . The highest root carbon content was *A. marina* (4.49 g.C) and the lowest was *R. apiculata* (1.26 g.C).

 CO_2 absorption of mangrove leaves and roots

The estimated carbon dioxide uptake in the leaf organs of the Gili Meno mangrove ecosystem is in the range of 130.36 g.CO_2 - 168.27 g.CO_2 or with an average of $154.34\text{ g.CO}_2 \pm 14.376$ (Table 6). The species with the highest adsorption capacity was found in the mangrove species *R. apiculata* (268.27 g.CO_2) and the lowest in the *L. racemosa* species (130.36 g.CO_2).

Table 1. The biomass of mangroves tissue

Sample	Mangrove tissue of leaf biomass				Mangrove tissue of root biomass			
	GW (g)	DW (g)	Water content (%)	Biomass (g)	GW (g)	DW (g)	Water content (%)	Biomass (g)
<i>Avicennia marina</i>	6.45	5.58	13.49	5.68	10.14	8.80	13.21	8.96
<i>Rhizophora apiculata</i>	7.95	6.80	14.42	6.95	3.14	2.79	11.30	2.82
<i>Excoecaria agallocha</i>	8.08	6.95	13.98	7.08	5.03	4.37	13.29	4.44
<i>Bruguiera cylindrica</i>	3.95	3.64	7.99	3.66	5.09	4.39	13.92	4.47
<i>Lumnitzera racemosa</i>	8.26	6.58	20.26	6.87	4.83	4.19	13.17	4.27
Average	6.94	5.91	14.029	6.05	5.65	4.91	12.98	4.99
Standard deviation	1.817	1.378	4.352	1.448	2.639	2.278	0.985	2.321

Table 2. The carbon content of leaves tissue

Sample	GW (g)	DW (g)	KL	FK	Abs	ppm Kurva	% C
<i>Avicennia marina</i>	6.45	5.58	15.60	116	0.30	184.39	42.63
<i>Rhizophora apiculata</i>	7.95	6.80	16.86	1.17	0.32	196.17	45.85
<i>Excoecaria agallocha</i>	8.08	6.95	16.26	1.16	0.31	189.35	44.03
<i>Bruguiera cylindrica</i>	3.95	3.64	8.69	1.09	0.32	194.31	42.24
<i>Lumnitzera racemosa</i>	8.26	6.58	25.40	1.25	0.23	141.61	35.52

Note: GW: Gross weight; DW: Dry weight; KL: Soil moisture content; FK: Correction factor; Abs: absorbance; %C: The percentage of carbon content

Table 3. The carbon content of roots tissue

Sample	GW (g)	DR (g)	KL	FK	Abs	ppm Kurva	% C
<i>Avicennia marina</i>	10.14	8.80	15.22	1.15	0.35	217.25	50.06
<i>Rhizophora apiculata</i>	3.14	2.79	12.75	1.13	0.32	198.03	44.66
<i>Excoecaria agallocha</i>	5.03	4.37	15.32	1.15	0.30	185.01	42.67
<i>Bruguiera cylindrica</i>	5.09	4.39	16.19	1.16	0.31	190.59	44.29
<i>Lumnitzera racemosa</i>	4.83	4.19	15.15	1.15	0.23	139.75	32.19

Note: GW: Gross weight; DW: Dry weight; KL: Soil moisture content; FK: Correction factor; Abs: absorbance; %C: The percentage of carbon content

Table 4. Carbon content in leaf and roots of mangrove

Sample	Leaves carbon stored			Roots carbon stored		
	Biomass (g)	% C	Carbon stored (g.C)	Biomass (g)	% C	Carbon stored (g.C)
<i>Avicennia marina</i>	5.68	42.63	2.42	8.96	50.06	4.49
<i>Rhizophora apiculata</i>	6.95	45.85	3.19	2.82	44.66	1.26
<i>Excoecaria agallocha</i>	7.08	44.03	3.12	4.44	42.67	1.90
<i>Bruguiera cylindrica</i>	3.66	42.24	1.55	4.47	44.29	1.98
<i>Lumnitzera racemosa</i>	6.87	35.52	2.44	4.27	32.19	1.37
Average	6.05	42.05	2.54	2.54	42.77	2.20
Standard deviation	1.448	3.917	0.664	0.664	6.538	1.316

Table 5. The carbon stock area

Sample	Leaves carbon stored		Roots carbon stored	
	Carbon stored (g.C)	Carbon stored (ton.C.ha ⁻¹)	Carbon stored (g.C)	Carbon stored (ton.C.ha ⁻¹)
<i>Avicennia marina</i>	2.42	242.27	4.49	448.54
<i>Rhizophora apiculata</i>	3.19	318.66	1.26	126.03
<i>Excoecaria agallocha</i>	3.12	311.91	1.90	189.58
<i>Bruguiera cylindrica</i>	1.55	154.64	1.98	198.06
<i>Lumnitzera racemosa</i>	2.44	243.88	1.37	137.39
Average	2.54	254.27	2.20	219.92
Standard deviation	0.664	66.419	1.316	131.616

Table 6. CO₂ Absorption of mangrove leaves and roots

Sample	Leaves carbon absorption		
	% C	Mr CO ₂ / Ar C	Carbon absorption (g.CO ₂)
<i>Avicennia marina</i>	42.63	3.67	156.45
<i>Rhizophora apiculata</i>	45.85	3.67	168.27
<i>Excoecaria agallocha</i>	44.03	3.67	161.59
<i>Bruguiera cylindrica</i>	42.24	3.67	155.02
<i>Lumnitzera racemosa</i>	35.52	3.67	130.36
Average	42.05	3.67	154.34
Standard deviation	3.917	0.000	14.376

Discussion

Based on the identification results, there are 5 (five) mangrove species found on Gili Meno, namely: *Avicennia marina*, *Lumnitzera racemosa*, *Bruguiera cylindrica*, *Rhizophora apiculata*, and *Excoecaria agallocha*. The results of this study were less than the 12 species of mangrove vegetation in Gerupuk Bay (*Rhizophora apiculata*, *Rhizophora stylosa*, *Rhizophora mucronata*, *Bruguiera gymnorrhiza*, *Ceriops decandra*, *Sonneratia alba*, *Avicennia marina*, *Avicennia lanata*, *Aegiceras corniculatum*, *Osbornia octodonta*, *Lumnitzera racemosa* and *Xylocarpus moluccensis*), while in Sereweh Bay there were 13 species *Avicennia lanata*, *Lumnitzera racemosa*, *Excoecaria agallocha*, *Pemphis acidula*, *Bruguiera gymnorrhiza*, *Ceriops decandra*, *Ceriops tagal*, *Rhizophora apiculata*, *Rhizophora mucronata*, *Rhizophora stylosa*, *Scyphiphora hydrophyllacea*, *Sonneratia alba*, and *Sonneratia caseolaris* (Rahman et al. 2020). The small number of mangrove species found on Gili Meno can be caused by environmental factors that are quite extreme, namely high levels of salinity in the waters with an average of 54 ± 0.82 ppt (Rahman and Hadi 2021).

Several things that can affect the mangrove biomass content of Gili Meno Lake are the condition of the chemical parameters of the waters, especially in conditions of quite extreme salinity (Rahman and Hadi 2021), so that it can affect the physiology and morphology of mangroves, especially the photosynthesis mechanism as part of the CO_2 binding process and plant biomass production. This is reinforced by the report of Shannon (1999) that extreme salinity conditions can affect the decrease in leaf area, leaf number, root thinning, and affect root growth. In addition, the osmotic effect of salinity can cause a decrease in plant growth rate, changes in leaf color, and a decrease in the development of the root/shoot ratio. This is by the report of Zhang et al. (2016); Tam et al. (2009); Abdelhakeem et al. (2016); Barreto et al. (2016); Hilmi et al. (2017); Shiau et al. (2017); Hilmi et al. (2019) that salinity, phosphate, soil nitrate, fertility, pH, and temperature can affect the growth rate.

In addition, the total biomass of an area can be influenced by vegetation characteristics, including vegetation strata (trees, poles, saplings, and seedlings), species density, species dominance, and leaf cover (Rahman et al. 2018), such as the high leaf biomass content of mangrove *E. agallocha* could be caused by leaf samples taken from the tree strata, while the type of *B. cylindrica* (3.66 g) was still in the sapling strata. In addition, Sheil et al. (2017) and Scales and Friess (2019) have that the biomass content of a mangrove species is influenced by the diameter of the trunk. The total biomass of the Gili Meno lake mangrove ecosystem is 110.42 tons.ha⁻¹ or the equivalent of 331.26 tons.C in an area of 3 ha of the Gili Meno mangrove ecosystem. The total biomass of the Gili Meno lake ecosystem is still lower than the mangrove forest biomass of Alas Purwo National Park at 438.79 tons.ha⁻¹ (equivalent to 219.53 tons.C.ha⁻¹ or 805.68 tons.CO₂.ha⁻¹); and the mangrove biomass of Dukuh Tapak, Semarang city of 1507.91 tons.ha⁻¹ (Irsadi et al. 2017). However, it is greater than the biomass of mangrove

forests on Kemujan Island, Karimunjawa National Park, which is 91.31 tons.C and mangrove forest biomass in Bandar Baru Dumai area of 78.6 tons.ha⁻¹ or equivalent to 39.3 tons.C.ha⁻¹ (Mandari et al. 2016).

The high percentage of carbon content stored in the mangrove of *R. apiculata* mangrove could be caused by the thick leaf morphology and wider leaf cross-section compared to the other 4 species in Gili Meno lake. This refers to the report by Hairiah and Rahayu (2007) that the carbon content of mangroves contained in the biomass is 46-50%. Meanwhile, the leaf carbon content of *L. racemosa* (35.53% C) has a low value due to the smaller leaf cross-sectional area with the largest percentage of water content compared to 4 other types of mangrove, namely 20.262% C.

The percentage of carbon content on the roots and leaves of *L. racemosa* both had the lowest values, this could be due to the relatively thin morphology of the roots of *L. racemosa* with a root diameter of $\pm 0.2-0.5$ cm. It was different with the root morphology of *A. marina* (50.06% C) and *R. apiculata* (44.66% C), and *B. cylindrica* (44.29% C) with larger diameters and root volumes.

The carbon content contained in the leaves of *R. apiculata* (3.19 g.C) with the largest content was not correlated with the low carbon content of the roots (1.26 g.C), this was inversely proportional to the carbon content of the mangrove *A. marina* which was greater in the roots (4.49 g.C) compared to leaves (2.42 g.C). The high and low carbon content in the Gili Meno mangrove organs is influenced by the percentage of water content, one of which is the low carbon content of the roots of *R. apiculata* (2.82 g) caused by the samples taken in the form of breath roots that are still relatively young and are always flooded by water, so that when After oven drying the sample, the lowest dry weight result (2.79 g) with the difference between wet rice and dry weight was 0.355 g, this will affect the carbon content results even though *R. apiculata* mangrove has the second highest % carbon content of root tissue among the mangrove root samples, Gili Meno lake. Overall, the average carbon content of the roots (2.20 g.C \pm 1.316) was lower than that of the leaves (2.54 g.C \pm 0.664) with a ratio of 1:1.156.

Mangroves are one of the vegetation with the largest potential carbon stock, this is supported by the report of Sad et al. (2010); Lunstrum and Chen (2014); Matsui et al. (2015); Dahl et al. (2016); Wang et al. (2016); and Nyanga (2020) that mangrove forests can store carbon three to four times greater than forests on land because the organic matter contained in mangrove ecosystems sinks and is stored in the substrate, in contrast to terrestrial forest ecosystems which can easily release carbon through mechanisms. Weathering, combustion, and source of food for decomposer organisms. Several studies have observed the factors that influence carbon conservation in mangrove ecosystems, namely Matsui et al. (2015); Weiss et al. (2016); Martuti et al. (2017); Asadi et al. (2018); Perez et al. (2018); Gao et al. (2019); and Kida and Fujitake (2020).

Alongi et al. (2016) reported that the average carbon stock of mangroves in Indonesia is around 950 Mg.C.ha⁻¹ with details of the soil carbon stock at 774.7 Mg.C.ha⁻¹,

25
 above ground at 159.1 Mg.C.ha⁻¹ and below ground at 16.7 Mg.C.ha⁻¹. Meanwhile, the average mangrove forest carbon in the Kelantan Delta of Peninsular Malaysia is 156.35 Mg.C.ha⁻¹ (Rozainah et al. 2018); and mangrove carbon stock in Honda Bay, the Philippines at 47.9 Mg.C.ha⁻¹ (Castillo et al. 2018). While overall, the average carbon stock in the 3 ha area of the Gili Meno mangrove ecosystem was 762.81 tons.C±199.257 on the leaves and 659.76 tons.C±394.848 on the roots. The largest potential carbon stock in leaf organs is *R. apiculata* mangrove, which is 318.91 tons.C.ha⁻¹ and at the roots is *A. marina* mangrove at 448.54 tons.C.ha⁻¹ (Table 5), this is by field conditions that one of the dominant species in Gili Meno is *A. marina*.

The ability of carbon storage in mangrove type *A. marina* in Gili Meno lake has the same ability as the results of research by Kathiresan et al. (2013) in the mangrove ecosystem of the South Coast of India, which was 75% higher than *R. mucronata*, and while Alimbon and Manseguiao (2021b) also found that aboveground carbon stocks of *A. marina* were higher than *R. mucronata* and *S. alba*. Likewise, the research results of Purwanto et al. (2021) showed that *A. marina* trees in the Pangarengan mangrove forest store the highest amount of carbon than *A. alba*, *R. mucronata*, and *S. caseolaris* species.

Amount estimates of carbon stored in living plants (biomass) can reflect the CO₂ absorbed by plants from the atmosphere (Saderne et al. 2019). The ability of carbon absorption in each type of mangrove can be influenced by the age of the mangrove species, leaf cross-sectional area, water physicochemical factors, and the morphology of the mangrove strata.

Based on the research, it can be concluded that the high and low carbon content in the mangrove species of Gili Meno lake can be influenced by the biomass content, species strata, and water chemistry factors.

REFERENCES

- Abdelhakeem G, Abouloos SA, Kamel MM. 2016. Performance of a vertical subsurface flow constructed wetland under different operational conditions. *J Adv Res* 7 (5): 803-814. DOI: 10.1016/j.jare.2015.12.002.
- Aksornkoae S, Kato S. 2011. Mangroves for the people and environmental conservation in Asia. *Bull Soc Sea Water Sci* 65 (1): 3-9.
- Alimbon JA, Manseguiao MRS. 2021a. Community knowledge and utilization of mangroves in Panabo Mangrove Park, Panabo City, Davao del Norte, Philippines. *Intl J Bonorowo Wetl* 11 (2): 51-57. DOI: 10.13057/bonorowo/w110201.
- Alimbon JA, Manseguiao MRS. 2021b. Species composition, stand characteristics, aboveground biomass, and carbon stock of mangroves in Panabo Mangrove Park, Philippines. *Biodiversitas* 22 (6): 3130-3137. DOI: 10.13057/biodiv/d220615.
- Alongi DM, Murdiyoso D, Fourqurean JW, Kauffman JB, Hutahaean A, Crooks S, Wagey T. 2016. Indonesia's blue carbon: A globally significant and vulnerable sink for seagrass and mangrove carbon. *Wetl Ecol Manag* 24 (1): 3-13. DOI: 10.1007/s11273-015-9446-y.
- Asadi MA, Yona D, Saputro SE. 2018. Species diversity, biomass, and carbon stock assessments of mangrove forest in Labuhan, Indonesia. *IOP Conf Ser Earth Environ Sci* 151 (1): 012009. DOI: 10.1088/1755-1315/151/1/012009.
- Barreto MB, Lo Mónaco S, Díaz R, Barreto-Pittol E, López L, Peralba M, do CR. 2016. Soil organic carbon of mangrove forests (*Rhizophora* and *Avicennia*) of the Venezuelan Caribbean coast. *Org Geochem* 100: 51-61. DOI: 10.1016/j.orggeochem.2016.08.002.
- Castillo JAA, Apan AA, Maraseni TN, Salmó SG. 2018. Tree biomass quantity, carbon stock and canopy correlates in mangrove forest and land uses that replaced mangroves in Honda Bay, Philippines. *Reg Stud Mar Sci* 24: 174-183. DOI: 10.1016/j.rsma.2018.08.006.
- Dahl M, Deyanova D, Gütschow S, Asplund ME, Lyimo LD, Karamfilov V, Santos R, Björk M, Gullström M. 2016. Sediment properties as important predictors of carbon storage in *Zostera marina* meadows: A comparison of four European areas. *Plos One* 11 (12): e0167493. DOI: 10.1371/journal.pone.0167493.
- Estrada GCD, Soares MLG. 2017. Global patterns of aboveground carbon stock and sequestration in mangroves. *An Acad Bras Cienc* 89 (2): 973-989. DOI: 10.1590/0001-3765201720160357.
- Gao Y, Zhou J, Wang L, Guo J, Feng J, Wu H, Lin G. 2019. Distribution patterns and controlling factors for the soil organic carbon in four mangrove forests of China. *Glob Ecol Conserv* 17: e00575. DOI: 10.1016/j.gecco.2019.e00575.
- Giri C, Long J, Abbas S, Murali RM, Qamer FM, Pengra B, Thau D. 2015. Distribution and dynamics of mangrove forests of South Asia. *J Environ Manag* 148: 101-111. DOI: 10.1016/j.jenvman.2014.01.020.
- Hamilton SE, Casey D. 2016. Creation of a high spatio-temporal resolution global database of continuous mangrove forest cover for the 21st century (CGMFC-21). *Glob Ecol Biogeogr* 25 (6): 729-738. DOI: 10.1111/geb.12449.
- Hairiah K, Rahayu S. 2007. Pengukuran Karbon Tersimpan di Berbagai Macam Penggunaan Lahan. World Agroforestry Centre, ICRAFSA, Bogor.
- Hilmi E, Kusmana C, Suhendang E, Iskandar. 2017. Correlation analysis between seawater intrusion and mangrove greenbelt. *Indon J For Res* 4 (2): 151-168. DOI: 10.20886/ijfr.2017.4.2.151-168.
- Hilmi E, Kusmana C, Suhendang E, Iskandar. 2019. The carbon conservation of mangrove ecosystem in Indonesia. *Biotropia* 26 (3): 1-16. DOI: 10.11598/btb.2019.26.3.1099.
- Howard J, Hoyt S, Isensee K, Telszewski M, Pidgeon E. 2014. Coastal Blue Carbon: Methods for Assessing Carbon Stocks and Emissions Factors in Mangroves, Tidal Salt Marshes, and Seagrasses. Conservation International, Intergovernmental Oceanographic Commission of UNESCO, International Union for Conservation of Nature, Arlington, Virginia, USA.
- Irsadi A, Matuti NKT, Nugraha SB. 2017. Estimasi stok karbon mangrove di Dukuh Tapak Kelurahan Tugurejo Kota Semarang. *Jurnal Sainteknologi* 15 (2): 120-127. DOI: 10.15294/sainteknologi.v15i2.12402. [Indonesian]
- Kathiresan K, Anburaj R, Gomathi V, Saravanakumar K. 2013. Carbon sequestration potential of *Rhizophora mucronata* and *Avicennia marina* as influenced by age, season, growth and sediment characteristics in southeast coast of India. *J Coast Conserv* 17 (3): 397-408. DOI: 10.1007/s11852-013-0236-5.
- Kida M, Fujitake N. 2020. Organic carbon stabilization mechanisms in mangrove soils: A review. *Forests* 11 (9): 981. DOI: 10.3390/f11090981.
- Kusumaningtyas MA, Hutahaean AA, Fischer HW, Pérez-mayo M, Ransby D, Jennerjahn TC. 2019. Variability in the organic carbon stocks, sources, and accumulation rates of Indonesian mangrove ecosystems. *Estuar Coast Shelf Sci* 218: 310-323. DOI: 10.1016/j.ecss.2018.12.007.
- Lunstrum A, Chen L. 2014. Soil carbon stocks and accumulation in young mangrove forests. *Soil Biol Biochem* 75: 223-232. DOI: 10.1016/j.soilbio.2014.04.008.
- Mandari DZ, Gunawan H, Isda MN. 2016. Penaksiran biomassa dan karbon tersimpan pada ekosistem hutan mangrove di Kawasan Bandar Bakau Dumai. *Jurnal Riau Biologia* 1 (3): 17-23. [Indonesian]
- Martuti NKT, Setyowati DL, Nugraha SB, Mutiatari DP. 2017. Carbon stock potency of mangrove ecosystem at Tapak Sub-village, Semarang, Indonesia. *AACL Bioflux* 10 (6): 1524-1533.
- Matatula J, Afandi AY, Wirabuana PYAP. 2021. Short communication: Comparison of stand structure, species diversity and aboveground biomass between natural and planted mangroves in Sikka, East Nusa Tenggara, Indonesia. *Biodiversitas* 22 (3): 1098-1103. DOI: 10.13057/biodiv/d220303.
- Matsui N, Meepool W, Chukwamdee J. 2015. Soil organic carbon in mangrove ecosystems with different vegetation and sedimentological conditions. *J Mar Sci Eng* 3 (4): 1404-1424. DOI: 10.3390/jmse3041404.
- Mcleod E, Chmura GL, Bouillon S, Salm R, Björk M, Duarte CM, Lovelock CE, Schlesinger WH, Silliman BR. 2011. A blueprint for

- blue carbon: Toward an improved understanding of the role of vegetated coastal habitats in sequestering CO₂. *Front Ecol Environ* 9 (10): 552-560. DOI: 10.1890/110004.
- Murdiyasarso D, Purbopuspito J, Kauffman JB, Warren MW, Sasmito SD, Donato DC, Manuri S, Krisnawati H, Taberima S, Kurnianto S. 2015. The potential of Indonesian mangrove forests for global climate change mitigation. *Nat Clim Change* 5 (12): 1089-1092. DOI: 10.1038/nclimate2734.
- Murray BC, Pendleton L, Jenkins WA, Sifleet S. 2011. Green Payments for Blue Carbon: Economic Incentives for Protecting Threatened Coastal Habitats. Duke University, North Carolina, US.
- Nordhaus I, Toben M, Fauziyah A. 2019. Impact of deforestation on mangrove tree diversity, biomass and community dynamics in the Segara Anakan lagoon, Java, Indonesia: A ten-year perspective. *Estuar Coast Shelf Sci* 227: 106300. DOI: 10.1016/j.ecss.2019.106300.
- Nyanga C. 2020. The role of mangroves forests in decarbonizing the atmosphere. In: Bartoli M, Frediani M (eds). *Carbon-Based Material for Environmental Protection and Remediation*. IntechOpen, United Kingdom.
- Pendleton L, Donato DC, Murray BC, Crooks S, Aaron JW, Sifleet S, Craft C, Fourqurean JW, Kauffman JB, Marba N, Magonigal P, Pidgeon E, Herr D, Gordon D, Baldera A. 2012. Estimating global "blue carbon" emissions from conversion and degradation of vegetated coastal ecosystems. *Plos One* 7: e43542. DOI: 10.1371/journal.pone.0043542.
- Pérez A, Libardoni BG, Sanders CJ. 2018. Factors influencing organic carbon accumulation in mangrove ecosystems. *Biol Lett* 14 (10): 20180237. DOI: 10.1098/rsbl.2018.0237.
- Prasad MBK, Dittmar T, Ramanathan AL. 2010. Organic matter and mangrove productivity. In: Ramanathan AL, Bhattacharya P, Dittmar T, Prasad MBK, Neupane BR (eds). *Management and Sustainable Development of Coastal Zone Environments*. Springer, Dordrecht.
- Purwanto RH, Mulyana B, Sari PI, Hidayatullah MF, Marpaung AA, Putra ISR, Putra AG. 2021. The environmental services of Pangarengan mangrove forest in Cirebon, Indonesia: Conserving biodiversity and storing carbon. *Biodiversitas* 22 (12): 5609-5616. DOI: 10.13057/biodiv/d221246.
- Rahman FA, Hadi AP. 2021. Kandungan c-organik substrat ekosistem mangrove di Danau Air Asin Gili Meno Kabupaten Lombok Utara. *Jurnal Bioscientist: Jurnal Ilmiah Biologi* 9 (2): 516-526. DOI: 10.33394/bioscientist.v9i2.4276. [Indonesian]
- Rahman FA, Qayim I, Wardiatno Y. 2018. Carbon storage variability in seagrass meadows of Marine Poton Bako, East Lombok, West Nusa Tenggara, Indonesia. *Biodiversitas* 19 (5): 1626-1631. DOI: 10.13057/biodiv/d190505.
- Rahman FA, Rohyani IS, Surtiyo, Hadi AP, Lestari DP. 2020. Analisis kualitas perairan terhadap kemelimpahan strata pertumbuhan vegetasi mangrove di Teluk Sereweh, Kabupaten Lombok Timur, Nusa Tenggara Barat, Mataram, Indonesia. *Prosiding Seminar Nasional Forum Ilmiah Pengelolaan Perikanan Berkelanjutan Nusa Tenggara Barat* 2019. Mataram, 3 Desember 2019. [Indonesian]
- Rozainah MZ, Nazri MN, Sofawi AB, Hemati Z, Juliana WA. 2018. Estimation of carbon pool in soil, above and below ground vegetation at different types of mangrove forests in Peninsular Malaysia. *Mar Pollut Bull* 137: 237-245. DOI: 10.1016/j.marpolbul.2018.10.023.
- Saderne V, Gerald NR, Macreadie PI, Maher DT, Middelburg JJ, Serrano O. 2019. Role of carbonate burial in Blue Carbon budgets. *Nat Commun* 10 (1): 1106. DOI: 10.1038/s41467-019-08842-6.
- Sadono R, Soeprijadi D, Susanti A, Matatula J, Pujiono E, Idris F, Wirabuana PYAP. 2020. Local indigenous strategy to rehabilitate and conserve mangrove ecosystem in the southeastern Gulf of Kupang, East Nusa Tenggara, Indonesia. *Biodiversitas* 21 (3): 1250-1257. DOI: 10.13057/biodiv/d210353.
- Scales IR, Friess DA. 2019. Patterns of mangrove forest disturbance and biomass removal due to small-scale harvesting in southwestern Madagascar. *Wetl Ecol Manag* 27 (5): 609-625. DOI: 10.1007/s11273-019-09680-5.
- Shannon MC. 1999. Salinity and horticulture. *Intl J Intl Soc Hort Sci* 78: 1-4.
- Sheil D, Eastaugh CS, Vlam M, Zuidema PA, Groenendijk P, van der Sleen P, Jay A, Vanclay J. 2017. Does biomass growth increase in the largest trees? Flaws, fallacies and alternative analyses. *Funct Ecol* 31 (3): 568-581. DOI: 10.1111/1365-2435.12775.
- Taillardat P, Friess DA, Lupascu M. 2018. Mangrove blue carbon strategies for climate change mitigation are most effective at the national scale. *Biol Lett* 14 (10): 20180251. DOI: 10.1098/rsbl.2018.0251.
- Tam NFY, Wong AHY, Wong MH, Wong YS. 2009. Mass balance of nitrogen in constructed mangrove wetlands receiving ammonium-rich wastewater: Effects of tidal regime and carbon supply. *Ecol Eng* 35 (4): 453-462. DOI: 10.1016/j.ecoleng.2008.05.011.
- Wang J, Bai J, Zhao Q, Lu Q, Xia Z. 2016. Five-year changes in soil organic carbon and total nitrogen in coastal wetlands affected by flow-sediment regulation in a Chinese delta. *Sci Rep* 6: 21137. DOI: 10.1038/srep21137.
- Weiss C, Weiss J, Boy J, Iskandar I, Mikutta R, Guggenberger G. 2016. Soil organic carbon stocks in estuarine and marine mangrove ecosystems are driven by nutrient colimitation of P and N. *Ecol Evol* 6 (14): 5043-5056. DOI: 10.1002/ece3.2258.
- Widyastuti A, Yani E, Nasution EK, Rochmatino. 2018. Diversity of mangrove vegetation and carbon sink estimation of Segara Anakan Mangrove Forest, Cilacap, Central Java, Indonesia. *Biodiversitas* 19 (1): 246-252. DOI: 10.13057/biodiv/d190133.
- Worthington T, Spalding M. 2018. *Mangrove Restoration Potential: A Global Map Highlighting A Critical Opportunity*. Cambridge University, United Kingdom.
- Zhang CG, Leung KK, Wong YS, Tam NFY. 2007. Germination, growth and physiological responses of mangrove (*Bruguiera gymnorhiza*) to lubricating oil pollution. *Environ Exp Bot* 60 (1): 127-133. DOI: 10.1016/j.envexpbot.2006.09.002.

1. Variabilities

ORIGINALITY REPORT

14%

SIMILARITY INDEX

7%

INTERNET SOURCES

7%

PUBLICATIONS

8%

STUDENT PAPERS

PRIMARY SOURCES

1	Submitted to Udayana University Student Paper	2%
2	Submitted to Universitas Diponegoro Student Paper	2%
3	Ratna Herawatiningsih, Eddy Thamrin. "Diversity of meranti (Shorea spp) in secondary forest of tropic area in Mempawah District, West Kalimantan, Indonesia", IOP Conference Series: Earth and Environmental Science, 2021 Publication	1%
4	e-journal.undikma.ac.id Internet Source	1%
5	www.cirebonpower.co.id Internet Source	1%
6	Submitted to SDM Universitas Gadjah Mada Student Paper	1%
7	www.wrc.org.za Internet Source	1%

8	Submitted to Universitas Sam Ratulangi Student Paper	1 %
9	repositorio.udf.edu.br Internet Source	<1 %
10	core.ac.uk Internet Source	<1 %
11	icslca.sil.ui.ac.id Internet Source	<1 %
12	ric.zntu.edu.ua Internet Source	<1 %
13	"Blue Carbon Dynamics of the Indian Ocean", Springer Science and Business Media LLC, 2022 Publication	<1 %
14	www.frontiersin.org Internet Source	<1 %
15	Rebeca Hernández-Gutiérrez, Susana Magallón. "The timing of Malvales evolution: Incorporating its extensive fossil record to inform about lineage diversification", Molecular Phylogenetics and Evolution, 2019 Publication	<1 %
16	dyuthi.cusat.ac.in Internet Source	<1 %
17	repository.unmul.ac.id Internet Source	<1 %

<1 %

18

vbook.pub

Internet Source

<1 %

19

real-j.mtak.hu

Internet Source

<1 %

20

repository.unair.ac.id

Internet Source

<1 %

21

Jesús Jaime Guerra-Santos, Rosa María Cerón-Bretón, Julia Griselda Cerón-Bretón, Diana Lizett Damián-Hernández et al. "Estimation of the carbon pool in soil and above-ground biomass within mangrove forests in Southeast Mexico using allometric equations", *Journal of Forestry Research*, 2014

Publication

<1 %

22

Joao B Xavier. "Social interaction in synthetic and natural microbial communities", *Molecular Systems Biology*, 04/12/2011

Publication

<1 %

23

Submitted to Coventry University

Student Paper

<1 %

24

Ervina Indrayani, John Kalor, Maklon Warpur, Baigo Hamuna. "Using Allometric Equations to Estimate Mangrove Biomass and Carbon Stock in Demta Bay, Papua Province,

<1 %

Indonesia", Journal of Ecological Engineering, 2021

Publication

25

Submitted to University of Melbourne

Student Paper

<1 %

26

N Manglili, B Nurkin, S A Paembonan, S Millang, M Restu, S H Larekeng. "The potential of carbon deposits to residual stand in Tongkonan lembang Buri' garden of Tana Toraja", IOP Conference Series: Earth and Environmental Science, 2019

Publication

<1 %

27

Submitted to Universitas Borneo Tarakan

Student Paper

<1 %

28

Yustiawati, M S Syawal, Rosidah. "Carbon, Nitrogen and C/N ratio of sediment in a floodplain lake: Lake Tempe, South Sulawesi", IOP Conference Series: Earth and Environmental Science, 2022

Publication

<1 %

29

"Mangroves: Ecology, Biodiversity and Management", Springer Science and Business Media LLC, 2021

Publication

<1 %

30

"Proceeding of the 1st International Conference on Tropical Agriculture", Springer Science and Business Media LLC, 2017

Publication

<1 %

Exclude quotes On

Exclude bibliography On

Exclude matches < 10 words