1 ***Please list the reviewers' comments and author's response into a Table.***

2 ***Reviewer A***

Page number	Line number	Reviewer's Comment	Author's Revision
	Abstract		
3	15	What type of manure Is this rice straw	Cattle manure; chopped rice straw
3	22	This is very poorly written. Please rewrite for clarity. Also use correct English .	I have rewritten it . The revision of the writing has been proof read by ASN
	Introduction		
3	33-38	Please rewrite the statement and include the source.	I did rewrite and include source (Suwardji et al., 2012 and Sukartono, et al., 2013)
4	55	What type of manure	Cattle manure
	Material and Methods		
4	86	Is it available P or total P	Available-P (Bray 1)
5	90	Font not italic	I have revised
5	91	Sentence not clear	I revised sentence through proof reading
5	101	Format not italic	2.2. Experimental design and treatments
5	105-112	Not clear. Which height? English problem. Not clear.	40 cm is the high of bed (from surface plot to the farrow). revised the sentence
5	112	state the water content also.	Soil was kept moist at 80% of water field capacity
5	114	Not clear. Please re-write.	I had rewritten sentence
5	126	Is it by the macro or micro Kjeldahl method	Total N was determined using Kjeldahl method (Rayment and Lyons, 2011). Kjeldahl glass used was 250 ml volume
5	124-133	Sentence not clear	I revised sentence
	Results section		
6	150-177	These results are not well reported. Please report your results properly and also eliminate the references since you are not discussing the results.	I made correction (rewrite) according to the reviewer suggestion
6	155	MWD is usually measured in mm. please present your result in mm, so that one can actually know the MWD of water stable aggregates	MWD is the mean weight diameter of aggregate (mm). I used the obtained MWD value (equation 1) to calculate aggregate stability (%) (equation 2).
6	162	What is TDB	TDB: top dry biomass already stated clearly in methods section (Agronomy measurements)
6	150-177	This presentation of the results is very poor. You can improve on it	I did correction for improving the results presentation
	Discussion section		
7	180	What is proposed treatment?	The organic amendment treatments
7	180-190	Are you presenting the results again or you should be discussing your results??	I already revised the sentence
7	191-225	Repetition of results instead of Discussion	I already revised the sentence
	Conclusion		
9	266	Conclusion: mention the soil properties	I already revised conclusion

3

Reviewer B

Page number	Line number	Reviewer's Comment	Author's Revision
	ABSTRACT		
3	11	Change word: A field trial to field exp.	Yes - A field experiment
3	14	Add the volume	Yes I added the amount of amendment material applied for treatments
	INTRODUCTION		
3	36	Can you add any reference about low nutrient and depletion of SOC in dryland?	Sukartono, et al (2013)
3	44	Every single statement here, I think you can add any reference.	This is general condition in the northern Lombok dry land until now. (no reference for this information)
	MATERIAL AND METHODS		
4	85	Please change all comma to dot = 1,14 g cm^{-3} to 1.14 g cm^{-3}. And please change in all text	Yes I already changed
5	91	Preparation of biochar, cattle manure and rice straw. Please add more information about it	Yes I already add more detail information dealing with biochar preparation
5	110-111	Did you any treatment for rice straw before incorporate to the soil wit biochar, like cutting in some centimeter.	Dry rice straw was chopped into size of approximately 3 cm
	RESULTS		
6	152	So the cattle manure and rice straw is fresh?	Dry cattle manure used had C/N of 25. Rice straw was dry. I revised the sentence to make it more clear
	DISCUSSION		
7	180-190	Discussion: Why and How the treatment can increase SOC; Why and How the treatment can improve Total N, available P, Ca, CEC and aggregate stability	I already explain briefly in the discussion section how the treatment can improve the soil characteristics of sandy loams soil in particular N, P, CEC and soil aggregate stability
8	251	How soc can improve the soil aggregate stability?	I already explained in the discussion section
8	226-dst	Evident about effect of biochar and other organic matter on soil quality actually already published in many journal. But more important is how biochar and other organic material can improve the soil quality its more important to explain especially in sandy soil	In the discussion section, I also have provided confirmation from several publications as evidence of the effect of the application of biochar and organic matter on improving soil quality, especially sandy soil
7	208	How you can state this statement without data? Did you measure the soil microorganism? If not how you can know?	Actually, I did not measure the microorganisms. However, theoretically, I would expect that rice straw as mulch mineralizes more slowly than when it is incorporated thoroughly with soil and/or manure.

Author ~~Response~~ ~~re~~vision ~~to~~send to Reviewr ~~Author~~

Influence of biochar amendments on the soil quality indicators of sandy loam soils under cassava - peanut cropping sequence in the semi-arid tropics of Northern Lombok, Indonesia

ABSTRACT

Low nutrient retention and soil organic matter depletion are the major challenges of the cropping system in the sandy loam soils of Northern Lombok, Indonesia. A field trial was conducted to evaluate the influence of biochar-based organic amendments on the soil quality of sandy loam soils under cassava (*Manihot Esculenta, Crants*)-peanut (*Arachis Hypogeae L.*) cropping sequence. The treatments were as follows: biochar and rice straw (B1), biochar, cattle manure and rice straw (B2), biochar and cattle manure (B3), biochar and cattle manure, plus rice straw mulch applied on surface soils (B4), and without organic amendments (B0) as control. Results showed that biochar-based organic amendments significantly improved several soil quality indicators such as SOC, total-N, available P, Ca, cation exchange capacity (CEC) and aggregate stability, but had no significant effect on pH, K and Mg. Improvement in soil quality was strongly indicated by an increase in the growth, and yields of cassava and peanuts. Treatments B1, B2, B3 and B4, generally had a comparable effect on soil parameters and tended to improve the growth and yield of cassava and peanuts. Cassava was responsive to treatments B2 (biochar, cattle manure and rice straw) and B3 (biochar and cattle manure) with its actual yield of 27 ton ha^{-1}, which is a 40% increase compared with that in the control. As a secondary crop growing after cassava, peanuts also exhibit higher yields in all amended plots compared with that in the control. The highest yield was obtained in B2 (1.38 ton ha^{-1}), followed by B4 (1.36 ton ha^{-1}), B1 (1.33 ton ha^{-1}), and B3 (1.25 ton ha^{-1}). In conclusion, the incorporation of biochar, cattle manure and crop residues (rice straw) into soils is a promising option to maintain soil quality and sustainably produce cassava and peanuts in the sandy loam soils of the semi-arid tropics of Lombok, Indonesia.

Keywords: biochar, cattle manure, crop residues, soil quality

1. INTRODUCTION

Indonesia has a great opportunity to increase production of cassava and peanuts by optimizing and developing sustainable agriculture practices in the dryland area. However, sustainable agriculture in the dry land in particular on sandy soils, generally faces large constraints due to low nutrient retention capacity and soil organic matter depletion.

West Nusa Tenggara, located in the eastern part of Indonesia, has potential dry lands of about 1,807,463 ha; of which, 335, 136 ha is relatively suitable for ~~productive~~ agriculture and about 38,000 ha is located in North Lombok (Suwardji, et al., 2012). This area is favorable for food crops such as cassava, peanuts and maize. Soils in this area are dominated by entisols, which are predominately formed from volcanic ash materials derived from the Mount Rinjani eruption. The ~~C~~characteristics of the ~~The~~ soils ~~,~~are as follows: ~~has a~~ light texture with a sand fraction of more than - 50%, poor soil structure, low soil organic-C (SOC) content, ~~-~~infertility and low water retention (Sukartono et al., 2013)

Formatted: English (United States)

Commented [A1]: What type of Manure? Eg poultry manure etc

Commented [A2R1]: Cattle manure

Commented [A3]: What manure

Commented [A4R3]: Cattle manure

Commented [A5]: Is it the rice straw?

Commented [A6R5]: Yes we used the rice straw

Commented [A7]: What type of Manure? Eg poultry manure etc

Commented [A8R7]: Cattle manure

Commented [A9]: What manure

Formatted: Superscript

Commented [A10]: This is very poorly written. Please rewrite for clarity. Also use correct English .

Commented [A11R10]: Yes I did rewrites

Formatted: Indent: Left: 0 cm, Numbered + Level: 1 + Numbering Style: 1, 2, 3, … + Start at: 1 + Alignment: Left + Aligned at: 0,63 cm + Indent at: 1,27 cm

Formatted: Line spacing: Multiple 1,15 li

Field Code Changed

Traditional farmers in the dry land of North Lombok commonly grow cassava as the first crop in early wet season, followed by peanuts as a secondary crop soon after harvesting of cassava. Hence, the common cropping pattern in the area is cassava-peanut-fallow. Peanuts are selected as a secondary crop after cassava due to several considerations: (i) peanut is a legume crop that generates biomass for good quality green manure; (ii) peanut as a part of rotational crops contributes significantly to improve soil fertility, especially nitrogen and SOC (iii) this crop has promising economic value. Soil and cropping management based on organic amendments seems to be an appropriate strategy to achieve sustainable production for both cassava and peanuts.

For sustainable production of these two crops in North Lombok, the limiting factors of soil fertility (i.e. low SOC content and poor nutrient retention and soil structure) must be overcome by implementing conservation-based soil management including addition of organic amendments materials such as biochar and other fresh organic materials (i.e., cattle manure and crop residues). Soil management through the addition of fresh organic matter such as cattle manure has been widely reported to improve soil fertility (Bhatt et al., 2019; Rayne and Aula, 2020) and crop yield in dry land (Sukartono et al. 2011), however, the effect mostly lasts for only one growing season. The use of these organic sources combined with biochar for a cropping rotation of cassava-peanuts has not been carried out.

Biochar is a recalcitrant and stable carbon material in soils. It is a good option as soil amendment for

Previous studies showed that under tropical conditions, the addition of biochar into the soil significantly improved soil chemical properties (Sukartono et al., 2013; Kartika et al., 2018), water retention, and soil aggregates (Zhang et al., 2017; Blanco-Canqui, 2017).Increased SOC content and soil water retention under maize cropping system was also reported in North Lombok by Sukartono et al., (2013). Unfortunately, the incorporation of biochar combined with local fresh organic matter such as cattle manure and rice straw in the root zone of the cassava-peanut cropping sequence in North Lombok has not been explored. Cassava and peanut have a typical root system that requires crumb soil structure and good aggregate, both of which can be induced by supplementing biochar and fresh organic matter. These organic amendments may have a positive impact on the growth and yields of both crops. The present study aimed to evaluate the influence of biochar-based organic amendments (biochar, cattle manure and rice-straw) in improving soil quality of sandy loam soils under cassava-peanut cropping sequence in Northern Lombok.

Commented [A12]: Please rewrite the statement and include the source.

Commented [A13R12]: Source : Suwardji et al (2012) and Sukartono et al (2013)

Commented [A14]: What type of manure. Give the specific one used

Commented [A15R14]: Cattle manure

Commented [A16]: ??

Commented [A17R16]:

2. MATERIAL AND METHODS

A field experiment was carried out at an agricultural dry land in North Lombok, East Indonesia. The experimental site was located at Akar-Akar Village, Subdistrict of Bayan (08° 25'S, 116° 23' E) at 21 m above sea level. The soil developed from volcanic ash and pumice from Mount Rinjani eruption. The topsoil (0-15 cm) has a sandy loam texture (57% sand, 33% silt and 10% clay), 1.14 g cm^{-3} bulk density (BD), pH of 5.98, and low contents of SOC (0.95%), total N (0.12%); available P (14.24 $mgkg^{-1}$), exchangeable K (0.57 cmol kg^{-1}) and cation exchange capacity (CEC) (11.65 cmol kg^{-1}). The trial was conducted under cassava- peanut cropping sequence with casava as the first crop and peanut as the secondary crop.

Formatted: Superscript

2.1. Preparation of biochar.

Biochar was produced using a traditional method by combusting coconut shells in an earth pit with dimensions of 1.0 m depth 0.80 m diameter. Coconut husk was used as the fuel source (Sukartono et al., 2011). Combustion was performed from 195°C to 340°C with an average of 310°C for 5 to 6 hours until the feedstock had completely changed into black charcoal. The chars was then cooled by water spraying and dried for one day. The chars was ground and sieved using a 1.0 mm mesh sieve. The final product of biochar contained 8.5% water, 70.20% C, 0.15% P, 0.76% K, 8.12% ash with pH 8.9 and potential CEC of 12.08 cmol kg^{-1}. Cattle manure had pH 6.8, and contained 11% water, 10.18% C, 0.95% total N, 0, 70% available P, and and 0.65% K.

Commented [A19]: Sentence not clear

Commented [A20R19]: I did rewrite

Commented [A21]: Is it available P or total P

Commented [A22R21]: Available-P (Bray 1)

2.2. Experimental design and treatments.

Commented [A23]: Not Italic

Formatted: Font: Not Italic

Field experiment was set up using a randomized complete block design with five treatments replicated four times. The experiment was carried out in one cycle of the cassava-peanut cropping sequence from February 2015 to April 2016.

The organic amendments were as follows: incorporated biochar and rice straw (B1), incorporated biochar, cattle manure and rice straw (B2); incorporated biochar and cattle manure (B3); incorporated biochar, cattle manure, and rice straw on surface soil (B4); and a control treatment without organic amendments (B0). The size of each plot was 4 m long, 3.5 m wide, and 40 cm high with a space of 0.5 m between plots. Biochar (10 tons ha^{-1}) combined with manure (10 ton ha^{-1}) and rice-straw (3 ton ha^{-1}) was incorporated into each plot at a depth of 10 cm during tillage operation. All treated plots were incubated for 7 days by watering the soil at approximately 80% field capacity.

Commented [A24]: ?? what manure

Commented [A25R24]: Cattle manure

Commented [A26]: Check. repetition

2.3. Agronomic activities for cassava – peanuts.

Seedling from 12-month-old cassava stems (20 cm length and diameter of 2.5 cm) were planted at depth of 5 cm and a spacing of 100 cm x 50 cm at 7 days post treatment (February 2015), and the soil was kept moist at 80% field capacity. Cassava was fertilized by Urea at rates of 300 kg Urea ha^{-1}, SP 36 at 200 kg ha^{-1} and KCl at 150 kg ha^{-1}. Urea at 100 kg ha^{-1} was applied three times at 10, 90 and 150 days after planting (DAP). SP-36 (200 kg ha^{-1}) and KCl (150 kg ha^{-1}) were basally applied at 5 cm from the stems and 10 cm deep in the soil.

Cassava was harvested at 330 DAP by pulling the tubers out from soils. At 7 days post cassava harvesting in January 2016, a local variety of peanut seeds were sown using wooden steaks with a row spacing of 20 cm x 20 cm and a depth of 5 cm.

2.4. Soil sample collection and analysis

Soil samples were collected from each plot at 15 cm top soil before harvest of cassava at 330 DAP. SOC was measured by Walkley and Black method, pH was detected using a pH meter in 1:2.5 soil : water solution, total N was determined by the Kjeldahl method, extractable P was analyzed using Bray-1, and exchangeable cations of K, Ca and Mg and CEC were studied by the NH_4OAc method (Rayment and Lyons, 2011). Soil aggregate stability was measured using a dry and wet sieving method and a modified Yoder sieving machine (Sun and Lu, 2014) with sieves in diameters of 8.00, 4.76, 2.83, 2.0, 1.0, 0.5, and 0.30 mm. The subsamples for aggregate stability analysis were sieved using a 10 mm diameter sieve. Approximately 400 g of the sieved samples were used to determine mean size of the aggregates retained at each sieve. The mean weight diameter (MWD) of soil samples was computed using equation 1 (Sun and Lu, 2014):

$$MWD = \sum_{i}^{j} Xi.Wi \quad (1)$$

Where MWD is the mean weight diameter of aggregate (mm), *Xi* is the mean diameter of ith size fraction, and *Wi* is the proportion of the total sample weight in the corresponding size fraction. The obtained MWD value was used to calculate aggregate stability as follows:

Commented [A29]: Not Italic

Commented [A30]: State the water content also.

Commented [A31R30]:

Formatted: Superscript

Formatted: Font: Not Italic

Commented [A32]: Not clear. Please re-write.

Commented [A33R32]: I had rewrite

Formatted: Font: Not Italic

Commented [A34]: Is it by the macro or micro Kjeldal method

Commented [A35R34]: Total N was determined using Kjeldahl method (Rayment and Lyons, 2011). Kjeldhal glass used was 250 ml volume

Formatted: Subscript

Commented [A36]: Not clear

Commented [A37R36]: I already rewrite

Commented [A38]: Not clear

Commented [A39R38]: I did rewrite

Formatted: Font: Not Italic

Aggregate stability = {1: (MWD_{dry} -MWD_{wet})} x 100% (2)

2.5. Agronomic measurements

The agronomic parameters for cassava were top dry biomass (TDB) and weight of fresh tubers harvested at 330 DAP, and those for peanuts were TDB , weight of dry pods (WDP) and grains (WDG) and N uptake. N-uptake was determined by multiplying the TDB with N concentration in plant tissue at 60 DAP. The effects of treatments on soil and agronomics parameters were analyzed using ANOVA, and significance was tested by Fischer's least significant difference (p=0,05) using Minitab program version 18.

3. RESULTS

3.1. Soil chemical characteristics.

Table 1 shows that the addition of biochar + fresh organic matter based soil amendments had no significant effect on pH, K, and Mg, but affected concentration of SOC, total N, P, Ca and CEC. These parameters were higher in the amended group than those in the control. Meanwhile, total N in B2 plot was higher than that in the control and was similar to those in B1, B3 and B4 plots.

.

.

.

.

Commented [A40]: These results are not well reported. Please report your results properly and also eliminate the references since you are not discussing the results.

Commented [A41R40]: I did rewrite

3.2. Soil aggregate stability.

Soil aggregates stability in unit percent (%) was evaluated using MWD values (Sun & Lu, 2014). As shown in Fig. 1, the soil aggregate stability was 59.24, 59.33, 58.21, and 58.95 (% MWD) for B1, B2, B3 and B4 plots respectively. These values were significantly higher than the 56.59% MWD of no-amendment plot (B0). No significant difference in soil aggregate stability was observed among the plots under the four amendments.

Commented [A42]: MWD is usually measured in mm. please present your result in mm, so that one can actually know the MWD of water stable sggregates

Commented [A43R42]: MWD is the mean weight diameter of aggregate (mm) . The obtained MWD value (equation 1) was used to calculate aggregate stability (%) as stated in the equation 2.

Commented [A44]: Not clear

3.3. Growth and yields of cassava and peanuts.

The biochar-based organic amendments had a significant effect on the growth and yield of cassava as the first crop and peanuts as the secondary crop (Table 2). The TDB of cassava increased significantly by 16% in B1 plot and 20% in B2, B3, and B4 plots relative to that in the control. No significant difference in harvested biomass was observed among the plots under the four amendments. However, tuber yield under all treatments significantly differed from that in the control (18.53 ton

Registrasi pada Jurnal (tanggal 11 Juli 2022)



Revisi Artikel

Mengirimkan hasil Revisi (tanggal 15 September 2022)



ISSN: 1412-3606 (Print) 2356-1424 (Online)
SAINS TANAH - Journal of Soil Science and Agroclimatology
https://jurnal.uns.ac.id/tanah

Submission

Authors	Sukartono Sukartono, Bambang Hari Kusumo, Suwardji Suwardji, Arifin Aria Bakti, Mahrup Mahrup, Lolita Endang Susilowati, Fahrudin Fahrudin
Title	Influence of biochar amendments on the soil quality indicators of sandy loam soils under cassava–peanut cropping sequence in the semi-arid tropics of Northern Lombok, Indonesia
Original file	65452-182051-1-SM.doc 2022-09-19
Supp. files	None Add a Supplementary File
Submitter	Sukartono Sukartono
Date submitted	September 19, 2022 - 10:32 PM
Section	RESEARCH
Editor	Komariah Komariah
Author comments	Herewith a publication manuscript that has passed the ICOSATA review to be published in Sains Tanah

Status

Status	In Editing
Initiated	2022-12-24
Last modified	2022-12-29

Submission Metadata

ISSN
1412-3606 (Print)

Manuscript Template

INDEXING

» My Journals

» My Profile

» Log Out

Information

- For Readers
- For Authors
- For Librarians

Journal Content

Search

Search Scope

All

Search

Browse

- By Issue
- By Author
- By Title
- Other Journals
- Categories

TOOLS

Round 1

Review Version	65452-182054-2-RV.docx	2022-09-20
Initiated	2022-09-20	
Last modified	2022-10-24	
Uploaded file	Reviewer A 65452-184383-1-RV.docx	2022-10-07
	Reviewer A 65452-184383-2-RV.docx	2022-10-07
	Reviewer B 65452-185760-1-RV.docx	2022-10-24

Editor Decision

Decision	Accept Submission 2022-12-24	
Notify Editor	Editor/Author Email Record 2022-12-01	
Editor Version	65452-182103-1-ED.docx	2022-09-20
Author Version	65452-186906-1-ED.docx	2022-11-07 Delete
	65452-186906-2-ED.docx	2022-11-17 Delete
	65452-186906-3-ED.docx	2022-12-05 Delete
	65452-186906-4-ED.docx	2022-12-29 Delete
Upload Author Version	Telusuri... Tidak ada berkas dipilih.	Upload

Editorial Team of Sains Tanah <jurnal@mail.uns.ac.id>
kepada saya

24 Okt 2022, 14.44

Inggris > Indonesia Terjemahkan pesan Nonaktifkan untuk: Inggris

Dear Sukartono:

We have reached a decision regarding your submission to SAINS TANAH - Journal of Soil Science and Agroclimatology, "The influence of biochar-based organic amendments on changes in soil quality of sandy loam soils under cropping sequence of cassava-peanut in the tropical-semi-arid of northern Lombok, Indonesia".

Our decision is: Revisions Required (due date is November 07, 2022)

Please revise your article according to the comments. We kindly ask you to resubmit corrected article under the same identification number. To do so, login into the system, click on this article and fill in "Upload Author Version" input field.

The revised version must include highlighted changes and modifications recommended in the first revision to ensure that all reviewer(s)' comments were considered.

Should you have any questions about the system or other functions please do not hesitate to contact us.

Best regards.

Dr. Komariah
Associate Editor in Chief of SAINS TANAH
Department of Soil Science, Faculty of Agriculture, Sebelas Maret University
(Scopus Author ID: 48661102400)
sainstanah@mail.uns.ac.id

Reviewer A:

50 dari 372

Revision Sukartono Article

Eksternal Kotak Masuk ×

Dr. Ir. Sukartono <kartono1962@unram.ac.id> Rab, 26 Okt 2022, 20.19

kepada Komariah

Dear editor, Thanks very much for sending me the revision article from reviewer. I am going to make great revision as recommended by reviewer and I should resubmit before 7 of Nov. I would like to ask whether you have proof reading services. I am happy to spent compensation fee for the proof reading if possible. I am trying hard to revise very soon.
Best Regards,
Sukartono

J. Soil Sci. & Agric <sainstanah@mail.uns.ac.id> Rab, 26 Okt 2022, 20.35

kepada komariah, saya

Inggris > Indonesia Terjemahkan pesan Nonaktifkan untuk: Inggris ×

You can contact our partner Akses Sains Nasional (E-mail address: asn.office17@gmail.com, Whatsapp/Phone: +62 882-2950-2650) who has partnership with enago in proofread services.

Thank you. Regards.

Dr. Ir. Sukartono <kartono1962@unram.ac.id> Jum, 28 Okt 2022, 07.48

kepada Soil

Ok terima kasih banyak,
Salam

Balas Teruskan

Mengirimkan hasil Revisi (tanggal 26 Oktober 2022)



Gmail

Telusuri dalam email

Aktif

Mail

Chat

Spaces

Meet

Tulis

Kotak Masuk 278

Berbintang

Ditunda

Terkirim

Draf 22

Selengkapnya

Label

[Imap]/Sent

11 dari 439

Inggris > Indonesia Terjemahkan pesan

Nonaktifkan untuk: Inggris

You can contact our partner Akses Sains Nasional (E-mail address: asn.office17@gmail.com, Whatsapp/Phone: +62 882-2950-2650) who has partnership with enago in proofread services.

Thank you. Regards.

Dr. Ir. Sukartono <kartono1962@unram.ac.id>

kepada Soil

28 Okt 2022 07.48

Ok terima kasih banyak,

Salam

Pada tanggal Rab, 26 Okt 2022 pukul 20.35 J. Soil Sci. & Agric <sainstanah@mail.uns.ac.id> menulis:

> You can contact our partner Akses Sains Nasional (E-mail address: asn.office17@gmail.com, Whatsapp/Phone: +62 882-2950-2650) who has partnership with enago in proofread services.
>
> Thank you. Regards.
>
> Pada tanggal Rab, 26 Okt 2022 19:19, Dr. Ir. Sukartono <kartono1962@unram.ac.id> menulis:
>
> > Dear editor, Thanks very much for sending me the revision article from reviewer. I am going to make great revision as recommended by reviewer and I should resubmit before 7 of Nov. I would like to ask whether you have proof reading services. I am happy to spent compensation fee for the proof reading if possible. I am trying hard to revise very soon.
> > Best Regards,
> > Sukartono

Balas Teruskan

[STJSSA] Editor Decision (REVISIONS REQUIRED)

Eksternal Kotak Masuk ×

Editorial Team of Sains Tanah <jurnal@mail.uns.ac.id> 14 Nov 2022 14.10

kepada saya

Inggris › Indonesia Terjemahkan pesan Nonaktifkan untuk: Inggris ×

Dear Sukartono:

The revised version must include highlighted changes and modifications recommended in the first revision to ensure that all reviewer(s)' comments were considered.

Please list the reviewers' comments and your response into a Table.

Should you have any questions about the system or other functions please do not hesitate to contact us.

Best regards.

Dr. Komariah
Associate Editor in Chief of SAINS TANAH
Department of Soil Science, Faculty of Agriculture, Sebelas Maret University
(Scopus Author ID: 48661102400)
sainstanah@mail.uns.ac.id

SAINS TANAH - Journal of Soil Science and Agroclimatology
http://jurnal.uns.ac.id/tanah

Gmail

Telusuri dalam email

Aktif

Mail 99+

Chat

Spaces

Meet

Tulis

Kotak Masuk 278

Berbintang

Ditunda

Terkirim

Draf 22

Selengkapnya

Label

[Imap]/Sent

8 dari 439

Dr. Ir. Sukartono <kartono1962@unram.ac.id>

kepada Komariah

15 Nov 2022 13.52

Dear Editor,
Ok I will sent the revised version which include highlighted changes and modifications recommended by reviewer
Best Regards,

Sukartono

•••

Pada tanggal Sen, 14 Nov 2022 pukul 14.10 Editorial Team of Sains Tanah <jurnal@mail.uns.ac.id> menulis:

> Dear Sukartono:
>
> The revised version must include highlighted changes and modifications recommended in the first revision to ensure that all reviewer(s)' comments were considered.
>
> Please list the reviewers' comments and your response into a Table.
>
> Should you have any questions about the system or other functions please do not hesitate to contact us.
>
> Best regards.
>
> Dr. Komariah
> Associate Editor in Chief of SAINS TANAH
> Department of Soil Science, Faculty of Agriculture, Sebelas Maret University
> (Scopus Author ID: 48661102400)
> sainstanah@mail.uns.ac.id
>
> _____
> SAINS TANAH - Journal of Soil Science and Agroclimatology

Mail
Chat
Spaces
Meet
Tulis
Kotak Masuk 278
Berbintang
Ditunda
Terkirim
Draf 22
Selengkapnya
Label
[Imap]/Sent

SAINS TANAH - Journal of Soil Science and Agroclimatology
http://jurnal.uns.ac.id/tanah

Dr. Ir. Sukartono <kartono1962@unram.ac.id>
kepada Komariah

17 Nov 2022 09.23

Dear Editor Jurnal Sains Tanah

Here, I send the revised article based on reviewer recommendation.
I also have submitted the article to the web of sains tanah journal.
Wassalam

Author
Sukartono

Satu lampiran • Dipindai dengan Gmail

65452-184383-2-...

Balas Teruskan

[STJSSA] Editor Decision (REVISIONS REQUIRED)

Eksternal Kotak Masuk ×

Editorial Team of Sains Tanah <jurnal@mail.uns.ac.id> Kam, 1 Des 2022, 13.45

kepada saya, komariah

Inggris > Indonesia Terjemahkan pesan Nonaktifkan untuk: Inggris ×

Dear Sukartono,

Please list the reviewers' comments and your response into a Table.

Should you have any questions about the system or other functions please do not hesitate to contact us.

Best regards.

Dr. Komariah
Associate Editor in Chief of SAINS TANAH
Department of Soil Science, Faculty of Agriculture, Sebelas Maret University
(Scopus Author ID: 48661102400)
sainstanah@mail.uns.ac.id

SAINS TANAH - Journal of Soil Science and Agroclimatology
http://jurnal.uns.ac.id/tanah

Satu lampiran • Dipindai dengan Gmail



Korespondensi Copyediting (tanggal 29 Desember 2022)



Mengirimkan hasil review Copyediting (tanggal 30 Desember 2022)



[STJSSA] Copyediting Review Acknowledgement

Eksternal Kotak Masuk ×

Editorial Team of Sains Tanah <jurnal@mail.uns.ac.id> Jum, 30 Des 2022, 14.00

kepada saya

Inggris > Indonesia Terjemahkan pesan Nonaktifkan untuk: Inggris ×

Dear Dr. Sukartono:

Thank you for reviewing the copyediting of your manuscript, "Influence of biochar amendments on the soil quality indicators of sandy loam soils under cassava–peanut cropping sequence in the semi-arid tropics of Northern Lombok, Indonesia," for SAINS TANAH - Journal of Soil Science and Agroclimatology. We look forward to publishing this work.

Dr. Komariah
Associate Editor in Chief of SAINS TANAH
Department of Soil Science, Faculty of Agriculture, Sebelas Maret University
(Scopus Author ID: 48661102400)
sainstanah@mail.uns.ac.id

_____
SAINS TANAH - Journal of Soil Science and Agroclimatology
http://jurnal.uns.ac.id/tanah

Dr. Ir. Sukartono <kartono1962@unram.ac.id> Jum, 30 Des 2022, 14.02

kepada Komariah

Matur nuwun bu Dr. Komariah atas perhatian dan kerjasamanya
Salam hormat

Sukartono

•••

[STJSSA] Proofreading Acknowledgement (Author)

Eksternal Kotak Masuk ×

Editorial Team of Sains Tanah <jurnal@mail.uns.ac.id> Jum, 30 Des 2022, 14.34

kepada saya

Inggris > Indonesia Terjemahkan pesan Nonaktifkan untuk: Inggris ×

Dear Dr. Sukartono:

Thank you for proofreading the galleys for your manuscript, "Influence of biochar amendments on the soil quality indicators of sandy loam soils under cassava–peanut cropping sequence in the semi-arid tropics of Northern Lombok, Indonesia," in SAINS TANAH - Journal of Soil Science and Agroclimatology. We are looking forward to publishing your work shortly.

If you subscribe to our notification service, you will receive an email of the Table of Contents as soon as it is published. If you have any questions, please contact me.

Dr. Komariah
Associate Editor in Chief of SAINS TANAH
Department of Soil Science, Faculty of Agriculture, Sebelas Maret University
(Scopus Author ID: 48661102400)
sainstanah@mail.uns.ac.id

SAINS TANAH - Journal of Soil Science and Agroclimatology
http://jurnal.uns.ac.id/tanah

SAINS TANAH – Journal of Soil Science and Agroclimatology

Journal homepage: http://jurnal.uns.ac.id/tanah

Influence of biochar amendments on the soil quality indicators of sandy loam soils under cassava–peanut cropping sequence in the semi-arid tropics of Northern Lombok, Indonesia

Sukartono*, Bambang Hari Kusumo, Suwardji, Arifin Aria Bakti, Mahrup, Lolita Endang Susilowati, Fahrudin

Department of Soil Science, Faculty of Agriculture, University of Mataram, Indonesia

ARTICLE INFO

Keywords:
Biochar
Cattle manure
Crop residues
Soil quality

Article history
Submitted: 2022-09-19
Accepted: 2022-12-24
Available online: 2022-12-30
Published regularly: Dec 2022

* *Corresponding Author*
Email address:
kartono1962@unram.ac.id

ABSTRACT

Low nutrient retention and soil organic matter depletion are the major challenges of the cropping system in the sandy loam soils of Northern Lombok, Indonesia. A field experiment was conducted to evaluate the influence of biochar-based organic amendments on the soil quality of sandy loam soils under cassava (*Manihot esculenta*, Crantz)–peanut (*Arachis hypogaea* L.) cropping sequence. The treatments were as follows: biochar (10 ton ha^{-1}) and rice straw (3 ton ha^{-1}) (B1); biochar (10 ton ha^{-1}), cattle manure (10 ton ha^{-1}), and rice straw (3 ton ha^{-1}) (B2); biochar (10 ton ha^{-1}) and cattle manure (10 ton ha^{-1}) (B3); biochar (10 ton ha^{-1}) and cattle manure (10 ton ha^{-1}) plus rice straw mulch (3 ton ha^{-1}) applied on surface soils (B4), and without organic amendments (B0) as control. Results showed that the biochar-based organic amendments significantly improved several soil quality indicators such as SOC, total N, available P, Ca, cation-exchange capacity (CEC), and aggregate stability but had no significant effect on pH, K, and Mg. Improvement in soil quality was strongly indicated by an increase in the growth and yield of cassava and peanuts. Treatments B1, B2, B3, and B4 generally had a comparable effect on soil parameters and tended to improve the growth and yield of cassava and peanuts. Cassava was responsive to treatments B2 (biochar, cattle manure, and rice straw) and B3 (biochar and cattle manure) with its actual yield of 27 tons ha^{-1}, which is a 40% increase compared with that in the control. As a secondary crop growing after cassava, peanuts also exhibited higher yields in all amended plots compared with that in the control. The highest yield was obtained in B2 (1.38 ton ha^{-1}), followed by B4 (1.36 ton ha^{-1}), B1 (1.33 ton ha^{-1}), and B3 (1.25 ton ha^{-1}). In conclusion, the incorporation of biochar, cattle manure, and crop residues (rice straw) into soils is a promising option to maintain soil quality and sustainably produce cassava and peanuts in the sandy loam soils of the semi-arid tropics of Lombok, Indonesia.

How to Cite: Sukartono., Kusumo, B H., Suwardji., Bakti, A A., Mahrup., Susilowati, L E., Fahrudin. (2022). Influence of biochar amendments on the soil quality indicators of sandy loam soils under cassava–peanut cropping sequence in the semi-arid tropics of Northern Lombok, Indonesia. Sains Tanah Journal of Soil Science and Agroclimatology, 19(2): 205-210. https://dx.doi.org/10.20961/stjssa.v19i2.65452

1. INTRODUCTION

Indonesia has a great opportunity to increase its production of cassava and peanuts by optimizing and developing sustainable agriculture practices in the dryland area. However, sustainable agriculture in dry land, particularly on sandy soils, generally faces large constraints due to low nutrient retention capacity and soil organic matter depletion (Sukartono, 2011). West Nusa Tenggara, located in the eastern part of Indonesia, has potential dry lands of about 1,807,463 ha; of which, 335, 136 ha is relatively suitable for agriculture and about 38,000 ha is located in North Lombok (Sukartono, 2011). This area is favorable for food crops such as cassava, peanuts, and maize. Soils in this area are dominated by entisols, which are predominately formed from volcanic ash materials derived from the Mount Rinjani eruption. The characteristics of the soils are as follows: light texture with a sand fraction of more than 50%, poor soil structure, low soil organic C (SOC) content, infertility, and low water retention (Sukartono et al., 2013).

Traditional farmers in the dry land of North Lombok commonly grow cassava as the first crop in early wet season,

Scopus Preview

Source details

Feedback > Compare sources >

Sains Tanah

Open Access

Scopus coverage years: from 2019 to 2022

Publisher: Sebelas Maret University - Department of Soil Science, Faculty of Agriculture

ISSN: 1412-3606 E-ISSN: 2356-1424

Subject area: Agricultural and Biological Sciences: Agronomy and Crop Science | Agricultural and Biological Sciences: Soil Science | Environmental Science: Pollution | Earth and Planetary Sciences: Atmospheric Science

Source type: Journal

View all documents > | Set document alert | Save to source list | Source Homepage

CiteScore 2021
0.9

SJR 2021
0.282

SNIP 2021
2.838

CiteScore | CiteScore rank & trend | Scopus content coverage

Improved CiteScore methodology

CiteScore 2021 counts the citations received in 2018-2021 to articles, reviews, conference papers, book chapters and data papers published in 2018-2021, and divides this by the number of publications published in 2018-2021. Learn more >

CiteScore 2021

0.9 = 46 Citations 2018 - 2021 / 49 Documents 2018 - 2021

Calculated on 05 May, 2022

CiteScoreTracker 2022

1.4 = 134 Citations to date / 95 Documents to date

Last updated on 03 March, 2023 • Updated monthly

CiteScore rank 2021

Category	Rank	Percentile
Agricultural and Biological Sciences — Agronomy and Crop Science	#272/370	26th
Agricultural and Biological Sciences — Soil Science	#110/145	24th

View CiteScore methodology > | CiteScore FAQ > | Add CiteScore to your site