

B11

by Adhitya Wisnu

Submission date: 16-May-2022 04:23PM (UTC-0500)

Submission ID: 1837870066

File name: Lampiran_B11.pdf (2.02M)

Word count: 2543

Character count: 10855

Journal of Fundamental Mathematics and Applications (JFMA)

Editor in Chief : Sunarsih

Publisher : Mathematics Department, Faculty of Science and Mathematics, Universitas Diponegoro

Subject : Engineering and Technology, Computers, Informations and Mathematics

[View Journal](#)



About this Journal

Journal of Fundamental Mathematics and Applications (JFMA) is an Indonesian journal published by the Department of Mathematics, Diponegoro University, Semarang, Indonesia. JFMA has been published regularly in 2 scheduled times (June and November) every year. JFMA is established to highlight the latest update of mathematical researches in both theoretical and applied works. The scope in JFMA is pure mathematics and applied mathematics. All accepted papers will be published both in print and online versions. The online version can be accessed via the DOI link of each article. The print version can be ordered to the journal administrator.

p-ISSN: 2621-6019

e-ISSN: 2621-6035



[Make Submissions](#)

[Focus and Scope](#)

[Peer Review Process / Policy](#)

[Current Issue](#)

[Editorial Team](#)

Journal Performance

Accredited



EJOURNAL UNDIP

UNDIP E-Journal Systems (UEJS) Portal is an e-journal management and publishing system published by Diponegoro University.

LINKS

[Home](#)
[About](#)
[Undip](#)
[LPPM Undip](#)

CONTACT

🏠 Department of Mathematics, Faculty of Science and Mathematics, Diponegoro University
Mailing address: Jl. Prof Soedarto, SH, Tembalang, Semarang, Indonesia 50275
☎ (+6224) 70789493
✉ s.sutrisno@live.undip.ac.id



An Indonesian Journal Published by
Department of Mathematics Diponegoro University



HOME ABOUT LOGIN REGISTER SEARCH CURRENT ARCHIVES
ANNOUNCEMENTS STATISTICS DOWNLOAD

Home > Archives > Vol 3, No 1 (2020)

Vol 3, No 1 (2020)

Table of Contents

FUNDAMENTAL MATHEMATICS AND APPLICATIONS

KARAKTERISASI MODUL CYCLICALLY PURE (CP)-INJEKTIF Nia Yulianti, Hanni Garminia Y	PDF 1-14
COPRIME GRAPH OF INTEGERS MODULO n GROUP AND ITS SUBGROUPS Rina Juliana, Masriani Masriani, I Gede Adhitya Wisnu Wardhana, Ni Wayan Switrayni, Irwansyah Irwansyah	PDF 15-18
FUNGSI WRIGHT SEBAGAI SOLUSI ANALITIK PERSAMAAN DIFUSI-GELOMBANG FRAKSIONAL PADA MEDIA VISKOELASTIS Ray Novita Yasa, Agus Yodi Gunawan	PDF 19-33
SOME PROPERTIES OF COPRIME GRAPH OF DIHEDRAL GROUP D_{2n} WHEN n IS A PRIME POWER Abdul Gazir S, I Gede Adhitya Wisnu Wardhana, Ni Wayan Switrayni, Quratul Aini	PDF 34-38
EASY ENSEMBLE WITH RANDOM FOREST TO HANDLE IMBALANCED DATA IN CLASSIFICATION Sarini Abdullah, GV Prasetyo	PDF 39-46
RUANG BERNORMA LENGKAP ATAS OPERATOR LINEAR TERBATAS PADA RUANG FUNGSI TERINTEGRAL DUNFORD Solikhin Solikhin, YD Sumanto, Abdul Aziz, Susilo Hariyanto, R. Heri Soelistyo Utomo	PDF 47-55
ANALISIS TARIF PREMI DAN ASUMSI LOADING PADA PRODUK ASURANSI DWIGUNA BEASISWA Ira Rosianal Hikmah, Yulial Hikmah	PDF 56-64
RELASI GREEN PADA GAMMA-SEMIGRUP YANG DIBANGKITKAN DARI SUATU SEMIGRUP YD Sumanto, Abdul Aziz, Solikhin Solikhin, R. Heri Soelistyo Utomo	PDF 65-71
NUMERICAL SIMULATION OF STOCHASTIC DIFFERENTIAL EQUATIONS USING IMPLICIT MILSTEIN METHOD Ratna Herdiana	PDF 72-83

Publisher:

Department of Mathematics, Faculty of Science and Mathematics, Diponegoro University
Mailing address: Jl. Prof Soedarto, SH, Tembalang, Semarang, Indonesia 50275
Telp./Fax : (+6224) 70789493 / (+62224) 76480922
Website : www.math.fsm.undip.ac.id
E-mail : admin.math@live.undip.ac.id

Indexed in:



USER

Username
Password
 Remember me

JOURNAL CONTENT

Search

Search Scope
All

Browse

- By Issue
- By Author
- By Title
- Other Journals

CURRENT ISSUE



ABOUT THE JOURNAL

- Editorial Team
- Focus and Scope
- Section Policies
- Open Access Policy
- Publication Ethics
- Copyright Notice
- Copyright Transfer Form
- Indexing and Abstracting
- Article Proc. Charge (APC)
- Manuscript Preparation
- Submission Guidance
- Reviewer Guideline
- Order JFMA hardcopy
- DOWNLOAD

RECOMMENDED TOOLS:



VISITOR COUNTER:





An Indonesian Journal Published by
Department of Mathematics Diponegoro University



HOME ABOUT LOGIN REGISTER SEARCH CURRENT ARCHIVES
ANNOUNCEMENTS STATISTICS DOWNLOAD

Home > About the Journal > Editorial Team

Editorial Team

Chief Editor

Sunarsih Sunarsih, Dept. of Mathematics, Diponegoro University, Indonesia

Editorial Board

- Ahmad Faisol, Dept. of Mathematics, Universitas Lampung, Indonesia
- Arief Fatchul Huda, Dept. of Mathematics, UIN Sunan Gunung Djati, Indonesia
- Bayu Surarso, Dept. of Mathematics, Diponegoro University, Indonesia
- Budi Waluya, Dept. of Mathematics, Universitas Negeri Semarang, Indonesia
- Esther Sorta Mauli Nababan, Dept. of Mathematics, Universitas Sumatera Utara, Indonesia
- Fadilah Ilahi, Dept. of Mathematics, UIN Sunan Gunung Djati, Indonesia
- Farikhin Farikhin, Dept. of Mathematics, Diponegoro University, Indonesia
- Fitriani Fitriani, Dept. of Mathematics, Universitas Lampung, Indonesia
- Herry Pribawanto Suryawan, Dept. of Mathematics, Universitas Sanata Dharma, Indonesia
- Idha Sihwaningrum, Dept. of Mathematics, Universitas Jenderal Soedirman, Indonesia
- Imam Marzuki Shofi, Departemen Teknik Informatika, UIN Jakarta, Indonesia
- Kris Suryowati, IST AKPRIND YOGYAKARTA, Indonesia
- Muhammad Muhajir, Department of Statistics, Universitas Islam Indonesia, Indonesia
- Nanang Susyanto, Dept. of Mathematics, Universitas Gadjah Mada, Indonesia
- Nikken Prima Puspita, Dept. of Mathematics, Diponegoro University, Indonesia
- Puguh Wahyu Prasetyo, Dept. of Mathematics, Universitas Ahmad Dahlan, Indonesia
- Putranto Hadi Utomo, Departemen Matematika, Universitas Sebelas Maret, Indonesia
- Roberd Saragih, Dept. of Mathematics, Institut Teknologi Bandung, Indonesia
- Susilo Hariyanto, Dept. of Mathematics, Diponegoro University, Indonesia
- Sutrisno Sutrisno, Dept. of Mathematics, Universitas Diponegoro, Indonesia
- Widowati Widowati, Dept. of Mathematics, Diponegoro University, Indonesia

Publisher:

Department of Mathematics, Faculty of Science and Mathematics, Diponegoro University
 Mailing address: Jl. Prof Soedarto, SH, Tembalang, Semarang, Indonesia 50275
 Telp./Fax : (+6224) 70789493 / (+62224) 76480922
 Website : www.math.fsm.undip.ac.id
 E-mail : admin.math@live.undip.ac.id

Indexed in:



USER

Username
 Password
 Remember me

JOURNAL CONTENT

Search
 Search Scope
 All

Browse

- By Issue
- By Author
- By Title
- Other Journals

ABOUT THE JOURNAL

- Editorial Team
- Focus and Scope
- Section Policies
- Open Access Policy
- Publication Ethics
- Copyright Notice
- Copyright Transfer Form
- Indexing and Abstracting
- Article Proc. Charge (APC)
- Manuscript Preparation
- Submission Guidance
- Reviewer Guideline
- Order JFMA hardcopy
- DOWNLOAD

RECOMMENDED TOOLS:



VISITOR COUNTER:



00000047733
View JFMA's Stats

1
**Some Properties of Coprime Graph of Dihedral Group D_{2n}
 When n is a prime power**

Abdul Gazir S¹, I Gede Adhitya Wisnu Wardhana^{2*}, Ni Wayan Switrayni³,
 Qurratul Aini⁴

^{1,2,3,4} *Departement of Mathematics, Universitas Mataram*

Email: ¹abdgazirsyazir@gmail.com, ^{2*}adhitya.wardhana@unram.ac.id, ³niwayan.switrayni@unram.ac.id,
⁴qurratul.aini@unram.ac.id

*Corresponding author

Abstract. Research on algebra structures represented graph theory led the way to a new topic of research in recent years. In this paper, the algebraic structure that will be represented in the coprime graph is the dihedral group and its subgroup. The coprime graph of a group G , denoted by Γ_G is a graph whose vertices are elements of G and two distinct vertices a and b are adjacent if only if $(|a|, |b|) = 1$. Some properties of coprime graphs from a dihedral group D_{2n} are obtained. Some of the results is if n is prime then $\Gamma_{D_{2n}}$ is complete bipartite graph and if n is composite then $\Gamma_{D_{2n}}$ is multipartite graph.

Keywords: complete bipartite graph, coprime graph, dihedral group, multipartite graph.

I. INTRODUCTION

Study of algebraic structures represented in a graphs raises many recent and in interesting results. This area is relatively new, and over the years different types of graphs of a group were defined. For example, prime graph, the non-commuting graph, and Cayley graphs, which have a long history. In 2014, Ma et al define a coprime graph of a group [1]. The coprime graph of a group G , denoted by Γ_G is a graph whose vertices are elements of G and two distinct vertices a and b are adjacent if only if $(|a|, |b|) = 1$ [2]. Dorbidi in 2016 classify all the groups which Γ_G is a complete t -partite graph or a planar graph, he also studied the automorphism group of Γ_G [3]. In this paper, we will study some properties of coprime graph of a dihedral group.

II. RESULT

Some definitions of a group and a graph that used in this paper are given as follows.

Definition 1 ([4]) *A dihedral group G with identity e , is a group generated by two elements a, b with properties*

$$G = \{a, b | a^n = e, b^2 = e, bab^{-1} = a^{-1}\}$$

It is easy to see that the order of dihedral group is $2n$ and we denote dihedral with order $2n$ by D_{2n} . We can also rewrite $D_{2n} = \{e, a, a^2, a^3, \dots, a^{n-1}, b, ab, a^2b, a^3b, \dots, a^{n-1}b\}$. For example, for $n = 4$ we have $D_8 = \{e, a, a^2, a^3, b, ab, a^2b, a^3b\}$.

Definition 2 *If G is a group with identity e and $x \in G$, the order of x is the least natural number k such that $x^k = e$ and we write $|x| = k$.*

Definition 3 ([1]) The coprime graph of a group G , denoted by Γ_G is a graph whose vertices are elements of G and two distinct vertices u and v are adjacent if and only if $(|a|, |b|) = 1$.

2.1. Coprime Graph of D_{2n}

Given dihedral group D_{2n} , we will give some properties of $\Gamma_{D_{2n}}$ and Γ_S for all non trivial subgroup S of D_{2n} . The result is quite interesting, the coprime graph of D_{2n} always a tripartite graph except for n is a power of two.

First result, if n is prime then the coprime graph of D_{2n} is given by the following Theorem.

Theorem 1 Let n is an odd prime number, then the coprime graph $\Gamma_{D_{2n}}$ is a complete tripartite graph.

Proof. Let n is an odd prime number, Define three partitions by $V_1 = \{e\}$, $V_2 = \{a, a^2, \dots, a^{p-1}\}$ and $V_3 = \{b, ab, a^2b, \dots, a^{p-1}b\}$. Clearly $|e| = 1$, as n is prime number. Since the order of a is n and the order of b is 2 then we have $|a| = |a^2| = \dots = |a^{n-1}| = n$, and $|b| = |ab| = |a^2b| = \dots = |a^{n-1}b| = 2$. Hence for each $x, y \in V_i$ then $(|x|, |y|) = p \neq 1$ or $(|x|, |y|) = 2 \neq 1$, $\in \{2, 3\}$. Then for any $u \in V_i$ dan $v \in V_j$ where $i \neq j$ we have $(|u|, |v|) = 1$, thus u and v are adjacent so the coprime graph of the dihedral group is complete tripartite graph. \square

Second case is, if n is the power of 2, then the coprime graph of D_{2n} is given by the following Theorem.

Theorem 2 Let $n = 2^k$, for some $k \in \mathbb{N}$ then the coprime graph $\Gamma_{D_{2n}}$ is a complete bipartite graph.

Proof. Let $D_{2n} = \{e, a, a^2, a^3, \dots, a^{2^k-1}, b, ab, a^2b, a^3b, \dots, a^{2^k-1}b\}$ is a dihedral group. Define two partitions $V_1 = \{e\}$, and $V_2 = \{a, a^2, \dots, a^{2^k-1}, b, ab, a^2b, a^3b, \dots, a^{2^k-1}b\}$. Clearly $|e| = 1$, as $n = 2^k$ then we have $|a| = |a^2| = \dots = |a^{2^k-1}| = 2^s$ and $|b| = |ab| = |a^2b| = \dots = |a^{2^k-1}b| = 2$. Hence for each $x, y \in V_2$ we have $(|x|, |y|) = 2^s \neq 1$ for some $s \in \mathbb{N}$, thus x and y are not neighbors. Since $|e| = 1$ and for any $v \in V_2$ we have $(|e|, |v|) = 1$. We have e is adjacent to $v \in V_2$, then the Coprime graph of the dihedral group is a complete bipartite graph. \square

Third case is, if n is prime power, then the coprime graph of D_{2n} is given by the following Theorem.

Theorem 3 Let $n = p^k$, $p \neq 2$ and p is prime number, for some $k \in \mathbb{N}$ then the coprime graph $\Gamma_{D_{2n}}$ is a complete tripartite graph.

Proof. Let $D_{2n} = \{e, a, a^2, \dots, a^{p^k-1}, b, ab, a^2b, \dots, a^{p^k-1}b\}$ is a dihedral group. Define three partitions $V_1 = \{e\}$, $V_2 = \{a, a^2, \dots, a^{p^k-1}\}$ and $V_3 = \{b, ab, a^2b, \dots, a^{p^k-1}b\}$. Clearly $|e| = 1$, since $n = p^k$, $p \neq 2$ then $|a|, |a^2|, \dots, |a^{p^k-1}|$ is divided by p and $|b| = |ab| = |a^2b| = \dots = |a^{p^k-1}b| = 2$. Hence for each $x, y \in V_2$ we have $(|x|, |y|)$ is power of p or $(|x|, |y|) = 2 \neq 1$, then for any $u \in V_i$ and $v \in V_j$ where $i \neq j$ then $(|u|, |v|) = 1$. Hence u and v are adjacent, and the coprime graph of the dihedral group is a complete tripartite graph. \square

2.2. Coprime Graph of Nontrivial Subgroup of D_{2n}

In this section we give the coprime graph of a subgroup of a dihedral group. The result is quite interesting, the coprime graph of nontrivial subgroups of D_{2n} is always a complete bipartite graph except for n is a power of odd prime. If n is a power of odd prime then there is a chance that the coprime graph of subgroup D_{2n} is a complete tripartite graph.

First, if n is prime then coprime graph of any subgroup of D_{2n} is given by the following Theorem.

Theorem 4 Let S is a nontrivial subgroup of D_{2n} . If n is prime number, then the coprime graph Γ_S is a complete bipartite graph.

Proof. From [5] we know all the subgroup of D_{2n} can be categorized into several subgroups, so we will divided into 2 cases

Case 1 (Rotation subgroup)

The subgroup contain identities and all rotation element, write $S_1 = \{e, a, a^2, \dots, a^{n-1}\}$. Define two partitions $V_1 = \{e\}$, $V_2 = \{e, a, a^2, \dots, a^{n-1}\}$. Clearly $|e| = 1$, as n is prime number and $|a| = |a^2| = \dots = |a^{n-1}| = n$. Hence for each $x, y \in V_2$ then $(|x|, |y|) = n \neq 1$, then x and y are not neighbors. Since $|e| = 1$, then for any $v \in V_2$ we have $(|e|, |v|) = 1$. Then the coprime graph of subgroup S_1 is complete bipartite graph.

Case 2 (Reflection subgroup)

The subgroups contains identity and one reflection element, they are $S_0 = \{e, b\}$, $S_1 = \{e, ab\}$, \dots , $S_{n-1} = \{e, a^{n-1}b\}$. Let $i \in \{0, \dots, n-1\}$. Define two partitions $V_1 = \{e\}$, and $V_2 = \{a^i b\}$. Since $|e| = 1$ and $|a^i b| = 2$ then for any $v \in V_2$ we have $(|e|, |v|) = 1$. So e is adjacent to any $v \in V_2$. Then the coprime graph of the subgroups S_i is complete bipartite graph. \square

Second, if n is the power of 2, then coprime graph of any subgroup of D_{2n} is given by the following Theorem.

Theorem 5 Let S is a nontrivial subgroup of D_{2n} . If $n = 2^k$, for some $k \in \mathbb{N}$ then the coprime graph of Γ_S is a complete bipartite graph.

Proof. From [5] we know all the subgroup of D_{2n} can be categorized into several subgroups, so we will divided into 3 cases

Case 1 (Rotation subgroup)

The subgroup contain identities and some rotation element, they are $S_0 = \{e, a, a^2, \dots, a^{2^k-1}\}$ the subgroup with all rotation element and $S_i = \{e, a^{2^i}, a^2 \cdot 2^{2^i}, \dots, a^{n-2^i}\}$ for $i \in \{2, \dots, k-1\}$. Define two partitions $V_1 = \{e\}$ and $V_2 = S_i - \{e\}$. Clearly $|e| = 1$, and since $n = 2^k$ then $|a|, |a^2|, \dots, |a^{2^k-1}|$ is divided by 2. Hence for each $x, y \in V_2$ then $(|x|, |y|)$ is power of 2, then x and y are not neighbors. Clearly for any $v \in V_2$ we have $(|e|, |v|) = 1$, so e is adjacent to any $v \in V_2$. Hence the coprime graph of the subgroup S_i is complete bipartite graph.

Case 2 (Reflection subgroup)

The subgroups contains identity and one reflection element, they are $S_1 = \{e, b\}$, $S_2 = \{e, ab\}$, \dots , $S_n = \{e, a^{2^{k-1}}b\}$. Define two partitions $V_1 = \{e\}$, and $V_2 = \{a^i b\}$, for some $3 \leq i \leq n-1$. Since $|e| = 1$ and $|a^i b| = 2$ then for any $v \in V_2$ we have $(|e|, |v|) = 1$. Hence the coprime graph of the subgroups S_i is complete bipartite graph.

Case 3 (Mixed subgroup)

The subgroups containing identity with some reflection element and some rotation element, they are such as $S_i = \{e, a^{2^i}, a^{2^{2^i}}, \dots, a^{n-2^i}b, a^{j+2^i}b, \dots, a^{n-2^i+j}b\}$. Define two partitions $V_1 = \{e\}$, and $V_2 = S_i - \{e\}$. Clearly $|e| = 1$, $|a^{2^i}| = \dots = |a^{n-2^i}| = 2^s$, for some $s \in \mathbb{N}$ and $s \leq k$, and $|b| = |a^{2^i}b| = \dots = |a^{n-2^i}b| = 2$. Hence for each $x, y \in V_2$ we have $(|x|, |y|) = 2 \neq 1$, then x and y are not neighbors. Since $|e| = 1$, for any $v \in V_2$ we have $(|e|, |v|) = 1$. Then e is adjacent to any $v \in V_2$. Hence the coprime graph of the subgroup S_i is a complete bipartite graph. \square

Third, if n is a prime power, then coprime graph of any subgroup of D_{2n} is given by the following Theorem.

Theorem 6 Let S is a nontrivial subgroup of D_{2n} . If $n = p^k$, p is prime number, for some $k \in \mathbb{N}$ then the coprime graph Γ_S is a complete bipartite graph or a complete tripartite graph.

Proof. From [5] we know all the subgroup of D_{2n} can be categorized into several subgroups, so we will divided into 3 cases

Case 1 (Rotation subgroup)

The subgroup contain identities and some rotation element, they are $S_0 = \{e, a, \dots, a^{p^k-1}\}$ the subgroup with all rotation element and $S_i = \{e, a^{p^i}, a^{2 \cdot p^i}, \dots, a^{n-p^i}\}$ for $i \in \{2, \dots, k-1\}$ Define two partitions $V_1 = \{e\}$ and $V_2 = S_i - \{e\}$. Clearly $|e| = 1$, and since $n = p^k$ then $|a|, |a^2|, \dots, |a^{2^{k-1}}|$ is divided by p . Hence for each $x, y \in V_2$ then $(|x|, |y|)$ is the power of p , then x and y are not neighbors. Clearly for any $v \in V_2$ we have $(|e|, |v|) = 1$, so e is adjacent to any $v \in V_2$. Hence the coprime graph of the subgroup S_i is complete bipartite graph.

Case 2 (Reflection subgroup)

The subgroups contains identity and one reflection element, they are $S_1 = \{e, b\}$, $S_2 = \{e, ab\}$, \dots , $S_n = \{e, a^{p^{k-1}}b\}$. Define two partitions $V_1 = \{e\}$, and $V_2 = \{a^i b\}$, for some $3 \leq i \leq n-1$. Since $|e| = 1$ and $|a^i b| = 2$ then for any $v \in V_2$ we have $(|e|, |v|) = 1$. Hence the coprime graph of the subgroups S_i is complete bipartite graph.

Case 3 (Mixed subgroup)

The subgroups containing identity with some reflection element and some rotation element, they are such as $S_i = \{e, a^{p^i}, a^{2 \cdot p^i}, \dots, a^{n-p^i}b, a^{j+p^i}b, \dots, a^{n-p^i+j}b\}$. Define three partitions $V_1 = \{e\}$, $V_2 = \{a^{2^i}, a^{2^{2^i}}, \dots, a^{n-2^i}\}$, and $V_3 = \{b, a^{p^i}b, \dots, a^{n-p^i}b\}$. Clearly $|e| = 1$, since $n = p^k$ then $|a^{p^i}| = \dots = |a^{n-p^i}| = p^s$, for some $s \in \mathbb{N}$ and $s \leq k$. And $|b| = |a^{p^i}b| = \dots = |a^{n-p^i}b| = 2$. Hence for each $x, y \in V_i$ we have $(|x|, |y|) = p^s \neq 1$ or $(|x|, |y|) = 2 \neq 1$ for $i \in \{2, 3\}$. For any $u \in V_i$ and $v \in V_j$ where $i \neq j$ then $(|u|, |v|) = 1$. So, u and v are adjacent. Hence the coprime graph of subgroup S_i is a complete tripartite graph. \square

The last result valid for every subgroup of any dihedral group, even for any group.

Theorem 7 If the coprime graph of the dihedral group is a complete bipartite graph, then the coprime graph of the nontrivial subgroups is a complete bipartite graph.

Proof. Suppose the coprime graph of the dihedral group is in the form of a complete bipartite graph then $\exists V_1, V_2 \subset D_{2n}$ so that for each $v_1 \in V_1$ and $v_2 \in V_2, (|v_1|, |v_2|) = 1$ and for every $a, b \in V_i, i = 1, 2, (|a|, |b|) \neq 1$. Let $H < D_{2n}$, choose $W_1 = V_1 \cap H$ and $W_2 = V_2 \cap H$, hence if $x \in W_1$ then $x \in V_1$ and $y \in W_2$ thus $y \in V_2$ as a result $(|x|, |y|) = 1$. So, the coprime graph of the subgroups is a complete bipartite graph. \square

III. CONCLUSIONS AND FUTURE RESEARCH DIRECTION

The obtained result show that the coprime graph of a dihedral group or its subgroup will be tripartite graph or bipartite graph, depending on n . For the next research, it is interesting to see if n is multiplication more than two different prime numbers.

ACKNOWLEDGEMENT

This paper is dedicated to Professor Ahmad Erfanian who inspired this research.

REFERENCES

- [1] X.L. Ma, H.Q. Wei, and L.Y. Yang, "The coprime graph of a group," *International Journal of Group Theory*, vol. 3, no. 3, pp. 13–23, 2014.
- [2] R. Munir, *Matematika Diskrit*. Bandung: Penerbit Informatika, 2010.
- [3] H.M. Dorbidi, "A note on the coprime graph of a group," *International Journal of Group Theory*, vol. 5, no. 4, pp. 17–22, 2016.
- [4] J. B. Fraleigh, *A First Course in Abstract Algebra*. Harlow: Pearson New International Edition, 2014.
- [5] A. Gazir S and I.G.A.W. Wardhana, "Subgrup Non Trivial Dari Grup Dihedral," *Eigen Mathematics Journal*, vol. 2, no. 2, pp. 73–76, 2019.

ORIGINALITY REPORT

24%

SIMILARITY INDEX

21%

INTERNET SOURCES

20%

PUBLICATIONS

10%

STUDENT PAPERS

PRIMARY SOURCES

1	ejournal2.undip.ac.id Internet Source	9%
2	Submitted to Universitas Diponegoro Student Paper	7%
3	A G Syarifudin, I G A W Wardhana, N W Switrayni, Q Aini. "The Clique Numbers and Chromatic Numbers of The Coprime Graph of a Dihedral Group", IOP Conference Series: Materials Science and Engineering, 2021 Publication	6%
4	A G Syarifudin, Nurhabibah, D P Malik, I G A W Wardhana. " Some characterizations of coprime graph of dihedral group D ", Journal of Physics: Conference Series, 2021 Publication	3%

Exclude quotes OnExclude matches < 3%Exclude bibliography On