

JHP effect

by Ardiana Ekawanti

Submission date: 07-Apr-2023 12:28AM (UTC-0500)

Submission ID: 2058192906

File name: JHP_effect.pdf (862.75K)

Word count: 5235

Character count: 27508

Effect of Mercury Exposure on Renal Function and Hematological Parameters among Artisanal and Small-scale Gold Miners at Sekotong, West Lombok, Indonesia

Ardiana Ekawanti¹, Baiq Dewi Krisnayanti²

¹ Medical Faculty, University of Mataram, Mataram, Indonesia

² Agriculture Faculty, University of Mataram, Mataram, Indonesia

Corresponding Author:

¹⁶ Dewi Krisnayanti
Agriculture Faculty,
University of Mataram,
Jl. Pendidikan No. 37
Mataram, Indonesia

bdewi.krisnayanti@gmail.com

Introduction

²⁰ Artisanal and small-scale gold mining (ASGM) activities in West Nusa Tenggara (WNT) province began in mid-2009. Hydrargyrum (mercury) amalgamation has been the most common method used to recover gold. Besides the availability of mercury (Hg), this technique is widely used because it is considered to be efficient, effective and affordable.¹ In Indonesia, at least 250,000 miners were estimated to be directly involved in more than 1,000 areas of ASGM, spread throughout the country.² This report quantified 713 locations of illegal small-scale mining throughout Sumatra, Java, Kalimantan and Sulawesi.² In addition, there are around 300,000 to 500,000 people affected by the ASGM activities, the majority of which are non-registered, informal operations

³ **Background.** Mercury is a toxic metal with effects on human health ranging from acute to chronic in a very short time of exposure. Artisanal and small-scale gold mining (ASGM) is the main source of direct human exposure to mercury. Human exposure to mercury (Hg) can occur through both direct inhalation of mercury vapor and consumption of material taken from contaminated areas. To protect the health of ASGM workers and surrounding communities, a health assessment of mercury exposure and its effects is urgently needed. However, analysis of hair and urine samples as a proof test for mercury toxicity is very expensive. Therefore other tests must be considered to identify the first symptoms of mercury toxicity in miners and the surrounding community.

Objectives. The present study aimed to determine the effects of mercury exposure on renal function along with the hematological parameters of gold miners and the community as a first indication of mercury exposure symptoms.

Methods. The study was designed as a purposive field sampling study and was conducted in 3 main villages in Sekotong District, West Lombok Regency, West Nusa Tenggara Province, Indonesia. The 100 subjects were miners that have been exposed to mercury for at least 5 years and their wives and children (non-miners) who lived around the gold processing area. Blood and urine samples were then obtained from the subjects. The miners and non-miners were questioned about their mercury exposure over the previous 5 years, duration of exposure, and how mercury was handled in their daily life. Blood and urine samples were collected at the time of the study, around 10 ml of urine and 0.1 ml of blood (2 drops) were collected per subject. In order to determine the parallel results between the blood-urine and hair results, hair from the miners was collected at a different time for analysis.

Results. The results showed that the subjects had low proteinuria, hemoglobin and hematocrit concentrations as a consequence of chronic mercury intoxication. This finding was parallel with results of high mercury concentrations in urine (>7 – 273.3 µg/l) and miners' hair (>1 – 12.93 µg/g). Miners and non-miners in the exposure area were found to have proteinuria levels of more than 0.3 g/L. Proteinuria (≥0.3 g/L) was observed in 92.6% of miners and 72.4% of non-miners.

Conclusions. The results of the present study suggest that urinalysis of proteinuria and hemoglobin values can be used as a screening test to detect renal impairment due to mercury intoxication.

Competing Interests. The authors declare no competing financial interests.

Keywords. ASGM, proteinuria, hemoglobin, hematocrit

J Health Pollution 9: 25-32 (2015)

located in remote areas in Indonesia.³

On Lombok Island alone, in 2012 there were around 22,500 people estimated to be involved in this activity.⁴

The amalgamation method of ASGM results in Hg emissions that enter the environment in several ways. Mercury is unintentionally or intentionally spilled onto the ground or agriculture land. Atmospheric transport and

deposition at normal temperature is another common way for Hg to enter many water systems. In addition, Hg is often discharged together with other wastes into inadequate tailings ponds, or directly disposed into rivers and water systems. Moreover, when purifying the amalgam by burning, vaporized Hg is released into the atmosphere.⁵ In addition, mercury gets into the human body through the food chain when fish are eaten or used in agricultural products.⁶⁻⁸ Many studies have shown that mercury emissions into the environment affect human health.⁹⁻¹³

Mercury is a highly toxic metal that directly affects the nervous and cardiovascular system. Nausea, vomiting, diarrhea and severe kidney damage may occur due to exposure to high doses of mercury over a short period of time. Hallucinations, memory loss, nerve damage and the inability to concentrate can also occur. Furthermore, mercury toxicity symptoms include tremors, loss of dermal sensitivity, and slurred speech.^{14,15}

Mercury exposure can also come through inhalation of inorganic metallic mercury, ingestion of inorganic complexed mercury, or ingestion of organic forms of mercury. The factors regarding the occurrence and severity effects of mercury on human health include: the chemical form and dose of mercury; the age or the developmental stage of the person exposed, and the duration and exposure route, including inhalation, ingestion, and dermal contact. Fish consumption patterns can also increase the chance of mercury exposure when fish and seafood are contaminated with mercury. Mercury toxicity interferes with vital body systems: the nervous system, kidney function, and cardiovascular system. It has been

Abbreviations			
ASGM	Artisanal and small-scale gold mining	Hg	Mercury
Hb	Hemoglobin	SD	Standard deviation
HCT	Hematocrit	WHO	World Health Organization
		WNT	West Nusa Tenggara

reported that the development of the fetal nervous system is of high risk to the toxic effects of mercury. Other systems that may be affected by mercury toxicity are the respiratory, gastrointestinal, hematologic, immune, and reproductive systems. Following exposure to elemental mercury and methylmercury, kidney damage is the end-point of exposure to inorganic mercury compounds. Exposure damages the secretory organ of erythropoietin—a hormone that stimulates erythrocyte synthesis in the kidney. As kidney function declines, this affects the amount of red blood cells.^{5,11,16,17} In addition, the developing nervous system has been identified as the most sensitive toxicological endpoint.^{5,11,12,17}

Since mercury vapor is easily absorbed into the human body, an assessment of miners and people living in the vicinity of ASGM is urgently needed. However, in Indonesia, the analysis of hair and urine samples as a proof test for total mercury toxicity is expensive, as few laboratories are able to analyze total mercury or even methylmercury. Based on the authors' experience, samples must be sent to a laboratory out of West Nusa Tenggara Province or even overseas. Hence, other tests must be considered to identify the first symptoms of mercury toxicity among miners

in the surrounding community. Bose-O'Reilly et al. recommended four neuropsychological tests to assess memory and motor function problems in ASGM miners.¹⁸ A physical was aimed at identifying neurologic disturbances such as ataxia, tremor and coordination problems related to the neurotoxic effects of mercury. Furthermore, preliminary symptoms of mercury exposure can be determined by examining kidney function through a simple urine test and hemoglobin level can be assessed with a simple blood test kit. Holmes, et al. reported that there was an increased level of excretion of low molecular weight proteins in urine from ASGM workers of as low as 5–10 µg/g creatinine.¹⁹ These creatinine levels were only slightly above those found in the general population. Even though this level was only slightly higher than normal, the significance of such changes in renal excretory profile is of toxicological importance. Furthermore, mercury ingested or inhaled as elemental mercury binds to structural protein-hemoglobin and causes impairment in the erythrocyte function of oxygen carrying capacity and induces hemolysis. Both mechanisms could cause decreases in either hemoglobin (Hb) or hematocrit (HCT) levels.²⁰ These findings could be used to take further action for workers and the entire community




Figure 1 — The study site at Sekotong District West Nusa Tenggara Province

in the vicinity of ASGM. This study aimed to determine the association between mercury exposure, renal function and hematological parameters of gold miners and the surrounding community as a first indication of mercury exposure symptoms.

18 Methods

Study Location

The research was conducted at Sekotong ASGM, West Lombok Regency, West Nusa Tenggara Province, Indonesia (Figure 1). The subjects were miners and families from 3 different villages: Sekotong Tengah, Tawun and Pelangan. The

villages were the main gates to the mining sites.

Sample Collection and Preparation

The samples were recruited using a non-probability sampling technique (purposive sampling) by collecting 33 samples in each village, and in total, 100 people were recruited from the 3 villages. The subject criteria were: 1) miners who had been mining and using mercury for at least 5 years; and 2) non-miners, including wives and children, who had lived in the gold processing area for at least one year. Before conducting the study, ethical clearance was obtained from the Ethical Code Board of the Medical Faculty, University of Mataram,

Indonesia, where subjects were asked to sign an informed consent form in order to be involved in this study. Blood and urine samples were then obtained from the miners and non-miners. Following the study procedure, the miners and non-miners were questioned about their mercury exposure over the previous 5 years, duration of exposure, and how mercury was handled in their daily life. Blood and urine samples were collected at the time of the study, around 10 ml of urine and 0.1 ml of blood (2 drops) were collected per subject. In order to determine the parallel results between the blood-urine and hair results, hair from the miners was collected at a different time for analysis.¹

Sample Analysis

Proteinuria was assessed immediately upon collection, but urine was stored at -4°C until analysis for mercury. Proteinuria level was assessed using dip-stick urinalysis and the results were stated as g/L. The hemoglobin and hematocrit via capillary blood sample were assessed using a hemoglobinometer and stated as mg/dL.

For determining the parallel results of proteinuria and hemoglobin and hematocrit, some of the miners' hair and urine samples were analyzed. Analysis of the hair samples was performed at State Key Laboratory of Environmental Geochemistry in Guiyang, China.¹ Hair samples for total mercury examination were directly assessed using a LUX RA915+ mercury analyzer. The detection limit of the instrument was 0.2–5 ng g⁻¹. The urine samples were directly collected, and placed and sealed in a polyethylene container and kept in the freezer at -20°C until transport to the laboratory of the Institute and Outpatient Clinic for Occupational, Social and Environmental Medicine in Munich,

Characteristics	Percentage (%)
Sex	
Male	75
Female	25
Age (year)	
<18	10
18-60	85
>60	5
Occupation	
Miner (directly exposed)	71
Non-miner (indirectly exposed)	29
Duration of exposure (years)	
Mean	5.5
<1 (n=0)	0
1-5 (n=54)	53
6-10 (n=4)	47
>10 (n=1)	0

Table 1 — Characteristics of Research Participants

Germany. The analysis of mercury in urine was performed with cold-vapor atomic absorption spectrometry and the determination limit was 0.25 µg/l.⁹ Blood and urine data were analyzed using descriptive statistical methods and Spearman's correlation test. It was found that after a short time (4 years) of mercury exposure, the concentration of total Hg in miners' hair and urine from Sekotong ASGM were above the World Health Organization (WHO) standard.^{1,6,8}

Results

Characteristic of Research Participants

This research involved one hundred ASGM workers and their families, the characteristics of whom are shown in Table 1.

Table 1 shows that most participants were male (75%), and 90% were adults and 10% were children. Most participants were miners (71%) working in processes such as amalgamation, cyanidation, and smelting. Hence, they were directly exposed to mercury. Non-miners, the family of miners (wives and children), were considered to be indirectly exposed to mercury, as they lived in the vicinity of the smelting process within a radius of less than 500 m. Their duration of exposure was more than five years (mean = 5.4 years).

Data shown in Table 2 are a compilation of all parameters measured in this research, including urinalysis, hemoglobin, and total mercury in hair and urine. Protein was assessed qualitatively using the dip-stick method and was categorized as negative, trace, +1, +2, +3, or +4 and quantitatively valued as negative, <0.3 g/L, 0.3 g/L, 1 g/L, 3 g/L and >20 g/L. The average amount of protein in urine was 1.56 ± 1.03 g/L (mean SD). This was considered to indicate

Parameter	Level			
	Minimum	Maximum	Mean ± Standard deviation	Median
Proteinuria	1.0	3.0	2.0	2.0
Level of protein in urine (g/L)	1.0	3.0	1.6±1.0	1.0
Hemoglobin (g/dL)				
Male	7.3	25.0	12.9±2.7	12.7
Female	9.6	16.9	13.1±1.5	13.0
Hematocrit (%)	21.9	75.0	38.9±7.3	38.4
Urine mercury level (µg/L)	2.6	178.1	41.0±52.2	19.3
Hair mercury level (µg/g)	0.0	6.6	2.6±1.7	2.2

Table 2 — Characteristics of Proteinuria, Urine Mercury and Hematological Parameters

Proteinuria (g/L)	Percentage (%)
Miners	
0	0
<0.3	0
0.3	7
1	56.3
3	36.3
≥20	0
Non-miners	
0	0
<0.3	0
0.3	27.6
1	48.3
3	24.1
≥20	0

Table 3 — Proteinuria Profile in Sekotong Area ASGMs

proteinuria since this level was higher than 300 mg/L (0.3 g/L). The mean value of hemoglobin found in men (12.9 ± 2.68 g/dL) was lower than in women (13.11 ± 1.45 g/dL). The mean hematocrit value was $38.95 \pm 7.27\%$, and mercury levels in urine and hair were 41.04 ± 52.19 µg/L and 2.56 ± 1.71 (µg/g), respectively.

Table 4 shows that 57.7% of the miners suffered from anemia, as the cut-off point for this value was 13 g/dL among male non-smokers. However, if the cut-off value for anemia among smokers is taken into consideration (13.3 g/dL), 67.6% of the miners would be considered anemic. The results also showed that the Hb values of non-miners were predominantly normal compared to miners, as only 13.8% of non-miners suffered from anemia.

Hemoglobin Level Group	Percentage (%)	Hematocrit Percentage	
		Percentage (%)	Percentage (%)
Miner			
<13 g/dL	57.7	<40%	67.6
≥13 g/dL	42.3	40-50%	28.2
Smoking			
<13.3 g/dL	67.6	≥50%	4.2
≥13.3 g/dL	32.4		
Non-miner			
<12 g/dL	13.8	<40%	51.7
≥12 g/dL	86.2	40-50%	41.4
		≥50%	6.9

Table 4 — Hemoglobin and Hematocrit Levels of Subjects in Sekotong Area ASGMs

Table 5 illustrates the correlations between mercury exposure and several health parameters such as urine protein level, Hb, HCT, urine and hair mercury levels in two different mercury exposure groups, miners and non-miners (23). Spearman's correlation test and the Kruskal-Wallis test were performed to confirm the relationships between the miners and non-miners and health parameters. Table 5 shows that miners and non-miners showed significant differences in all health parameters ($p < 0.05$ Kruskal-Wallis test and $p < 0.05$ Spearman's test) except for hair mercury value, which showed no significant difference between the miners and non-miners.

Discussion

All participants in the present study were directly exposed to mercury, either as workers (miners) or as family members (non-miners) living

in the contaminated atmosphere, water, food stuff and soil in the vicinity of the mining activities. Most participants were directly exposed to mercury, as they worked in the cyanidation process, since this process often used sludge from the amalgamation method which involves mercury.

Some study subjects were children that had direct contact with mercury from mercury panning after school and standing in close proximity to burning processes. This type of child activity has also been reported in other gold mining areas.⁹ Most of the women were of reproductive age, which means that their pregnancies were also at risk, as mercury affects intrauterine growth development, especially brain development.^{5,21,22} The duration of exposure to mercury contaminants was an average of 5.4 years. This period was much shorter than the 14.8 years needed to show

Exposure Group	Urine protein level (g/L)	Hemoglobin (g/dL)	Hematocrit (%)	Urine Mercury Level (µg/L)	Hair Mercury Level (µg/g)
Miners	1.68±1.023	12.74±2.39	38.21±7.18	69.39±62.41	2.77±1.68
Non-miners	1.29±1.03	13.59±2.43	40.77±7.29	12.7±11.50	2.37±1.82
Spearman's rho (ρ); coefficient correlation	0.031; -0.216	0.045; 0.201	0.045; 0.201	0.042; -0.550	0.517; -0.169
Kruskal-Wallis Test	0.032	0.045	0.045	0.047	0.500

Table 5 — Correlation Between Exposure Group and Urine Protein Level, Hemoglobin and Hematocrit, Urine and Hair Mercury Levels

specific clinical manifestations in previous reports.¹¹ This manifestation is affected by the type and dose of mercury, the age or developmental stage, duration of exposure and route of exposure.⁵ A weakness of the present study was that there was no identification of the dose and type of mercury exposure.

Protein levels in urine were obtained by converting the qualitative proteinuria dip-stick results categorized as negative, trace, +1, +2, +3, to quantitative values of 0 g/L, <0.3 g/L, 0.3 g/L, 1 g/L and ≥20 g/L, respectively. Proteinuria was identified by results of +2 and ≥0.3 g/L.²¹ Almost all of the subjects in the exposed areas, both miners and non-miners, were positive for proteinuria (miners, 92.6% and non-miners, 72.4%). This result was similar to findings of proteinuria in patients exposed to products containing mercury.²⁴ In addition, proteinuria is a clinical manifestation of mercury intoxication with elemental, inorganic and ethyl-mercury.¹⁷ High urinary (69.39 ± 62.41 µg/L) and hair (2.77 ± 1.68 µg/g) mercury levels in the miners indicate that a pathological process in the kidney may occur due to mercury exposure. Common causes of urinary

mercury excretion were elemental and organic mercury exposure, as miners were directly exposed to mercury vapor during the melting process.^{10,25} Once inhaled, mercury vapor is dispersed rapidly into the blood and might deposit in other organs e.g. brain, kidney, placenta thyroid and others.^{7,11} Significant differences in urinary levels between miners and non-miners indicate that a possible route of exposure may be ingestion of inorganic mercury through contaminated food or water.¹ In the study location, most of the water sources for amalgamation processes and daily life come from the same well, and often the tailing ponds are nearby the well. Although both groups were directly exposed to mercury, the level of mercury (urinary and hair samples) for the miners group was higher than for non-miners. One contributing factor in this case was smoking habit, as all miners were smokers, and the urinary mercury level of this group was more than five times higher than that of non-miners.¹³

Persistent proteinuria indicates kidney disease and the most common impacts of this condition are diabetes, hypertension, obesity and medicine or chemical substances.²⁶

One of the chemical substances that affects proteinuria is mercury, and to confirm this, repeated tests are required.²⁵ People with consistent positive results can be categorized as having persistent proteinuria and considered to have kidney disease. A weakness of the present study was that there were no repeated tests to confirm persistent proteinuria. Assessment of renal function is needed in future studies.

The mean of the hemoglobin concentration of miners (12.74 ± 2.39 g/dL) was lower than that of either non-miners (13.59 ± 2.43 g/dL) or the normal value for men (13–18 g/dL).⁶ The results of this study also indicated a correlation between the smoking habits of miners and hemoglobin concentration. Since all of the miners were smokers, if a correction factor for smokers (0.3) is used, then Hb values would appear normal (13.04). However, when the normal level for male smokers is used (13.3 d/dL), then the Hb values of the miners were below normal.

Hemoglobin concentration is affected by many factors such as high altitude, diet, pregnancy and smoking.²⁷ Since the Sekotong mining sites are in the lowlands, subjects were not pregnant

and had a homogenous diet, the remaining factors influencing Hb values were smoking and mercury exposure (route and magnitude). This finding was similar to that of previous studies reporting that mercury exposure could decrease hemoglobin levels.^{20,28} The mechanisms behind the decrease in Hb through mercury exposure are not only due to decreasing renal function, but also due to the influence on iron metabolism and haemolysis.^{16,29,30} To examine the iron metabolism factor, iron transfer and ferritin serum must be assessed, which was not performed in the present study. The Hb values and HCT concentration of miners ($38.21 \pm 7.18\%$) were lower than that of non-miners ($40.77 \pm 7.29\%$) as well as the normal value (40–50%). Hematocrit reflects the cell and plasma proportions, and low Hb is usually found in low HCT and vice versa.

Except for hair mercury level, all parameters measured in the present study showed significant differences between the two groups. The subjects in the miner group were men, and most subjects in the non-miner group were women. The results of the present study differed from those of previous reports that found that mercury in men's hair was higher than that in women's.²⁴

Conclusion

After five years of exposure to mercury, people in Sekotong area ASGMs, both miners and non-miners, showed proteinuria and low hemoglobin and hematocrit concentrations due to chronic mercury intoxication as indicated by high urinary and hair mercury levels.

References

- Krisnayanti BD, Anderson CW, Utomo WH, Feng X, Handayanto E, Mudarisna N, Ikram H, Khususiah. Assessment of environmental mercury discharge at a four-year-old artisanal gold mining area on Lombok Island, Indonesia. *J Environ Monit* [Internet]. 2012 Oct 26 [cited 2015 Oct 26];14(10):2598-607. Available from: <http://pubs.rsc.org/en/Content/ArticleLanding/2012/EM/c2em30515a#divAbstract> Subscription required to view.
- Grita A, Primanti A, Zaki K, Ismawati Y. Rapid assessment of the socio-economic impact and human rights aspect of mercury use in artisanal and small-scale gold mining hotspots in Indonesia [Internet]. Denpasar, Indonesia: Balifokus; 2012 Jul [cited 2015 Oct 26]. 46 p. Available from: http://www.academia.edu/4818655/Rapid_Assessment_of_the_social-economy_impact_of_ASGM_and_human_rights_aspect
- Mitigating mercury emissions from artisanal and small scale gold mining in Indonesia: final report [Internet]. New York: Blacksmith Institute; 2013 Aug [cited 2015 Jun 12]. 39 p. Available from: http://www.unep.org/chemicalsandwaste/Portals/9/Mercury/Final%20Report%20Narrative_June2014.pdf
- Krisnayanti BD, Anderson CW. Environmental impact assesment: illegal/informal gold mining in Lombok. Jakarta, Indonesia; GIZ-RED; 2012. 69 p.
- Limbong D, Kumampung J, Rimper J, Arai T, Miyazaki N. Emissions and environmental implications of mercury from artisanal gold mining in North Sulawesi, Indonesia. *Sci Total Environ* [Internet]. 2003 Jan 20 [cited 2015 Oct 26];302(1-3):227-36. Available from: <http://www.sciencedirect.com/science/article/pii/S0048969702003972> Subscription required to view.
- Mercury and health [Internet]. Geneva, Switzerland: World Health Organization; [updated 2013 Sep; cited 2015 Oct 25]. Available from: <http://www.who.int/mediacentre/factsheets/fs361/en/>
- Park JD, Zheng W. Human exposure and health effects of inorganic and elemental mercury. *J Prev Med Public Health* [Internet]. 2012 Nov [cited 2015 Oct 26];45(6):344-52. Available from: <http://jpmph.org/journal/view.php?doi=10.3961/jpmph.2012.45.6.344>
- Krisnayanti BD, Sukartono, Ardiana E. Alternative livelihood: report. Mataram, Indonesia: University of Mataram; 2013.
- Bose-O'Reilly S, Lettmeier B, Gothe RM, Beinhoff C, Siebert U, Drasch G. Mercury as a serious health hazard for children in gold mining areas. *Environ Res* [Internet]. 2008 May [cited 2015 Oct 28];107(1):89-97. Available from: <http://www.sciencedirect.com/science/article/pii/S0013935108000224> Subscription required to view.
- Burbure CD, Buchet JP, Leroyer A, Nisse C, Haguenoer JM, Mutti A, Smerhovsky Z, Cikrt M, Ochocka MT, Razniewska G, Jakubowski M, Bernard A. Renal and neurologic effects of cadmium, lead, mercury, and arsenic in children: evidence of early effects and multiple interactions at environmental exposure levels. *Environ Health Perspect* [Internet]. 2006 Apr [cited 2015 Oct 28];114(4):584-90. Available from: <http://www.ncbi.nlm.nih.gov/pmc/articles/PMC1440785/>
- Franko A, Budihna MV, Fikfak MD. Long-term effects of elemental mercury on renal function in miners of Idrija Mercury Mine. *Ann Occup Hyg* [Internet]. 2005 Aug [cited 2015 Oct 28];49(6):521-7. Available from: <http://annhyg.oxfordjournals.org/content/49/6/521.long>
- Kobal AB, Flisar Z, Miklavcic V, Dizdarevic T, Briski AS. Renal function in miner intermittently exposed to elemental mercury vapour. *Arch Ind Hygiene Toxicol* [Internet]. 2001 Mar [cited 2015 Oct 28];51(4):369-80. Available from: http://hrcak.srce.hr/index.php?show=clanak&id_clanak_jezik=1383
- Hong D, Cho SH, Park SJ, Kim SY, Park SB. Hair mercury level in smokers and its influence on blood pressure and lipid metabolism. *Environ Toxicol Pharmacol* [Internet]. 2013 Jul [cited 2015 Oct 28];36(1):103-7. Available from: <http://www.sciencedirect.com/science/article/pii/S1382668913000562> Subscription required to view.
- Nierenberg DW, Nordgren RE, Chang MB, Siegler RW, Blayney MB, Hochberg F, Toribara TY, Cernichiari E, Clarkson T. Delayed cerebellar disease and death after accidental exposure to dimethylmercury. *N Engl J Med* [Internet]. 1998 Jun 4 [cited 2015 Oct 28];338(23):1672-6. Available from: <http://www.nejm.org/doi/full/10.1056/NEJM199806043382305>
- Mohapatra SP, Nikolova I, Mitchell A. Managing mercury in the great lakes: an analytical review of abatement policies. *J Environ Manage* [Internet]. 2007 Apr [cited 2015 Oct 28];83(1):80-92. Available from: <http://www.sciencedirect.com/science/article/pii/S0301479706000648> Subscription required to view.
- Ribarov SR, Benov LC, Benchev IC. On the

- mechanism of mercury induced hemolysis. *Gen Physiol Biophys* [Internet]. 1983 Apr [cited 2015 Oct 28];2(2):81-4. Available from: http://www.gpb.sav.sk/1983/1983_02_81.pdf
17. **Clarkson TW.** The three modern faces of mercury. *Environ Health Perspect* [Internet]. 2002 Feb [cited 2015 Oct 28];110 Suppl 1:11-23. Available from: <http://www.ncbi.nlm.nih.gov/pmc/articles/PMC1241144/>
18. **Bose-O'Reilly S, McCarty MK, Steckling N, Lettmeier B.** Mercury exposure and children's health. *Curr Probl Pediatr Adolesc Health Care* [Internet]. 2010 Sep [cited 2015 Oct 28];40(8):186-215. Available from: <http://www.ncbi.nlm.nih.gov/pmc/articles/PMC3096006/>
19. **Holmes P, James KA, Levy LS.** Is low-level environmental mercury exposure of concern to human health? *Sci Total Environ* [Internet]. 2009 Dec 20 [cited 2015 Oct 28];408(2):171-82. Available from: <http://www.sciencedirect.com/science/article/pii/S0048969709009061> Subscription required to view.
20. **Maheswaran R, Devapaul A, Muralidharan S, Velmurugan B, Igacimuthu S.** Haematological studies of fresh water fish, *Clarias batrachus* (L.) exposed to mercury chloride. *Int Integr Biol* [Internet]. 2008 Jan [cited 2015 Oct 28];2(1):49-53. Available from: [http://www.researchgate.net/publication/26516925_Haematological_studies_of_fresh_water_fish_Clarias_batrachus_\(L.\)_exposed_to_mercuric_chloride](http://www.researchgate.net/publication/26516925_Haematological_studies_of_fresh_water_fish_Clarias_batrachus_(L.)_exposed_to_mercuric_chloride)
21. **Clarkson TW, Magos L, Myers GJ.** The toxicology of mercury--current exposures and clinical manifestations. *N Engl J Med* [Internet]. 2003 Oct 30 [cited 2015 Oct 28];349(18):1731-7. Available from: <http://www.nejm.org/doi/full/10.1056/NEJMra022471> Subscription required to view.
22. **Mahaffey KR.** Mercury exposure: medical and public health issues. *Trans Am Clin Climatol Assoc* [Internet]. 2005 [cited 2015 Oct 28];116:127-54. Available from: <http://www.ncbi.nlm.nih.gov/pmc/articles/PMC1473138/pdf/tacca11600127.pdf>
23. **Silva DA, Halstead AC, Cote AM, Sabr Y, Dadelszen P, Magee LA.** Urinary dipstick proteinuria testing: does automated strip analysis offer an advantage over visual testing? *J Obstet Gynaecol Can* [Internet]. 2014 Jul [cited 2015 Oct 28];36(7):605-12. Available from: http://www.researchgate.net/publication/263964224_Urinary_dipstick_proteinuria_testing_-_Does_automated_strip_analysis_offer_an_advantage_over_visual_testing
24. **Li SJ, Zhang SH, Chen HP, Zeng CH, Zheng CX, Li LS, Liu ZH.** Mercury-induced membranous nephropathy: clinical and pathological features. *Clin J Am Soc Nephrol* [Internet]. 2010 Mar [cited 2015 Oct 28];5(3):439-44. Available from: <http://www.ncbi.nlm.nih.gov/pmc/articles/PMC2827581/>
25. **Abdenour C, Khelili K, Boulakoud MS, Nezzal A, Boubil S, Slimani S.** Urinary markers of workers chronically exposed to mercury vapor. *Environ Res* [Internet]. 2002 Jul [cited 2015 Oct 28];89(3):245-9. Available from: <http://www.sciencedirect.com/science/article/pii/S0013935102943689> Subscription required to view.
26. **Interpreting urine dipstick test in adults: a reference guide for primary care.** *Best Tests* [Internet]. 2013 Jun [cited 2015 Oct 28];19:10-20. Available from: <http://www.bpac.org.nz/BT/2013/June/docs/BT19-pages-10-21.pdf>
27. **Haemoglobin concentration for the diagnosis of anaemia and assessment of severity** [Internet]. Geneva, Switzerland: World Health Organization; 2011 [cited 2015 Oct 28]. 6 p. Available from: <http://www.who.int/vmnis/indicators/haemoglobin/en/>
28. **Calisi A, Lionetto MG, Hernandez JC, Schettino T.** Effect of heavy metal exposure on blood haemoglobin concentration and methemoglobin percentage in *Lumbricus terrestris*. *Ecotoxicology* [Internet]. 2011 Jun [cited 2015 Oct 28];20(4):847-54. Available from: <http://link.springer.com/article/10.1007%2Fs10646-011-0641-1> Subscription required to view.
29. **Saljooghi AS, Mendi FD.** The effect of mercury in iron metabolism in rats. *Clin Toxicol* [Internet]. 2013 [cited 2015 Oct 28];3(5):1-5. Available from: <http://www.omicsonline.org/the-effect-of-mercury-in-iron-metabolism-in-rats-2161-0495.S3-006.pdf>
30. **Chitra S, Jayaprakash K.** The effect of mercury on blood component of fresh water edible fish *Labeo rohita*. *J Acad Indus Res* [Internet]. 2013 May [cited 2015 Oct 28];1(12):774-7. Available from: <http://jairjp.com/MAY%202013/11%20CHITRA.pdf>

JHP effect

ORIGINALITY REPORT

17%

SIMILARITY INDEX

11%

INTERNET SOURCES

12%

PUBLICATIONS

%

STUDENT PAPERS

PRIMARY SOURCES

- | | | |
|---|--|----|
| 1 | Raafat Abdeldayem. "Domestic water and accumulating mercury toxicity in the kidney", <i>Applied Water Science</i> , 2022
Publication | 2% |
| 2 | meridian.allenpress.com
Internet Source | 2% |
| 3 | link.springer.com
Internet Source | 1% |
| 4 | Baiq Dewi Krisnayanti, Christopher W. N. Anderson, Wani Hadi Utomo, Xinbin Feng et al. "Assessment of environmental mercury discharge at a four-year-old artisanal gold mining area on Lombok Island, Indonesia", <i>Journal of Environmental Monitoring</i> , 2012
Publication | 1% |
| 5 | docplayer.net
Internet Source | 1% |
| 6 | www.safasrl.it
Internet Source | 1% |
-

7	Siti Thomas Zulaikhah, Joko Wahyuwibowo, Arrizki Azka Pratama. "Mercury and its effect on human health: a review of the literature", International Journal of Public Health Science (IJPHS), 2020 Publication	1 %
8	www.semanticscholar.org Internet Source	1 %
9	D LIMBONG, J KUMAMPUNG, J RIMPER, T ARAI, N MIYAZAKI. "Emissions and environmental implications of mercury from artisanal gold mining in north Sulawesi, Indonesia", The Science of The Total Environment, 2003 Publication	1 %
10	www.ncbi.nlm.nih.gov Internet Source	1 %
11	www.frontiersin.org Internet Source	1 %
12	www.researchgate.net Internet Source	1 %
13	Justice Afrifa, Samuel Essien-Baidoo, Richard K.D. Ephraim, Daniel Nkrumah, Daniel Osei Dankyira. "Reduced egfr, elevated urine protein and low level of personal protective equipment compliance among artisanal small	<1 %

scale gold miners at Bibiani-Ghana: a cross-sectional study", BMC Public Health, 2017

Publication

14	biomed surg.com Internet Source	<1 %
15	www.biomedcentral.com Internet Source	<1 %
16	www.mdpi.com Internet Source	<1 %
17	www.scielo.br Internet Source	<1 %
18	www.rangeland.ir Internet Source	<1 %
19	Jarosinska, D.. "Urinary mercury and biomarkers of early renal dysfunction in environmentally and occupationally exposed adults: A three-country study", Environmental Research, 200810 Publication	<1 %
20	www.stepslinen.co.za Internet Source	<1 %
21	aoemj.org Internet Source	<1 %
22	Agus Abhi Purwoko, Veni Rori Setiawati, Saprizal Hadisaputra. "Metal-pigment complex derived from natural dye of	<1 %

anthocyanin: a potential candidate for DSSC photosensitizer", IOP Conference Series: Materials Science and Engineering, 2019

Publication

23

Arifin, Yayu, Masayuki Sakakibara, and Koichiro Sera. "Impacts of Artisanal and Small-Scale Gold Mining (ASGM) on Environment and Human Health of Gorontalo Utara Regency, Gorontalo Province, Indonesia", Geosciences, 2015.

Publication

<1 %

24

Comprehensive Guide to Autism, 2014.

Publication

<1 %

25

E. S. Prasedya, Y. Ambana, N. W. R. Martyasari, Ye'muh Aprizal, Nurrijawati, Sunarpi. "Short-term E-cigarette toxicity effects on brain cognitive memory functions and inflammatory responses in mice", Toxicological Research, 2020

Publication

<1 %

26

Kira Taux, Thomas Kraus, Andrea Kaifie. "Mercury Exposure and Its Health Effects in Workers in the Artisanal and Small-Scale Gold Mining (ASGM) Sector—A Systematic Review", International Journal of Environmental Research and Public Health, 2022

Publication

<1 %

27

Paruchuri, Y.. "Occupational and environmental mercury exposure among small-scale gold miners in the Talensi-Nabdam District of Ghana's Upper East region", *Science of the Total Environment*, 20101115

Publication

<1 %

28

Siska Nurfitriani, Endang Arisoesilaningih, Yulia Nuraini, Eko Handayanto. "Bioaccumulation of Mercury by Bacteria Isolated from Small Scale Gold Mining Tailings in Lombok, Indonesia", *Journal of Ecological Engineering*, 2020

Publication

<1 %

29

ueaeprints.uea.ac.uk

Internet Source

<1 %

30

Li, P.. "Mercury exposures and symptoms in smelting workers of artisanal mercury mines in Wuchuan, Guizhou, China", *Environmental Research*, 200805

Publication

<1 %

31

Nakazawa, Koyomi, Osamu Nagafuchi, Tomonori Kawakami, Takanobu Inoue, Kuriko Yokota, Yuka Serikawa, Basir Cyio, and Rosana Elvince. "Human health risk assessment of mercury vapor around artisanal small-scale gold mining area, Palu city, Central Sulawesi,

<1 %

Indonesia", Ecotoxicology and Environmental Safety, 2016.

Publication

Exclude quotes On

Exclude matches Off

Exclude bibliography On