

# Antibiotics susceptibility of Escherichia coli isolates from clinical specimens before and during COVID-19 pandemic

*by* Eustachius Hagni Wardoyo

---

**Submission date:** 01-May-2023 10:04PM (UTC-0500)

**Submission ID:** 2081624734

**File name:** LampB3\_Publikasi\_IJM\_Wardoyo\_2021.pdf (577.79K)

**Word count:** 2775

**Character count:** 15236

## Antibiotics susceptibility of *Escherichia coli* isolates from clinical specimens before and during COVID-19 pandemic

Eustachius Hagni Wardoyo<sup>1,2\*</sup>, I Wayan Suardana<sup>3</sup>, I Wayan Putu Sutirta Yasa<sup>4</sup>, I Dewa Made Sukrama<sup>5</sup>

<sup>1</sup>Department of Microbiology, School of Medicine, Universitas Mataram, Mataram, Indonesia

<sup>2</sup>Department of Doctoral Studies, School of Medicine, Universitas Udayana, Bali, Indonesia

<sup>3</sup>Department of Microbiology, School of Veterinary Medicine, Universitas Udayana, Bali, Indonesia

<sup>4</sup>Department of Clinical Pathology, School of Medicine, Universitas Udayana, Bali, Indonesia

<sup>5</sup>Department of Microbiology, School of Medicine, Universitas Udayana, Bali, Indonesia

Received: November 2020, Accepted: March 2021

### ABSTRACT

**Background and Objectives:** *Escherichia coli* is a Gram-negative organism causing mild to severe infections, with a wide spectrum range of organs involved. The study aimed to describe antibiotics susceptibility of *E. coli* from clinical specimens from October 11, 2019 to September 11, 2020.

**Materials and Methods:** Study was conducted retrospectively in a private microbiology laboratory in Mataram Indonesia. Period of study divided as two groups after WHO declared COVID-19 as pandemic by March 11, 2020; group A including the specimen related to September 2019 to March 11<sup>th</sup> 2020 and group B including the specimens related to March 11<sup>th</sup> 2020 to September 2020. All clinical specimens were subjected to identify *E. coli* isolates and their antibiotics susceptibility using WHO-NET 5.6 version.

**Results:** Totally, 148 *E. coli* isolates were found in group A and 62 isolates in group B. Prevalence of extended-spectrum beta lactamase (ESBL)-producing *E. coli* in group A was 50% and in group B was 20.9% with significantly difference ( $p < 0.05$ ). There was increase in susceptibility to 10/16 antibiotics; where 3 antibiotics ofloxacin, aztreonam, and fosfomycin were significant ( $p < 0.05$ ). There was a significant decrease in susceptibility to the antibiotics piperacillin ( $p = 0.012$ ), amoxicillin ( $p = 0.002$ ), cefadroxil ( $p = 0.036$ ) and ampicillin ( $p = 0.036$ ). Type of infections between two groups: musculoskeletal infections, pneumonia, urinary tract infections and sepsis were not significant.

**Conclusion:** Reduced number of *E. coli* isolates between two groups with decrease of ESBL-producing *E. coli* contribute in dynamics of antibiotics susceptibility. The longer period of analysis is needed to be done, due to the ongoing COVID-19 pandemic.

**Keywords:** Antimicrobial resistance; COVID-19 pandemic; *Escherichia coli*

\*Corresponding author: Eustachius Hagni Wardoyo, MSc, Department of Microbiology, School of Medicine, Universitas Mataram, Mataram, Indonesia; Department of Doctoral Studies, School of Medicine, Universitas Udayana, Bali, Indonesia.  
Tel: +62-370640874  
Fax: +62-370641717  
Email: wardoyo.eh@unram.ac.id

### INTRODUCTION

Antimicrobial resistance (AMR) is a complex health problem globally. The complexity is described as the cross boundary of AMR between hosts and interspecies within microorganisms. Resistance itself means a change in the response of pathogens to antibiotics which results in ineffective antibiotics, which

Copyright © 2021 The Authors. Published by Tehran University of Medical Sciences.



This work is licensed under a Creative Commons Attribution-Non Commercial 4.0 International license

(<https://creativecommons.org/licenses/by-nc/4.0/>). Noncommercial uses of the work are permitted, provided the original work is properly cited.

be used early when antibiotic is used (1, 2).

*Escherichia coli* is Gram negative bacterium causing urinary tract infection, enteritis, soft tissue infection, septicemia and other types of clinical infections. They can spread rapidly in community and hospital settings (3-5) and may be multi-drug resistant and capable of breaking down commonly used antibiotics. The extended-spectrum beta lactamase (ESBL) producing strains of *E. coli* is one of the important forms of multi-drug resistant phenotypes and categorized as a serious threat by CDC (3). ESBL-producing *E. coli* has experienced increased resistance to a wide variety of penicillins and cephalosporins (6).

There has been an increasing trend in antimicrobial resistance and mortality caused by *E. coli* infections. Between 1997-2011, in the US recorded ESBL-*E. coli* incidence from 0 per 10,000 patient-days to 16.4 per 10,000 patient-days discharged from hospitals (7). The incidence of bacteremia with *E. coli* in the UK has increased from 67.0 cases per 100,000 population in 2015 to 78.0 cases per 100,000 population in 2019 (8). In Indonesia, the prevalence of ESBL-producing *E. coli* in 4 major hospitals has increased from 29% in 2005 to 52% in 2012 (9, 10).

COVID-19 pandemic that has hit the world has affected various aspects of life, socio-economic, health policy, healthcare system, including the impact on bacterial patterns and antibiotic sensitivity (11, 12). The way of COVID-19 affected antimicrobial resistance comes from inappropriate antibiotics prescribing, shifting in antimicrobial stewardship programs, disruption in health services, and the wide use of environmental and personal disinfection (11). The resistance pattern of *E. coli* is interesting to study because of the following reasons: i) It is common Gram-negative microorganisms both as intestinal flora and hospital pathogen, ii) it causes a broad spectrum of infections. This study aimed to describe antibiotic susceptibility of *E. coli* isolates associated with two occasions, before and during the COVID-19 pandemic.

## MATERIALS AND METHODS

**Specimens.** Specimens consisting of blood, sputum, pus and urine were received in the microbiology laboratory of a private hospital in Mataram, Indonesia. This is a referral laboratory and gives services to

surrounding private and public hospitals and private practices.

**Cultures and antibiotic susceptibility tests.** Cultures and antibiotic susceptibility tests were performed according to Clinical and Laboratory Standard Institute (CLSI, 2018) (13). Microorganism identification was determined using the API20E biochemical identification system (BioMerieux France). The susceptibility of microorganisms to antibiotics was assessed using a disc diffusion method. The antibiotic susceptibility data was analyzed using WHO-NET Version 5.6 program.

**Antibiotics.** Our laboratory uses 21 antibiotics for Gram negative antibiotics panels, but only 16 were consistently available (stock shortage). The following antibiotic discs were used: penicillin (P) 10 U, amoxicillin (AMX) 20 µg, cefixime (CFM) 5 µg, ciprofloxacin (CIP) 5 µg, cefadroxil (CFR) 12 µg, amikacin (AMK) 30 µg, ceftriaxone (CRO) 30 µg, ampicillin (AMP) 10 µg, amoxicillin/clavulanic acid (AMC) 20/10 µg, cefuroxime (CXM) 30 µg, cefoperazone/sulbactam (CFP) 75/30 µg, levofloxacin (LVX) 5 µg, fosfomycin (FOS) 36 µg, ofloxacin (OFX) 5 µg, piperacillin (PRL) 100 µg and aztreonam (ATM) 30 µg.

## RESULTS

There were 308 Gram negative isolates in group A and 243 Gram negative isolates in group B. *E. coli* decreases proportionally in both groups from 48% in group A to 25.5% in group B (Table 1).

**Antibiotic susceptibility.** Susceptibility to most of antibiotics (10/16) in group B was higher than group A; these antibiotics were cefoperazone/sulbactam, cefuroxime, ceftriaxone, cefixime, aztreonam, amikacin, ciprofloxacin, ofloxacin, levofloxacin and fosfomycin. Five antibiotics were decreased in susceptibility: ampicillin, amoxicillin, piperacillin, amoxicillin/clavulanic acid and cefadroxil. In both groups, all isolates were resistant to penicillin (Table 2).

Among increased susceptibility antibiotics, three antibiotics fosfomycin, ofloxacin and aztreonam were significant ( $p < 0.05$ ). While in decreased susceptibility antibiotics, four antibiotics ampicillin, amoxicillin, piperacillin and cefadroxil were significant, while amoxicillin/clavulanic acid was not significant.

**Phenotypic ESBL producers.** Extended spectrum beta lactamase (ESBL)-producing *E. coli* was phenotypically identified. There were 74/148 (50%) ESBL-producing *E. coli* in group A and 13/62 (20.9%) in group B, which showed there was a significant decrease ( $p=0.000$ ).

**Type of infections.** Type of infections in group A, musculoskeletal 68/148 (45.9%), pneumonia 32/148 (21.6%), urinary tract infection 44/148 (29.7%) and sepsis 4/148 (2.7%). In group B soft tissue infection 27/62 (43.5%), pneumonia 20/62 (32.3%), urinary tract infection 15/62 (24.1%), sepsis 0. Chi-

**Table 1.** Gram negatives microorganisms isolated from group A and B

	Group A N (%)	Group B N (%)
<i>Enterobacter aerogenes</i>	0 (0)	10 (4.11)
<i>Enterobacter cloacae</i>	74 (24)	51 (20.9)
<i>Escherichia coli</i>	148 (48)	62 (25.5)
<i>Klebsiella pneumoniae</i>	42 (13.6)	21 (8.6)
<i>Pseudomonas aeruginosa</i>	21 (6.8)	80 (32.9)
<i>Serratia marcescens</i>	10 (3.2)	11 (4.5)
<i>Stenotrophomonas maltophilia</i>	0 (0)	1 (0.4)
Other	13 (4.2)	7 (2.8)
Total	308	243

**Table 2.** Prevalence of susceptible *E. coli* isolates in each group. NA=not applicable

	No. of susceptible <i>E. coli</i> isolates in group A (%) n = 148	No. of susceptible <i>E. coli</i> isolates in group B (%) n = 62	P value
Penicillin	0	0	NA
Ampicillin	11 (7.4)	0	0.036
Amoxicillin	18 (12.2)	0	0.002
Piperacillin	13 (8.7)	0	0.012
Amoxicillin/clavulanic acid	28 (18.6)	9 (14.5)	0.553
Cefadroxil	11 (7.4)	0	0.036
Cefoperazone/sulbactam	101 (68.2)	44 (70.9)	0.746
Cefuroxime	20 (13.5)	9 (14.5)	0.829
Ceftriaxone	32 (21.6)	18 (29)	0.287
Cefixime	46 (31.1)	21 (33.9)	0.746
Aztreonam	56 (37.8)	41 (66.1)	0.000
Amikacin	86 (58.1)	44 (70.9)	0.088
Ciprofloxacin	63 (42.6)	35 (56.4)	0.071
Levofloxacin	83 (56.1)	38 (61.3)	0.542
Ofloxacin	81 (54.7)	50 (80.6)	0.000
Fosfomycin	104 (70.3)	53 (85.5)	0.023

square test of both groups: musculoskeletal infection ( $p=0.764$ ), pneumonia ( $p=0.116$ ), urinary tract infection ( $p=0.502$ ) and sepsis ( $p=0.322$ ). All kinds of infection were not significant.

## DISCUSSION

The prevalence of extended spectrum beta-lactamase-producing *E. coli* has increased worldwide, with the highest rate in South Asia (11.5%) and Turkey (10.9%) before and after 2010 (14). The time separation before and during COVID-19 pandemic is important because the introduction of the social distancing policy simultaneously worldwide will automatically limit people's visits to the hospital, leading to a decrease in the number of requests for culture and susceptibility tests to the microbiology laboratory.

The actual impact of COVID-19 pandemic is an ongoing issue. Van Duin et al. (2020) stated there are two sides to debate regarding how COVID-19 affected antimicrobial resistance profile: increase or decrease antimicrobial resistance event. The decreased incidence of AMR is due to a direct impact of reduced outpatient / inpatient visits due to the implementation of social distancing policies carried out by the government. Increasing infection and preven-

tion control practice in community and hospital will reduce the spread of resistance (15).

Broad spectrum antibiotics are often used in a case of suspected secondary bacterial infection in COVID-19 patients, thus affecting unnecessary prescribing. The most common type of infection is pneumonia, followed by blood-stream and urinary tract infection (16).

This study describes the incidence of decreased antimicrobial resistance as a direct result of the implementation of the social distancing policy implemented by the government in the first 6 months of the COVID-19 pandemic.

Overall number of Gram-negative isolates were reduced between two groups, corresponding to reduced number outpatients/inpatients admission to the hospital (Table 1). Antibiotics susceptibility is dynamically changed between two groups, with the majority of antibiotics increase in susceptibility. This phenomenon is likely correlated with reduced consumption of antibiotics, similarly to restrict use of antibiotics is applied, then susceptibility tends to increase (17, 18). Increased number of aztreonam susceptible isolates seems to correspond to decreased number of ESBL-*E. coli* between two groups. Fosfomycin as one of drugs of choice for urinary tract infection (UTI) usage tends to decrease as lower number of UTI cases in group B, that has outcome in increasing its susceptibility.

Type of infections dominance between two groups showed difference in sequence, in group A, soft tissue infection, UTI, pneumonia and sepsis while in group B, pneumonia replace UTI position as runner up and no sepsis cases. This type is differed from elsewhere that urinary tract infections dominance the clinical samples (14, 19, 20). The number of pneu-

monia cases shifting from third in group A to second place in group B is not confirmed by the number of pulmonary manifestations of COVID-19, that is still needed for further investigation.

Collignon and Beggs (15) stated there were difference between developed and developing countries in resistance spread during COVID-19 pandemic, developed countries tends to decrease spread of resistance, while in developing countries with poor control on the spread of bacteria will increase, but in our study shows tendency resistance spread to decrease within given period.

The limitation of our study is the short period of time and limited area is not sufficient to draw a big picture impact of COVID-19 pandemic to *E. coli* susceptibility pattern. The subject in group B had no further investigation of SARS-CoV-2 infection status. The azithromycin antibiotics were not evaluated in this study.

CONCLUSION

Total *E. coli* isolates and ESBL-producing *E. coli* were in decline during COVID-19 pandemic. This phenomenon contributes to the dynamic of antibiotic susceptibility between two groups. longer period of analysis is needed to be done, due to the ongoing COVID-19 pandemic.

REFERENCES

1. WHO. Fact sheet Antibiotic Resistance. 2020. Available at: <https://www.who.int/news-room/fact-sheets/detail/antibiotic-resistance>
2. Aslam B, Wang W, Arshad MI, Khurshid M, Muzam-




Fig. 2. COVID-19 and antimicrobial resistance. AMR=antimicrobial resistance. Developed according to reference 11.

- mil S, Rasool MH, et al. Antibiotic resistance: a run-down of a global crisis. *Infect Drug Resist* 2018;11:1645-1658.
3. CDC. Antibiotic Resistance Threats in the United States 2019. <https://www.cdc.gov/drugresistance/pdf/threats-report/2019-ar-threats-report-508.pdf>
  4. Breijyeh Z, Jubeh B, Karaman R. Resistance of Gram-negative bacteria to current antibacterial agents and approaches to resolve it. *Molecules* 2020; 25:1340.
  5. Allocati N, Masulli M, Alexeyev MF, Di Ilio C. *Escherichia coli* in Europe: an overview. *Int J Environ Res Public Health* 2013;10:6235-6254.
  6. Ventola CL. The antibiotic resistance crisis: part 1: causes and threats. *P T* 2015;40:277-283.
  7. McDanel J, Schweizer M, Crabb V, Nelson R, Samore M, Khader K, et al. Incidence of extended-spectrum  $\beta$ -lactamase (ESBL)-producing *Escherichia coli* and *Klebsiella* infections in the United States: A systematic literature review. *Infect Control Hosp Epidemiol* 2017; 38:1209-1215.
  8. Public Health England (2020). English surveillance programme for antimicrobial utilisation and resistance (ESPAUR) 2015 to 2019.
  9. Hadi U, Kuntaman, Qiptiyah M, Paraton H. Problem of antibiotic use and antimicrobial resistance in Indonesia: are we really making progress? *IJTID* 2013; 4: 5-8.
  10. Hayati Z, Rizal S, Putri R. Isolation of extended-spectrum  $\beta$ -lactamase (ESBL) producing *Escherichia coli* and *Klebsiella pneumoniae* from Dr. Zainoel Abidin General Hospital, Aceh. *Int J Trop Vet Biomed Res* 2019;4:16-22.
  11. van Duin D, Barlow G, Nathwani D. The impact of the COVID-19 pandemic on antimicrobial resistance: a debate. *JAC Antimicrob Resist* 2020; 2: dlaa053.
  12. Getahun H, Smith I, Trivedi K, Paulin S, Balkhy HH. Tackling antimicrobial resistance in the COVID-19 pandemic. *Bull World Health Organ* 2020;98:442-442A.
  13. CLSI (2019). Performance Standards for Antimicrobial Susceptibility Testing. 29<sup>th</sup> ed. CLSI Supplement M100. Wayne PA: Clinical and Laboratory Institute.
  14. Lee DS, Lee SJ, Choe HS. Community-acquired urinary tract infection by *Escherichia coli* in the era of antibiotic resistance. *Biomed Res Int* 2018;2018: 7656752.
  15. Collignon P, Beggs JJ. CON: COVID-19 will not result in increased antimicrobial resistance prevalence. *JAC Antimicrob Resist* 2020; 2: dlaa051.
  16. Clancy CJ, Buehrle DJ, Nguyen MH. PRO: The COVID-19 pandemic will result in increased antimicrobial resistance rates. *JAC Antimicrob Resist* 2020; 2: dlaa049.
  17. Fair RJ, Tor Y. Antibiotics and bacterial resistance in the 21<sup>st</sup> century. *Perspect Medicin Chem* 2014; 6: 25-64.
  18. Manyi-Loh C, Mamphweli S, Meyer E, Okoh, A. Antibiotic use in agriculture and its consequential resistance in environmental sources: potential public health implications. *Molecules* 2018; 23:795.
  19. Kibret M, Abera B. Antimicrobial susceptibility patterns of *E. coli* from clinical sources in northeast Ethiopia. *Afr Health Sci* 2011; 11(Suppl 1): S40-S45.
  20. Kulkarni SR, Peerapur BV, Sailesh KS. Isolation and antibiotic susceptibility pattern of *Escherichia coli* from urinary tract infections in a tertiary care hospital of North Eastern Karnataka. *J Nat Sci Biol Med* 2017; 8:176-180.

# Antibiotics susceptibility of Escherichia coli isolates from clinical specimens before and during COVID-19 pandemic

## ORIGINALITY REPORT

**23%**  
SIMILARITY INDEX

**13%**  
INTERNET SOURCES

**20%**  
PUBLICATIONS

**%**  
STUDENT PAPERS

## PRIMARY SOURCES

- 1** Mahshid Lalvand, Seyed Jamal Hashemi, Mansour Bayat. "Effect of fluconazole and terbinafine nanoparticles on the treatment of dermatophytosis induced by Trichophyton mentagrophytes in guinea pig", Iranian Journal of Microbiology, 2021  
Publication **2%**
- 2** [ijm.tums.ac.ir](http://ijm.tums.ac.ir)  
Internet Source **2%**
- 3** Lucky H. Moehario. "Trend of antibiotics susceptibility of multidrugs resistant Pseudomonas aeruginosa in Jakarta and surrounding areas from 2004 to 2010", African Journal of Microbiology Research, 2012  
Publication **1%**
- 4** [daten-quadrat.de](http://daten-quadrat.de)  
Internet Source **1%**
- 5** [dspace.bracu.ac.bd](http://dspace.bracu.ac.bd)  
Internet Source **1%**

6	"Posters", <i>Clinical Microbiology and Infection</i> , 2007	1 %
Publication		
7	www.ncbi.nlm.nih.gov	1 %
Internet Source		
8	Eloísa Sevilla, Clara Marín, José F. Delgado - Blas, Bruno González - Zorn et al. " Wild griffon vultures ( ) fed at supplementary feeding stations: Potential carriers of pig pathogens and pig - derived antimicrobial resistance? ", <i>Transboundary and Emerging Diseases</i> , 2020	1 %
Publication		
9	Osman Sianipar, Widya Asmara, Iwan Dwiprahasto, Budi Mulyono. "Mortality risk of bloodstream infection caused by either <i>Escherichia coli</i> or <i>Klebsiella pneumoniae</i> producing extended-spectrum $\beta$ -lactamase: a prospective cohort study", <i>BMC Research Notes</i> , 2019	1 %
Publication		
10	Ibrahim A Naqid, Amer A Balatay, Nawfal Rasheed Hussein, Kurdistan Abdullah Saeed, Hiba Abdulaziz Ahmed, Shivan Hassan Yousif. "Antibiotic Susceptibility Pattern of <i>Escherichia coli</i> Isolated from Various Clinical Samples in Duhok City, Kurdistan Region of Iraq", <i>International Journal of Infection</i> , 2020	1 %



11 D. Dubois, J. Delmas, A. Cady, F. Robin, A. Sivignon, E. Oswald, R. Bonnet. "Cyclomodulins in Urosepsis Strains of Escherichia coli", Journal of Clinical Microbiology, 2010  
Publication 1 %

---

12 [www.scielo.br](http://www.scielo.br)  
Internet Source 1 %

---

13 [garuda.kemdikbud.go.id](http://garuda.kemdikbud.go.id)  
Internet Source 1 %

---

14 Usman Abubakar, Menier Al-Anazi, Zainab alanazi, Jesús Rodríguez-Baño. "Impact of COVID-19 pandemic on multidrug resistant gram positive and gram negative pathogens: A systematic review", Journal of Infection and Public Health, 2023  
Publication 1 %

---

15 [mdpi-res.com](http://mdpi-res.com)  
Internet Source 1 %

---

16 Dirayah Rauf Husain, Rihuh Wardhani. "Antibacterial activity of endosymbiotic bacterial compound from Pheretima sp. earthworms inhibit the growth of Salmonella Typhi and Staphylococcus aureus: in vitro and in silico approach", Iranian Journal of Microbiology, 2021 <1 %

17

Shena Y. C. Lim, Yvonne P. Zhou, Daphne Yii, De Zhi Chin et al. "Stemming the Rise of Antibiotic Use for Community-Acquired Acute Respiratory Infections during COVID-19 Pandemic", *Antibiotics*, 2022

Publication

<1 %

18

[www.medrxiv.org](http://www.medrxiv.org)

Internet Source

<1 %

19

Sidra Khan, Syed Shahzad Hasan, Stuart E. Bond, Barbara R Conway, Mamoon A. Aldayeb. "Antimicrobial consumption in patients with COVID-19: A systematic review and meta-analysis", *Expert Review of Anti-infective Therapy*, 2021

Publication

<1 %

20

[ejmed.org](http://ejmed.org)

Internet Source

<1 %

21

[repository.unair.ac.id](http://repository.unair.ac.id)

Internet Source

<1 %

22

[www.dovepress.com](http://www.dovepress.com)

Internet Source

<1 %

23

Asha Pai K. B, Brinda Rajagopalan. "Susceptibility of Urinary Escherichia coli to Fosfomycin and Nitrofurantoin- Re-Exploration of Old and Forgotten Oral

<1 %

# Antibiotics", Journal of Evolution of Medical and Dental Sciences, 2020

Publication

---

24

Cornelius J Clancy, Deanna J Buehrle, M Hong Nguyen. "PRO: The COVID-19 pandemic will result in increased antimicrobial resistance rates", JAC-Antimicrobial Resistance, 2020

Publication

---

<1 %

25

I Putu Gde Surya Adhitya, Wen-Yu Yu, Paul Bass, Gede Parta Kinandana, Mau-Roung Lin. "Effects of Kinesio Taping and Transcutaneous Electrical Nerve Stimulation Combined With Active Stretching on Hamstring Flexibility", Journal of Strength and Conditioning Research, 2021

Publication

---

<1 %

26

Marianna Savenko, Maryna Kryvtsova. "Urban Aquatic Ecosystems as a Factor of the Spread of Antibiotic Resistant Microorganisms and Resistance Genes", Transylvanian Review of Systematical and Ecological Research, 2021

Publication

---

<1 %

27

Marija Djukic, Nadine Sostmann, Thomas Bertsch, Marianne Mecke et al. "Vitamin D deficiency decreases survival of bacterial meningoencephalitis in mice", Journal of Neuroinflammation, 2015

Publication

---

<1 %

28

Robert Sinto, Khie Chen Lie, Siti Setiati, Suhendro Suwanto et al. "Blood culture utilization and epidemiology of antimicrobial-resistant bloodstream infections before and during the COVID-19 pandemic in the Indonesian national referral hospital", Antimicrobial Resistance & Infection Control, 2022

Publication

<1 %

29

Vikas Ambiya, Taraprasad Das, Savitri Sharma, Jay Chhablani, Vivek Dave, Subhadra Jalali, Raja Narayanan, Joveeta Joseph. "Comparison of clinico-microbiological profile and treatment outcome of in-house and referred post cataract surgery endophthalmitis in a tertiary care center in South India", Journal of Ophthalmic Inflammation and Infection, 2016

Publication

<1 %

30

[academic.oup.com](https://academic.oup.com)

Internet Source

<1 %

31

"13th European Congress of Clinical Microbiology and Infectious Diseases", Clinical Microbiology and Infection, 2003

Publication

<1 %

32

Ida Bagus Subanada, I. Made Bakta, I. Wayan Bikin Suryawan, Putu Astawa, Bagus Komang Satriyasa. "Association between Vitamin D

<1 %

Level, Vitamin D Receptor Gene Polymorphisms, and Cathelicidin Level to Acute Lower Respiratory Infections, and the Picture of Exon 2-Vitamin D Receptor Gene Polymorphisms in Children under 5 years old", Open Access Macedonian Journal of Medical Sciences, 2020

Publication

---

33

Sipei Wang, Sheng Zhao, Yangxiao Zhou, Shanshan Jin, Tinghua Ye, Xinling Pan. " Antibiotic resistance spectrum of strains from different samples and age-grouped patients: a 10-year retrospective study ", BMJ Open, 2023

Publication

---

<1 %

34

[digitalcommons.jsu.edu](https://digitalcommons.jsu.edu)

Internet Source

---

<1 %

35

"2-C3: Clinical Allergy & Immunology : Poster Sessions", Respirology, 2013.

Publication

---

<1 %

36

Al-Mayahie, Sareaa MG. "Phenotypic and genotypic comparison of ESBL production by Vaginal Escherichia coli isolates from pregnant and non-pregnant women", Annals of Clinical Microbiology and Antimicrobials, 2013.

Publication

---

<1 %

37

Edmond SK Ma, KH Kung, Hong Chen.  
"Combating antimicrobial resistance during  
the COVID-19 pandemic", Hong Kong Medical  
Journal, 2021

Publication

---

<1 %

38

Sofia D. Nyberg, Monica Österblad, Antti J.  
Hakanen, Sonja Löfmark, Charlotta Edlund,  
Pentti Huovinen, Jari Jalava. "Long-term  
antimicrobial resistance in Escherichia coli  
from human intestinal microbiota after  
administration of clindamycin", Scandinavian  
Journal of Infectious Diseases, 2009

Publication

---

<1 %

---

Exclude quotes      On

Exclude matches      Off

Exclude bibliography      On