

# Correlation Between Mannitol Intervention And Electrolyte Imbalance In Traumatic Brain Injury Patients Of Regional General Hospital Of West Nusa Tenggara Province

*by Rohadi Rohadi*

---

**Submission date:** 08-May-2023 03:15AM (UTC-0500)

**Submission ID:** 2087359211

**File name:** C16.pdf (397.85K)

**Word count:** 4339

**Character count:** 22743



**CORRELATION BETWEEN MANNITOL INTERVENTION AND ELECTROLYTE  
IMBALANCE IN TRAUMATIC BRAIN INJURY PATIENTS OF REGIONAL  
GENERAL HOSPITAL OF WEST NUSA TENGGARA PROVINCE**

I Gusti Ngruh Wira Narayana<sup>1</sup>, Bambang Priyanto<sup>2,3</sup>, Rohadi Muhammad Rosyidi<sup>2,3</sup>, Maz  
Isa Ansyori<sup>4,5</sup>, Arif Zuhan<sup>6,7</sup>, Ramses Indriawan<sup>8</sup>, Wahyu Nurchalamsah<sup>8</sup>, Akhada Maulana<sup>9</sup>,  
Lalu Shaktisila Fatrahady<sup>10</sup>, Sanditias Putrawan<sup>10</sup>

1 Medical Student of Mataram University

2 Department of Neurosurgery Faculty of Medicine Mataram University

3 Neurosurgeon of Regional General Hospital of West Nusa Tenggara Province

4 Department of Cardiothoracic Surgery West Nusa Tenggara General Provincial Hospital

5 Department of Cardiothoracic Surgery Mataram University

6 Department of Digestive Surgery Mataram University

7 Digestive Surgeon of Regional General Hospital of West Nusa Tenggara Province

8 Oncology Surgeon of Regional General Hospital of West Nusa Tenggara Province

9 Urology Surgeon of Regional General Hospital of West Nusa Tenggara Province

10 General Surgery Residency Training Program, Department Of Surgery, Medical Faculty Of  
Mataram University

**ABSTRACT**

**Background:** Traumatic brain injury (TBI) is a commonly occurring injury following traffic accident. The most common complication following TBI is electrolyte imbalance and might be happen due to mannitol use. This research aims to know the correlation between mannitol intervention and electrolyte imbalance (natrium, kalium and chloride) in TBI patients of Regional General Hospital of West Nusa Tenggara Province.

**Methods:** This is a retrospective research with cross-sectional design. This research was done in Regional General Hospital of West Nusa Tenggara Province. Samples were collected by medical records of TBI patients from 2018 to 2019 which had met the inclusion criteria and then tested with correlation comparative test.

**Results:** From 160 samples that were obtained, 146 samples had electrolyte data, then categorized into those who were given mannitol (73,29%) and those who were not given mannitol (26,71%). The most common electrolyte imbalance was chloride (70,55%) which in hyperchloremia condition, followed by kalium (30,82%) with hyperkalemia and natrium (28,08%) with hyponatremia. Mannitol is most commonly given for one day and has the mean intervention of 2.6 days. Length of mannitol intervention and electrolyte levels were not



normally distributed. There was no statistically significant difference between mannitol intervention and electrolyte imbalance ( $p=1.000$ ,  $p=1.000$ ,  $p=0.481$ ) and length of stay on patients with no mannitol intervention and electrolyte imbalance ( $p=0,856$ ,  $p=0,303$ ,  $p=0,347$ ).

**Conclusion:** This research shows no correlation between the length of mannitol intervention and electrolyte imbalance (represented by natrium, kalium, and chloride) in TBI patients of Regional General Hospital of West Nusa Tenggara Province.

**Keywords:** mannitol intervention, electrolyte imbalance, traumatic brain injury.

## Introduction

Traumatic brain injury is a brain disorder caused by outer mechanical force such as external impact or hit or penetrating injury.<sup>1,2</sup> The prevalence, morbidity and mortality rate of TBI is moderately high with the most common cause is traffic accident. The most common follow up event after traumatic brain injury is electrolyte disturbances. Various suspected etiologies such as intravenous fluid administration, the use of diuretics such as furosemide and mannitol, Syndrome of Inappropriate ADH Secretion (SIADH), and Cerebral Salt Wasting Syndrome (CSW) are said to cause disturbances in serum electrolytes of sodium, potassium, calcium and phosphate.<sup>3,5-7</sup>

Electrolyte disturbances may worsen due to mannitol administration, where mannitol administration in traumatic brain injury patients aims to prevent and treat intracranial hypertension, stabilization of cerebral perfusion pressure and brain oxygenation.<sup>3,7</sup> The use of mannitol can

cause polyuresis and is a possible source of loss of various electrolytes in patients.<sup>1,3</sup> Excessive reduction in intravascular fluid volume due to mannitol administration can lead to dehydration and electrolyte disturbances in the form of hyponatremia and hypokalemia.<sup>4</sup>

## Materials and Methods

This study was a retrospective unpaired comparative study with a cross sectional design. This study was conducted on traumatic brain injury patients at the Regional General Hospital of West Nusa Tenggara Province. Inclusion criteria were 1) Patients with head trauma who have medical records at Regional General Hospital of West Nusa Tenggara Province, 2) Patients with head trauma and diagnosed with traumatic brain injury, 3) Patients with a diagnosis of traumatic brain injury and accompanied by symptoms of increased intracranial pressure, 4) Patients with traumatic brain injury with increased intracranial pressure and were given



mannitol, and 5) Patients with traumatic brain injuries who had laboratory results of serum electrolytes (sodium, potassium, and chloride). The exclusion criteria were patients with a diagnosis of traumatic brain injury who did not have laboratory data on serum electrolytes and patients with electrolyte balance disorders before administration of mannitol. The minimum sample in this study is 150 samples. This study used a patient's medical record that was taken once. The medical record data recorded were data on the degree of traumatic brain injury measured using Glasgow Coma Scale (GCS), the duration of mannitol administration which was divided into  $\leq 7$  days and  $> 7$  days, the length of patient care in days, sodium, potassium and chloride levels which were divided into normal and impaired (normal sodium levels 135-145 mmol / L, potassium 3,4-5,4 mmol / L, chloride 95-105 mmol / L) and age divided into 18–24 years, 25–34 years, 35–44 years, 45–54 years, 55–65 years, and the gender of the patient.

### Results

In this study, 160 samples were obtained. This sample consisted of 111 men (69.4%) and 49 women (30.6%). The mean age was 38.5 years and the mode was 18

years. The 18–24-year age group had the greatest prevalence (25%). Of the total sample, 57 of them had severe brain injury (35.6%), 66 had moderate brain injury (41.3%) and 37 had mild brain injury (23.1%). The most frequent cause of injury was traffic accidents (87.5%). Leucocytosis occurred in 148 traumatic brain injury patients (92.5%). There were 27 traumatic brain injury patients who experienced comorbidities (16.9%) and 133 patients who did not experience comorbidities (83.1%). The accompanying injuries were mostly fractures. The average length of stay is 11 days. The condition of patients at the time of discharge from the hospital was still alive as much as 85.6%, so that the mortality rate for traumatic brain injury cases in this study was 14.4%.



**Table 1** Sample Characteristics

<b>Characteristics</b>		<b>Frequencies (N)</b>	<b>Percentage (%)</b>
<b>Gender</b>	Male	111	69.4
	Female	49	30.6
<b>Severity of brain injury</b>	Severe brain injury	57	35.6
	Moderate brain injury	66	41.3
	Mild brain injury	37	23.1
<b>Cause of trauma</b>	Traffic accident	140	87.5
	Attack	1	0.6
	Hit	2	1.2
	Fall	10	6.2
	Struck by an object	7	4.4
<b>Leukocytosis</b>	Yes	148	92.5
	No	12	7.5
<b>Accompanying trauma</b>	Yes	27	16.9
	No	133	83.1
<b>Conditions on discharge</b>	Dead	23	14.4
	Alive	137	85.6
<b>Age</b>	18 – 24 years	40	25.0
	25 – 34 years	30	18.8
	35 – 44 years	35	21.9
	45 – 54 years	22	13.8
	55 – 65 years	33	20.6
<b>Age (mean)</b>	38.6 years		
<b>Age (modus)</b>	18 years		
<b>Length of stay (mean)</b>	11 days		

In this study, 146 samples had electrolyte data. Of these 146 samples, 107 (73.29%)


were given mannitol and 39 (26.71%) were not given mannitol. In patients given mannitol, the electrolyte disturbances



mostly occurred in chloride as many as 79 people. Disruption to chloride level was dominated by hyperchloremia (72.9%) followed by hypochloreaemia (0.9%). The second most common problem is sodium, which happened to 33 people. Disorders of sodium consisted of hyponatremia (23.4%) and hypernatremia (7.5%). Imbalance in potassium occurred in 31 people consisting of hypokalemia (9.34%) and hyperkalemia

(19.6%). The average sodium content was 136.89 mmol / L, potassium was 11.73 mmol / L and chloride was 106.74 mmol / L. The highest sodium content was 155 mmol / L and the lowest was 123 mmol / L. The highest potassium content was 50 mmol / L and the lowest was 2.8 mmol / L. The highest chloride content was 134 mmol / L and the lowest was 65 mmol / L.

**Graph 1** Electrolyte Disorders



**Table 2** Variable analysis test results of the length of treatment for patients who were given mannitol, the length of treatment for patients who were not given mannitol, and the length of mannitol administration

	<b>Length of treatment</b>	<b>Length of treatment (without mannitol)</b>	<b>Length of mannitol administration</b>
	P value (Mann-Whitney)	P value (Mann-Whitney)	P value (Fisher)
<b>Natrium level</b>	0.080	0.856	1.000
<b>Kalium level</b>	0.440	0.303	1.000
<b>Chloride level</b>	0.941	0.347	0.481

### Discussion

In this study, it was found that the gender who experienced most traumatic brain injury were men. This is consistent with several previous studies which found that the incidence of traumatic brain injury was higher in men.<sup>8,9</sup> The median age with traumatic brain injury was 38.5 years. This is similar to a study by Marquez de la plata which found an average age of 36 years.<sup>10</sup> Traffic accident was the main cause of traumatic brain injury in this study. The same thing was found in a study by Dunne et al where they found that traffic accident was the most common cause of traumatic brain injury.<sup>11</sup> The incidence of leucocytosis occurred in 92.5% of the samples. This result is different from Guel's study which found 45.6% of traumatic

brain injury patients in his study had leucocytosis.<sup>12</sup> The average length of patient care in days in this study was 11 days. In a previous study by Tardif et al., It was found that the average length of stay was similar, which was 11.7 days.<sup>13</sup>

The mortality rate from traumatic brain injury in this study was 14.4%. Similar results were found in a study by Kourbeti, which found that mortality occurred in 12.8% of the sample.<sup>14</sup> This finding is different from that of Song et al., Which found that 4.7% mortality from traumatic brain injury in 2010–2014.<sup>15</sup> The difference was found in 2013–2017 by a study by Okidi et al which found that the mortality of traumatic brain injury reached 33%. This difference might be due to the





lack of health resources in the African region<sup>16</sup>. Mortality of traumatic brain injury also shows a decreasing trend each year along with the developments in the health sector.<sup>17</sup> Administration of mannitol is often used as a dehydrating agent in cases of traumatic brain injury.<sup>18</sup> The same thing was found in this study where mannitol was used more in cases of traumatic brain injury (107 versus 39). To reduce the increased intracranial pressure (ICP), mannitol is usually given 0.5–2 g / kg intravenously for 30 to 60 minutes where the effect will appear in 5 to 10 minutes and this mannitol is given every six hours.<sup>19</sup> It is similar with this study which found that mannitol was most often given every 4 hours. The effect of mannitol can work for 1.5 to 6 hours on ICP so that if long-term reduction of ICP is needed, mannitol administration can be done every 2 to 6 hours.<sup>20</sup> A study conducted by Tan et al also found that administering mannitol every four hours was effective at reducing ICP during the first four days.<sup>21</sup> Mannitol was most often given for one day. This is consistent with what Tan et al wrote that mannitol should not be given for more than eight days.<sup>21</sup> This is because mannitol can aggravate cerebral oedema. Although mannitol is difficult to cross the blood-brain barrier, it can easily pass-through gaps in the damaged endothelium. Prolonged use of

mannitol can cause mannitol leakage across the blood-brain barrier and exacerbate cerebral oedema. This is because the mannitol that has entered the brain draws water into the brain.<sup>19</sup>

In this study, hyperchloremia, hyponatremia, and hyperkalemia were mostly found in patients with traumatic brain injury. This is in accordance with a study conducted by Lee et al., 2016 which found that hyperchloremia occurred in 24.4% of traumatic brain injury patients. This is due to administration of high chloride fluids such as normal saline which can cause hyperchloremia. This hyperchloremic condition is associated with postoperative complications, acute renal failure and infection.<sup>22</sup> Hyponatremia is common in traumatic brain injury patients. Hyponatremia is the result of a syndrome of abnormal secretion of antidiuretic hormone (Syndrome of Inappropriate Antidiuretic Hormone Secretion / SIADH) and Cerebral Salt Wasting Syndrome (CSWS).<sup>23</sup> Hyponatremia due to SIADH usually occurs after traumatic brain injury and is mild. Chronic hyponatremia due to SIADH can result from damage to the infundibulum or posterior pituitary. This results in inappropriate ADH secretion.<sup>24</sup> CSWS is an electrolyte disorder characterized by hyponatremia and hypovolemia. The





underlying etiology has not been widely studied, but it is hypothesized to be caused by elevated levels of natriuretic peptides and changes in the sympathetic nervous system, renin – angiotensin – aldosterone, and adrenomedullin.<sup>25</sup> Hyperkalemia is the most common potassium electrolyte disturbance found in this study. This is different from the research by Syah et al. who found that hypokalemia was mostly found in traumatic brain injury patients. Hyperkalemia in this study may occur due to catecholamine secretion that accompanies head trauma resulting in stimulation of the Na / K ATPase pump.<sup>26</sup>

In this study, there was no significant difference between length of treatment and electrolyte disturbances, both in patients who were given mannitol or patients who were not given mannitol. Different results were obtained in a study by Ouf et al., 2015 which found that there was a significant relationship between length of treatment and electrolyte disturbances.<sup>27</sup> This difference is due to several confounding factors such as drugs or therapies that affect electrolyte levels such as diuretics and therapy, fluids, as well as dietary changes.<sup>28</sup> In this study, the results of the analysis test showed no significant difference between the duration of mannitol administration and electrolyte disturbances. This is similar to a study conducted by Czupryna in 2018

which examined the effect of mannitol with electrolyte concentrations in encephalitis patients and found that there was no significant difference in electrolyte balance after administration of mannitol.<sup>29</sup> This finding was also similar to the study conducted by Seo in 2010 which found that there was no difference in hypernatremia and hyponatremia after seven days of mannitol administration.<sup>30</sup>

This insignificant result was caused by several confounding factors. Electrolyte disturbances such as hyperchloremia can occur due to normal saline fluid resuscitation. Resuscitation of large volumes of normal saline can cause excess chloride ions in the blood.<sup>31</sup> Hypochloremia in TBI patients can be caused by the body's response by reducing renal perfusion, resulting in decreased Claudin-2 expression. Decreasing Claudin-2 expression will make it more difficult for chlorine to be absorbed.<sup>32</sup> Another confounding factor is an abnormal body response to trauma. An example is SIADH which occurs in response to trauma and causes hyponatremia. In addition, CSWS can also cause hyponatremia.<sup>23</sup> Hypokalemia occurs in patients with traumatic brain injury due to movement of potassium into cells in response to increased epinephrine after a traumatic event. This is evidenced by a research from



Beal et al who found high epinephrine levels in new traumatic brain injury patients.<sup>33,34</sup> The weakness of this study is the insufficient medical record data. Electrolyte data were only taken for a short period of time, while the data on mannitol administration were quite complete. This caused the researchers to be unable to analyse the exact length of mannitol administration. Another weakness is the incompleteness of medical records to record patient comorbidities such as diabetes mellitus and other diseases that can cause electrolyte disturbances. This study also did not exclude normal saline administration so that electrolyte disturbances can be caused by improper administration of normal saline.

### Conclusion

Based on the results of this study, it can be concluded that there is no relationship between the use of mannitol and the condition of serum electrolyte imbalance in Traumatic Brain Injury patients at the Regional General Hospital of West Nusa Tenggara Province.

### References

1. Jain A, Gauda B, Gajjar R, Gupta PB. Correlation Between the Severity of Head Injury and Electrolytes in Patients with Traumatic Brain Injury. *J Dent Med Sci*. 2017;16(9):55–8.
2. Capizzi A, Woo J, Verduzco-Gutierrez M. Traumatic Brain Injury. *Med Clin North Am* [Internet]. 2020 Mar;104(2):213–38. Available from: <https://linkinghub.elsevier.com/retrieve/pii/S0025712519301294>
3. Gupta S, Ahuja J, Sharma A. Electrolytes Imbalance in Traumatic Brain Injury Patients. *Int J Med Sci*. 2013;1(1):2349.
4. Abdul Rachman I, Rahardjo S, Chasnak Saleh S. Terapi Hiperosmolar pada Cedera Otak Traumatika. *J Neuroanestesi Indones* [Internet]. 2015;4(2):119–33. Available from: <http://inasnacc.org/ojs2/index.php/jni/article/view/110/pdf>
5. Jha AK, Kumari M, Kumar S, Tomar P, Prakash A, Kumar A. Serum Electrolyte Imbalances in Head Injury Patients: An Institutional Analysis. *Res Pap Neurosci India*, 2017;6(12):1233–5.
6. Lohani S, Devkota UP. Hyponatremia in Patients with Traumatic Brain Injury: Etiology, Incidence, and Severity Correlation. *World Neurosurg* [Internet]. 2011



- Sep;76(3–4):355–60. Available from:  
<https://linkinghub.elsevier.com/retrieve/pii/S1878875011004323>
7. Kumar N. Evaluation of Serum Electrolytes in Traumatic Brain Injury Patients: Prospective Randomized Observational Study. *J Anesth Crit Care Open Access* [Internet]. 2016 Jul 28;5(3). Available from:  
<https://medcraveonline.com/JACC/OA/evaluation-of-serum-electrolytes-in-traumatic-brain-injury-patients-prospective-randomized-observational-study.html>
  8. Munivenkatappa A, Agrawal A, Shukla DP, Kumaraswamy D, Devi BI. Traumatic brain injury: Does gender influence outcomes? *Int J Crit Illn Inj Sci* [Internet]. 2010;6(2):70–3. Available from:  
<http://www.ncbi.nlm.nih.gov/pubmed/27308254>
  9. Biswas RK, Kabir E, King R. Effect of sex and age on traumatic brain injury: a geographical comparative study. *Arch Public Health* [Internet]. 2017;75:43. Available from:  
<http://www.ncbi.nlm.nih.gov/pubmed/29043082>
  10. Marquez de la Plata CD, Hart T, Hammond FM, Frol AB, Hudak A, Harper CR, O’Neil-Pirozzi TM, Whyte J, Carlile M, Diaz-Arrastia R. Impact of age on long-term recovery from traumatic brain injury. *Arch Phys Med Rehabil*. 2008 May;89(5):896–903.
  11. Dunne J, Quiñones-Ossa GA, Still EG, Suarez MN, González-Soto JA, Vera DS, Rubiano AM. The Epidemiology of Traumatic Brain Injury Due to Traffic Accidents in Latin America: A Narrative Review. *J Neurosci Rural Pract* [Internet]. 2020 Apr;11(2):287–90. Available from:  
<http://www.ncbi.nlm.nih.gov/pubmed/32367985>
  12. Morán Guel E, Tamariz Becerra A, Ruiz Cereceres JI, Cisneros Castolo M. Correlación de la leucocitosis y lesión intracraneal en pacientes con traumatismo craneoencefálico. *Med Crítica*. 2018;32(4):208–16.
  13. Tardif P, Moore L, Boutin A, Dufresne P, Omar M, Bourgeois G, Lessard P, Lionel B, Kuimi B, Turgeon AF. Hospital length of stay following admission for traumatic brain injury in a Canadian integrated trauma system: A



- retrospective multicenter cohort study. *Injury*. 2016;1–7.
14. Kourbeti IS, Vakis AF, Papadakis JA, Karabetos DA, Bertias G, Filippou M, Ioannou A, Neophytou C, Anastasaki M, Samonis G. Infections in traumatic brain injury patients. *Clin Microbiol Infect* [Internet]. 2012 Apr;18(4):359–64. Available from: <https://linkinghub.elsevier.com/retrieve/pii/S1198743X14614476>
15. Song SY, Lee SK, Eom KS, KNTDB Investigators. Analysis of Mortality and Epidemiology in 2617 Cases of Traumatic Brain Injury: Korean Neuro-Trauma Data Bank System 2010–2014. *J Korean Neurosurg Soc* [Internet]. 2016 Sep;59(5):485–91. Available from: <http://www.ncbi.nlm.nih.gov/pubmed/27651867>
16. Okidi R, Ogwang DM, Okello TR, Ezati D, Kyegombe W, Nyeko D, Scolding NJ. Factors affecting mortality after traumatic brain injury in a resource-poor setting. *BJS open* [Internet]. 2020;4(2):320–5. Available from: <http://www.ncbi.nlm.nih.gov/pubmed/32207576>
17. Lu J, Marmarou A, Choi S, Maas A, Murray G, Steyerberg EW, Impact and Abic Study Group. Mortality from traumatic brain injury. *Acta Neurochir Suppl* [Internet]. 2005;95:281–5. Available from: <http://www.ncbi.nlm.nih.gov/pubmed/16463866>
18. Wang K, Sun M, Jiang H, Cao X-P, Zeng J. Mannitol cannot reduce the mortality on acute severe traumatic brain injury (TBI) patients: a meta-analyses and systematic review. *Burn trauma* [Internet]. 2015;3:8. Available from: <http://www.ncbi.nlm.nih.gov/pubmed/27574654>
19. Tenny S, Patel R, Thorell W. Mannitol [Internet]. Treasure Island (FL): StatPearls Publishing; 2020. Available from: <https://www.ncbi.nlm.nih.gov/books/NBK470392/>
20. Rangel-Castilla L, Rangel-Castillo L, Gopinath S, Robertson CS. Management of intracranial hypertension. *Neurol Clin* [Internet]. 2008 May;26(2):521–41, x. Available from: <http://www.ncbi.nlm.nih.gov/pubmed/18514825>
21. Tan G, Zhou J-Y, Mao Y-C, Liu B-W, Liao Q. [Most effective daily dose and use method of mannitol in patients with cerebral hemorrhage



- at acute stage]. *Zhonghua Yi Xue Za Zhi* [Internet]. 2008 Apr 1;88(13):889–91. Available from: <http://www.ncbi.nlm.nih.gov/pubmed/18756953>
22. Lee JY, Hong TH, Lee KW, Jung MJ, Lee JG, Lee SH. Hyperchloremia is associated with 30-day mortality in major trauma patients: a retrospective observational study. *Scand J Trauma Resusc Emerg Med* [Internet]. 2016 Dec 4;24(1):117. Available from: <http://sjtrem.biomedcentral.com/articles/10.1186/s13049-016-0311-7>
23. Sepehri P, Abbasi Z, Mohammadi NS, Bagheri S, Fattahian R. Hyponatremia in traumatic brain injury patients: Syndrome of Inappropriate Antidiuretic Hormone (SIADH) versus Cerebral Salt Wasting Syndrome (CSWS). *J Inj Violence Res* [Internet]. 2012 Nov;4(3 Suppl 1):17. Available from: <https://www.ncbi.nlm.nih.gov/pmc/articles/PMC3571543/>
24. Dick M, Catford SR, Kumareswaran K, Hamblin PS, Topliss DJ. Persistent syndrome of inappropriate antidiuretic hormone secretion following traumatic brain injury. *Endocrinol diabetes Metab case reports* [Internet]. 2015;2015:150070. Available from: <http://www.ncbi.nlm.nih.gov/pubmed/26527077>
25. Leonard J, Garrett RE, Salottolo K, Slone DS, Mains CW, Carrick MM, Bar-Or D. Cerebral salt wasting after traumatic brain injury: a review of the literature. *Scand J Trauma Resusc Emerg Med* [Internet]. 2015 Nov 11;23:98. Available from: <http://www.ncbi.nlm.nih.gov/pubmed/26561391>
26. Rafiq MFA, Ahmed N, Khan AA. Serum Electrolyte Derangements In Patients With Traumatic Brain Injury. *J Ayub Med Coll Abbottabad* [Internet]. 2013;25(1):162–4. Available from: [https://applications.emro.who.int/imemrf/J\\_Ayub\\_Med\\_Coll\\_Abbotabad\\_Pak/J\\_Ayub\\_Med\\_Coll\\_Abbotabad\\_Pak\\_2013\\_25\\_1\\_2\\_162\\_164.pdf](https://applications.emro.who.int/imemrf/J_Ayub_Med_Coll_Abbotabad_Pak/J_Ayub_Med_Coll_Abbotabad_Pak_2013_25_1_2_162_164.pdf)
27. Ouf FM, Aziz AFA, Wahba AH, Kabil AE, Alhusseiny MA. Electrolyte Disturbances and their Impact On Mechanically Ventilated Patients with Acute Exacerbations Of Chronic Obstructive Pulmonary





- Disease. Al-Azhar Assiut Med J. 2015;3(4):27-34.
28. Shrimanker I, Bhattarai S. Electrolytes. Treasure Island (FL): StatPearls Publishing; 2020.
29. Czupryna P, Moniuszko-Malinowska A, Grygorczuk S, Pancewicz S, Dunaj J, Król M, Borawski K, Zajkowska J. Effect of a single dose of mannitol on hydration status and electrolyte concentrations in patients with tick-borne encephalitis. J Int Med Res [Internet]. 2018 Dec;46(12):5083–9. Available from: <http://www.ncbi.nlm.nih.gov/pubmed/30124371>
30. Seo W, Oh H. Alterations in serum osmolality, sodium, and potassium levels after repeated mannitol administration. J Neurosci Nurs [Internet]. 2010 Aug;42(4):201–7. Available from: <http://www.ncbi.nlm.nih.gov/pubmed/20804114>
31. Sharma S, Hashmi MF, Aggarwal S. Hyperchloremic Acidosis [Internet]. Treasure Island (FL): StatPearls Publishing; 2020. Available from: <https://www.ncbi.nlm.nih.gov/books/NBK482340/>
32. Rodríguez-Triviño CY, Torres Castro I, Dueñas Z. Hypochloremia in Patients with Severe Traumatic Brain Injury: A Possible Risk Factor for Increased Mortality. World Neurosurg [Internet]. 2019 Apr;124:e783–8. Available from: <https://linkinghub.elsevier.com/retrieve/pii/S1878875019301196>
33. Beal AL, Deuser WE, Beilman GJ. A role for epinephrine in post-traumatic hypokalemia. Shock [Internet]. 2007 Apr;27(4):358–63. Available from: <http://www.ncbi.nlm.nih.gov/pubmed/17414416>
34. Ookuma T, Miyasho K, Kashitani N, Beika N, Ishibashi N, Yamashita T, Ujike Y. The clinical relevance of plasma potassium abnormalities on admission in trauma patients: a retrospective observational study. J Intensive Care [Internet]. 2015 Dec 13;3(1):37. Available from: <https://jintensivecare.biomedcentral.com/articles/10.1186/s40560-015-0103-6>

# Correlation Between Mannitol Intervention And Electrolyte Imbalance In Traumatic Brain Injury Patients Of Regional General Hospital Of West Nusa Tenggara Province

ORIGINALITY REPORT

**20%**  
SIMILARITY INDEX

**20%**  
INTERNET SOURCES

**3%**  
PUBLICATIONS

**%**  
STUDENT PAPERS

PRIMARY SOURCES

**1** [journal.unram.ac.id](http://journal.unram.ac.id)  
Internet Source

**20%**

Exclude quotes  On  
Exclude bibliography  On

Exclude matches  < 3%



# Correlation Between Mannitol Intervention And Electrolyte Imbalance In Traumatic Brain Injury Patients Of Regional General Hospital Of West Nusa Tenggara Province

---

GRADEMARK REPORT

---

FINAL GRADE

**/0**

GENERAL COMMENTS

**Instructor**

---

PAGE 1

---

PAGE 2

---

PAGE 3

---

PAGE 4

---

PAGE 5

---

PAGE 6

---

PAGE 7

---

PAGE 8

---

PAGE 9

---

PAGE 10

---

PAGE 11

---

PAGE 12

---

PAGE 13

---