

Growth performance of several promising local upland rice (*Oryza sativa* L.) genotypes of Southeast Sulawesi

by Ni Wayan Sri Suliartini

Submission date: 02-Apr-2023 09:04PM (UTC-0500)

Submission ID: 2054014641

File name: ijayanto_2021_IOP_Conf._Ser._Earth_Environ._Sci._807_042080.pdf (653.32K)

Word count: 3286

Character count: 16881

PAPER · OPEN ACCESS

Growth performance of several promising local upland rice (*Oryza sativa* L.) genotypes of Southeast Sulawesi

To cite this article: T Wijayanto *et al* 2021 *IOP Conf. Ser.: Earth Environ. Sci.* **807** 042080

View the [article online](#) for updates and enhancements.

You may also like

- [Growth performance of two superior line of local upland rice \(*Oryza sativa* L.\) from SE Sulawesi on the low light intensity](#)
G R Sadimantara, T Alawyah, N W S Suliartini et al.
- [Agronomic trait performance of some promising lines of local upland rice \(*Oryza sativa* L.\) from SE Sulawesi grown under shading conditions](#)
G R Sadimantara, R Adawiah, S Leomo et al.
- [The potential of Southeast Sulawesi local gogo rice genotypes](#)
T Wijayanto, A A Jaya, Nurleni et al.



The Electrochemical Society
Advancing solid state & electrochemical science & technology

241st ECS Meeting

Vancouver, BC, Canada. May 29 – June 2, 2022



ECS Plenary Lecture featuring
Prof. Jeff Dahn,
Dalhousie University

[Register now!](#)



Growth performance of several promising local upland rice (*Oryza sativa* L.) genotypes of Southeast Sulawesi

T Wijayanto¹, Ramdayani¹, Arditya¹, Asniah², N W S Suliartini¹, N M Rahni¹ and L O Afa¹

¹Department of Agrotechnology, Faculty of Agriculture, University of Halu Oleo, Kendari, South East Sulawesi, 93132, Indonesia

²Department of Plant Protection, Faculty of Agriculture, University of Halu Oleo, Kendari, South East Sulawesi, 93132, Indonesia

E-mail: wijayanto_teguh@yahoo.com

Abstract. Southeast Sulawesi, which is located in the southeastern part of Sulawesi island, is one of the upland rice producers in Indonesia, although the cultivation in the area is still not optimal because the upland rice productivity is still considered low. Southeast Sulawesi has an extensive dry land area and local upland rice genotypes have good growth and yield potential. This study aimed to determine the differences in the growth of several local upland rice genotypes of Southeast Sulawesi which were cultivated in Kendari City. This paper is still focused on the vegetative growth of the genotypes tested. This research was carried out in Experiment Field 2, Faculty of Agriculture, Halu Oleo University, which is located in Kambu District, Kendari City, Southeast Sulawesi. The research was conducted using a randomized block design (RBD) consisting of 10 genotypes, namely Wagamba (V1), Wangkariri (V2), Momea (V3), Waburi-buri (V4), Konkep (V5), Bakala (V6), Wakawonda (V7), Tinange (V8), Loiyo putih (V9) and Bombana (V10). Each treatment was repeated 3 times in order to obtain 30 experimental units. The results showed that potential genotypes that can be developed in Kendari City were Wakawonda genotype based on parameters of the number of tillers, plant height, and number of leaves, Wagamba genotype based on leaf width parameter, and Wangkariri genotype based on the flowering age and number of tiller parameter. Based on the overall performances, especially on the number of tillers (one of the most important parameters), in general Wakawonda is the most recommended genotype for further development.

1. Introduction

Rice (*Oryza sativa* L.) is the most important food crop commodity in Indonesia. Nearly 95% of Indonesia's population uses rice as a staple food. Indonesia's per capita rice consumption is among the highest in the world, including when compared to rice consuming countries in Asia, especially China, Japan and Malaysia [1, 2].

Lowland rice still receives primary attention to meet national food needs, due to the upland rice productivity is still relatively low. However, Indonesia actually has the potential for upland rice development because most areas of Indonesia have dry land. The development of upland rice in dry land which has not been optimally utilized can be a solution to the problem of food security [3].



Content from this work may be used under the terms of the Creative Commons Attribution 3.0 licence. Any further distribution of this work must maintain attribution to the author(s) and the title of the work, journal citation and DOI.

Southeast Sulawesi has dry land potential that can be optimized to support efforts to increase national rice production through the development of upland rice [4, 2].

Upland rice productivity can be improved by using good varieties including better growth performance and high yielding varieties planted in a specific environment. Southeast Sulawesi has many potential local upland rice germplasms, which have been cultivated by local farmers for decades [5]. Potential cultivars originating from regions in Southeast Sulawesi can possibly be cultivated in certain areas including in the Kendari region. However, those upland rice genotypes had not been intensively recorded and evaluated. The main purpose of this research was to determine the differences in the growth of several local Southeast Sulawesi's upland rice genotypes which in the future the most recommended genotype can be cultivated in the Kendari area. In this paper, only vegetative growth data are discussed, since the yield parameters are still under evaluation.

30

2. Materials and methods

2.1. Research materials

36

The research was carried out in Experimental Field 2, Faculty of Agriculture, Halu Oleo University, Kendari, Southeast Sulawesi. The research materials used included seeds from 10 local upland rice genotypes, namely Wagamba, Wangkariri, Waburi-buri, Wakawondu (from North Buton area), Bakala, Tinange, Momea, Loiyo putih (from South Konawe area), Konkep (from the region of Konawe Islands), and Bombana (from the Bombana area).

2.2. Research procedure

Soil analysis was carried out by taking soil samples (a-20 cm in depth) and analyzed in the Soil Science Laboratory of Agriculture Faculty, University of Halu Oleo. The analysis showed that soil pH was 4.36, the nitrogen content of 0.05%, and the P content of 5.02 ppm.

Land preparation began with clearing the land from weeds, wood and litter or plant debris. Subsequently, the soil was tilled 3 times and experimental plots were made. Plots were made with a size of 2 m x 3 m with a distance of 1.5 m between groups, the distance between the plots was 0.5 m. Liming was carried out after the first tillage by sprinkling lime over the cultivated land, and rotating it so that the lime is evenly distributed. Liming is carried out based on the pH of the soil analysis result (pH 4.36), with a dose of 1,280 kg ha⁻¹ or 768 g per plot with the intended pH of 5.0, according to the method of [6].

39

Before sowing the seeds were soaked in water for 24 hours. Media for nurseries in trays containing rice husk and sand in the ratio of 1:1. The nursery was held for 2 weeks until ready for transplanting. Planting was carried out when the seedlings were 18-20 days after sowing which begins with making planting holes with a spacing of 25 cm x 20 cm each. Each planting hole was planted with 1 seed to obtain 25 plants per plot.

Fertilization was carried out using organic and inorganic fertilizers. Organic fertilizer in the form of cow dung, with the dose of 450 kg ha⁻¹, was spread out on the plots as a base fertilizer, two weeks before seedlings sowing, to allow a good decomposition of the fertilizer. The inorganic fertilizers used were Urea (N), SP36 (P) and KCL (K). Fertilization doses were urea (N) 150 kg ha⁻¹ (90 g per plot), SP36 (P) 100 kg ha⁻¹ (60 g per plot) and KCl (K) 100 kg ha⁻¹ (60 g per plot).

2.3. Research design

34

This research was carried out using a randomized block design (RBD) consisting of 10 genotypes, namely: Wagamba (V1), Wangkariri (V2), Momea (V3), Waburi-buri (V4), Konkep (V5), Bakala (V6), Wakawondu (V7), Tinange (V8), Loiyo putih (V9) and Bombana (V10). Each treatment was repeated 3 times in order to obtain 30 experimental units.

29

The observed variables consisted of growth parameters, namely: plant height (cm), number of leaves (strands), leaf length (cm), leaf width (cm), number of tillers (plants), flowering age (DAP).

The data from the observations were analyzed using variance based on a randomized block design (RBD), followed by Duncan's Multiple Range Test (DMRT). Once the yield parameter data become available, further analysis such as Path Analysis by Singh and Chaudhary (1979) might be used to identify causal relationships (direct or indirect) between variables, especially between vegetative variables to yield variables.

3. Results and discussion

3.1. Results

The research result findings on some growth parameters of the ten upland rice genotypes tested are presented in table 1. The growth performances of the ten genotypes are varied. In most cases, three upland rice genotypes namely Wakawondu, Wangkariri, and Wagamba generally show better growth performances as compared to the other genotypes.

Table 1. Averages of plant height (cm), leaf length (cm), leaf width (cm), number of leaves (strands), number of tillers (plants) at the age of 77 days after sowing (DAS), and flowering age of several genotypes of local upland rice from Southeast Sulawesi planted in Kendari region.

Treatments	Plant height (cm) 77 DAS	Leaf length (cm) 77 DAS	Leaf width (cm) 77 DAS	Leaves number (strands) 77 DAS	Tiller number (plant) 77 DAS	Flowering ages (day)
V1 (Wagamba)	107.43 ^a	69.57 ^{ab}	1.73 ^a	72.27 ^b	16.90 ^{bc}	135 ^{ab}
V2 (Wangkariri)	101.20 ^a	62.40 ^{bc}	1.60 ^a	50.47 ^b	31.20 ^{ab}	118.2 ^c
V3 (Momea)	109.97 ^a	69.97 ^{ab}	1.53 ^a	54.60 ^b	12.90 ^c	159.1 ^a
V4 (Waburi-buri)	101.60 ^a	69.10 ^{ab}	1.67 ^a	69.83 ^b	19.23 ^{bc}	143.7 ^{ab}
V5 (Konkep)	113.43 ^a	70.70 ^a	1.60 ^a	81.70 ^b	18.57 ^{bc}	133.5 ^{ab}
V6 (Bakala)	82.20 ^b	53.03 ^d	1.23 ^b	87.77 ^b	26.87 ^{bc}	153.5 ^a
V7 (Wakawondu)	80.33 ^b	56.57 ^{cd}	1.23 ^b	151.40 ^a	45.13 ^a	141.5 ^{ab}
V8 (Tinangge)	103.30 ^a	68.87 ^{ab}	1.50 ^a	80.67 ^b	21.60 ^{bc}	135.3 ^{ab}
V9 (Loiyo Putih)	104.23 ^a	69.20 ^{ab}	1.63 ^a	77.30 ^b	17.80 ^{bc}	136.5 ^{ab}
V10 (Bombana)	105.03 ^a	69.90 ^{ab}	1.60 ^a	52.10 ^b	16.93 ^{bc}	124.7 ^{bc}

Note: The numbers followed by different letters in the same column are significantly different based on the DMRT with a 95% confidence level.

Data from observation and the Duncan's multiple range test (table 1) shows that the lowest average plant height is found in the Wakawondu (V7) and Bakala (V6) genotypes. The longest average leaf length was obtained in the Konkep genotype (V5), but it was not significantly different from the genotypes of Wagamba (V1), Momea (V3), Waburi-buri (V4), Tinangge (V8), Loiyo Putih (V9), Bombana (V10). The lowest average leaf length was found in the Bakala genotype (V6). The highest average leaf width was found in Wagamba (V1), but it was not significantly different from other genotypes, except for the Bakala (V6) and Wakawondu (V7) genotypes, which were the genotypes with the lowest leaf width. While the highest average number of leaves of upland rice was found in the Wakawondu genotype (V7), which was significantly different from the other genotypes, and the highest number of tillers was found in the Wakawondu genotype (V7), but it was not significantly different from the Wangkariri genotype (V2). The lowest average number of tillers was found in Momea (V3).

For flowering age, data from observations and the Duncan's test in Table 1 shows that the longest average flowering age is found in Momea (V3) and Bakala (V6), while the fastest average of age of

flowering is in genotypes Wangkariri (V2) and Bombana (V10), with the average values of 118.2 DAS and 124.7 DAS, respectively.



Figure 1. Plant growth condition in the vegetative phase (left) and towards the generative phase (flowering) (right).

3.2. Discussion

Each local upland rice genotype has an average plant height that varies according to its genetic characteristics. Various plant heights can be classified into high, low, and medium groups. Based on the IRRI Standard Evaluation System for rice [7], the plant height classes are short (<90 cm), medium (91-125 cm) and tall (> 126 cm). In this study, the genotypes of upland rice with the medium plant height category (91-125 cm) were the Konkep, Momea, Waburi-Buri, Wangkariri, Wagamba, Tinangge, Loiyo Putih and Bombana genotypes, while the genotypes with low plant height category (<90 cm) were the Bakala and Wakawondu genotypes. Leaf length characters also vary, which was not much different from the plant height. Generally, the higher the upland rice plant, the longer the leaves will be because the length of the upland rice leaves follows the height of the stem.

The number of tillers is a product of the vegetative phase of the plant which determines the yield of rice crops. According to [8], rice tillers are an indicator of the growth of healthy or sick rice plants. Although in character, the number of tillers is also influenced by plant genetics, the role of N is also very important in the process of producing the number of tillers. The results showed that the highest average number of tillers at the age of 77 DAS was found in the Wakawondu genotype, namely 45.1 tillers (table 1). For the flowering age variable, the genotypes with the fast-flowering time were Wangkariri genotype with an average of 118 DAS and Bombana with an average of 124 DAS. While the genotypes with a long flowering time were the genotypes of Momea and Bakala with averages flowering age of about 159 DAS and 153 DAS, respectively.

The new type of rice is characterized by the character of fewer tillers (8-10 tillers), all productive tillers, thick-skinned (200-250 grains per panicle), pithy, medium plant height (90-100 cm), and sturdy trunk, upright leaves, thick, and dark green, deep and deep roots, medium age (110-130 days), and resistant to pests and diseases [9]. These characteristics are intended to increase the yield potential by 20-25% [10]. [11] reported that local upland rice grown on marginal soils in North Buton, including the Wakawondu genotype, had an average panicle length of 37.32 cm, an average number of grains of 149.37 grains, an average wet weight of 53.80 g, an average dry weight of 44.88 g, Wangkariri genotype had an average panicle length of 35.57 cm, an average number of grains 101.35, an average wet weight of 34.20 g and a dry weight of 25.65 g. While the Waburi-buri genotype had an average panicle length of 31.69 cm, an average number of grains 119.90 and an average wet weight of 34.68 g and a dry weight of 28.74 g.

Based on the criteria for new types of rice, in this study, those with medium plant height were Wakawondu and Bakala genotypes. However, based on previous research [12], the Bakala and Wakawondu cultivars had a higher average plant height compared to Wangkariri cultivars [13, 14]. From the results of this study, the genotypes which had the characteristics of ideal type of rice based

on the New Type Rice based on the parameters of the number of tillers, plant height and number of leaves was the Wakawondu genotype, based on flowering age and the number of tillers was in the Wangkariri genotype, and based on leaf width parameter there was on the Wagamba genotype. The character appearance shown is not only influenced by genetic factors [15, 16], it is also greatly influenced by environmental factors on the growth of these plants [17, 18].

4. Conclusion

Based on the results of research and discussion, it can be concluded that there are differences in the characters of the vegetative growth of the local upland rice genotypes planted in Kendari City, especially based on parameters of plant height, leaf length, leaf area, number of leaves, number of tillers and flowering age. In this research, based on the overall performances, especially on the number of tillers (one of the most important parameters), Wakawondu genotype seemed to perform better, in terms of growth parameters, as compared to other genotypes tested. This genotype is potential, and might later be recommended for further evaluation and development. The findings, however, are still yet conclusive until the yield data are analyzed and evaluated.

18

Acknowledgment

The authors gratefully acknowledge the financial support from Directorate General of Higher Education, Ministry of Education and Culture, Indonesia, for funding supports of the research project through the grant for Excellent Higher Education Applied Research Program 2019-2020 & 2021-2022.

References

- [1] Norsalis E 2011 Padi Sawah dan Padi Gogo Tinjauan Secara Morfologi, Budidaya dan Fisiologi *Penelitian Pertanian Universitas Sumatera Utara* **13** 1 19-26
- [2] Badan Pusat Statistik Provinsi Sulawesi Tenggara 2018 *Luas Panen, Produksi, dan Produktivitas Padi Sawah dan Padi Ladang Menurut Kabupaten/Kota di Provinsi Sulawesi Tenggara* (Kendari: Badan Pusat Statistik Provinsi Sulawesi Tenggara)
- [3] Arinta K and Lubis I 2018 Pertumbuhan dan produksi beberapa kultivar padi lokal Kalimantan *Jurnal Agrohorti* **6** (2) 270-280
- [4] Sadimantara G R and Muhidin 2012 Daya hasil beberapa kultivar padi gogo lokal asal Sulawesi Tenggara pada cekama kekeringan *J. Agroteknologi* **2** (3) 121-125
- [5] Wijayanto T, Rembon F S and Susanti Y 2020 Seedling-stage screening method for tolerance of upland rice genotypes to low phosphorus (submitted for publication)
- [6] Krisnawati D and Bowo C 2019 Agricultural lime application to increase rice production in alluvial agricultural soil *Berkala Ilmiah PERTANIAN* **2** (1) 13-18
- [7] IRRI 2002 *Standard Evaluation System for Rice (SES)* (Manila: IRRI Philippines) p 56
- [8] Yoshida S 1981 *Fundamental of Rice Crop Science* (Los Banos, Laguna: The International Rice Research Institute)
- [9] Khush G S 2001 Green revolution: the way forward *Nat Rev Genet* **2** 815-822
- [10] Khan M M A, Haque E, Paul N C, Khaleque M A, Al-Garni S M S, Rahman M and Islam M T 2017 Enhancement of growth and grain yield of rice in nutrient deficient soils by rice probiotic bacteria *Rice Science* **24** (5) 264-273
- [11] Kadidaa B, Sadimantara G R, Suaib, Safuan L O and Muhidin 2017 Pengaruh pupuk organik dalam peningkatan produksi padi gogo lokal pada lahan marginal di Buton Utara Indonesia *Bioteknologi. Res. Asia* **14** (3) 1051-1054
- [12] Wahab A and Sabur A 2014 Karakteristik vegetatif enam kultivar padi gogo lokal Sulawesi Tenggara *Prosiding Seminar Nasional "Inovasi Teknologi Pertanian Spesifik Lokasi"*. Banjarbaru **17** 249-257
- [13] Anggia 2012 Pengaruh pemberian pupuk kandang sapi terhadap pertumbuhan dan produksi tanaman padi gogo lokal kultivar Wakawondu *Jurnal Akademika LPPM Universitas Dayanu Ikhsanuddin* **13** (1) 1-10

- [14] ² Meliala J H S, Basuki N and Seogianto A 2016 Pengaruh iradiasi sinar gamma terhadap perubahan fenotipik tanaman padi gogo (*Oryza sativa* L.) *Jurnal Produksi Tanaman* **4** (7) 585-594
- [15] ⁵ Borromeu M D R, Sadimantara I G R and Pasolon Y B 2018 Respon pertumbuhan dan produksi beberapa kultivar padi gogo lokal Sultra terhadap volume pemberian air *Jurnal Berkala Penelitian Agronomi* **6** (2) 25-36
- [16] ⁴ Nguyen A H, Wijayanto T, Erskine W and Barker S J 2016 Using green fluorescent protein shed light on *Lupinus angustifolius* L. transgenic shoot development *Plant Cell Tiss Organ Cult* **127** (3) 665-674
- [17] Sadimantara G R, Tanti T, Muhidin, Wayan N S S and Wijayanto T 2013 Pendugaan diversitas genetik dan korelasi antar karakter agronomi padi gogo (*Oryza sativa* L.) lokal Sulawesi Tenggara *Agriplus* **23** (3) 1-9
- [18] ¹⁴ Suliartini N W S, Wijayanto T, Madiki A, Muhidin and Tufaila M 2018 Yield potential improvement of upland red rice using gamma irradiation on local upland rice from Southeast Sulawesi Indonesia *Bioscience Research* **15** (3) 1673–1678

Growth performance of several promising local upland rice (*Oryza sativa* L.) genotypes of Southeast Sulawesi

ORIGINALITY REPORT

22%

SIMILARITY INDEX

18%

INTERNET SOURCES

11%

PUBLICATIONS

8%

STUDENT PAPERS

PRIMARY SOURCES

1	jurnal.yudharta.ac.id Internet Source	1%
2	www.jppipa.unram.ac.id Internet Source	1%
3	www.sciencegate.app Internet Source	1%
4	www.arccjournals.com Internet Source	1%
5	www.jim.unsyiah.ac.id Internet Source	1%
6	epubs.icar.org.in Internet Source	1%
7	sinta.ristekbrin.go.id Internet Source	1%
8	Achmad Gunawan, Nur Rosida, Khaerana Khaerana, Arif Muazam, Ani Mugiasih. "Evaluation of tungro-resistant lines based on tungro disease severity and agronomic	1%

characteristics", Jurnal Hama dan Penyakit
Tumbuhan Tropika, 2022

Publication

9	Submitted to Korea National University of Transportation Student Paper	1 %
10	www.ejournal.unmus.ac.id Internet Source	1 %
11	Yossita Fiana, Muhamad Hidayanto. "Adaptation test for shallots varieties in Kutai Kartanegara Regency, East Kalimantan Province", AIP Publishing, 2023 Publication	1 %
12	etd.repository.ugm.ac.id Internet Source	1 %
13	walennae.kemdikbud.go.id Internet Source	1 %
14	Submitted to Universitas Mulawarman Student Paper	1 %
15	(5-9-14) http://202.141.12.149/download/Annual%20Report%2007.pdf Internet Source	1 %
16	conference.unsri.ac.id Internet Source	1 %
17	repositori.umsu.ac.id	

	Internet Source	1 %
18	www.textroad.com Internet Source	1 %
19	pdfs.semanticscholar.org Internet Source	<1 %
20	popups.uliege.be Internet Source	<1 %
21	www.ricescience.org Internet Source	<1 %
22	jurnal.univpgri-palembang.ac.id Internet Source	<1 %
23	Prasajith Kapila Dissanayake, Wekumbure Gedara Chandima Wekumbura, Alge Watage Wijeratne, Douglas Siril Abeywickrama Wijesundara et al. "Morphological characterization, antioxidant capacity and diversity of Syzygium cumini trees from Sri Lanka", Horticultural Plant Journal, 2021 Publication	<1 %
24	academic-accelerator.com Internet Source	<1 %
25	helix.dnares.in Internet Source	<1 %
26	Submitted to Curtin University of Technology Student Paper	

<1 %

27

login.easychair.org

Internet Source

<1 %

28

hss.ulb.uni-bonn.de

Internet Source

<1 %

29

ijaast.com

Internet Source

<1 %

30

dspace.puet.edu.ua

Internet Source

<1 %

31

jpi.faterna.unand.ac.id

Internet Source

<1 %

32

www.chemijournal.com

Internet Source

<1 %

33

technodocbox.com

Internet Source

<1 %

34

A. Misra, N. K. Srivastava. "Effect of the triacontanol formulation ?Miraculan? on photosynthesis, growth, nutrient uptake, and essential oil yield of Lemongrass (Cymbopogon flexuosus) Steud. Watts", Plant Growth Regulation, 1991

Publication

<1 %

35

Marsela Anggita Ratri, Yuliawati Yuliawati. "Perbandingan Pendapatan Usahatani Padi

<1 %

Varietas Ciherang dan Mekongga di Desa Rogomulyo", Agritech: Jurnal Fakultas Pertanian Universitas Muhammadiyah Purwokerto, 2019

Publication

36

Ni Wayan Sri Suliartini, Kuswanto Kuswanto, Nur Basuki, Andy Soegianto. "Superior Lines Candidates Evaluation of Two Local Red Rice Southeast Sulawesi cultivars (Indonesia) Derived from Gamma rays Irradiation Techniques", International Journal of Plant Biology, 2017

Publication

<1 %

37

hyper.ahajournals.org

Internet Source

<1 %

38

www.banglajol.info

Internet Source

<1 %

39

D. K. JHA. "Mineral nutrition in the tripartite interaction between Frankia, Glomus and Alnus at different soil phosphorus regimes", New Phytologist, 2/1993

Publication

<1 %

40

M. Subandi, Alfi Muntafi I., Budy Frasetya T. Q., Sarbini, Aep Kusnawan. "The effect of various nutrient hydroponic formulation on growth and yield of three varieties of common bean (*Phaseolus vulgaris* L.) on hydroponic drip irrigation system", AIP

<1 %

Publishing, 2022

Publication

41	docplayer.info Internet Source	<1 %
42	journals.segce.com Internet Source	<1 %
43	kinfopolitani.com Internet Source	<1 %
44	onlinelibrary.wiley.com Internet Source	<1 %
45	inba.info Internet Source	<1 %

Exclude quotes Off

Exclude matches Off

Exclude bibliography Off