



Serum Homocysteine Level and Ankle-Brachial Index in Peripheral Arterial Disease

Yanna Indrayana[✉], Herpan Syafii Harahap

¹Department of Cardiology and Vascular Disease, Faculty of Medicine, University of Mataram

²Department of Neurology, Faculty of Medicine, University of Mataram

Article Info

Article History:

Submitted December 2019

Accepted February 2020

Published March 2020

Keywords:

Serum homosisteine level, ankle-brachial index, peripheral arterial disease

DOI

<https://doi.org/10.15294/kemas.v15i3.22261>

Abstract

Patients with peripheral arterial disease have a higher risk of death compared to normal populations. There are several relatively new risk factors significantly increase the vulnerability to suffering from peripheral arterial disease, one of which is homocysteine. Studies investigating the role of serum homocysteine level as a biomarker of the severity of peripheral arterial disease based on an ankle-brachial index (ABI) in the different populations were still limited and it was never been studied in Mataram. This was a cross-sectional study aimed to investigate the correlation between serum homocysteine level and ABI in 77 peripheral artery disease (PAD) outpatients in Siti Hajar Hospital, Mataram. The diagnosis of PAD was based on $ABI < 0.9$. Fasting serum homocysteine level was examined using the ELISA technique. Characteristic data collected were gender, age, hypertension, diabetes mellitus, and body mass index (BMI) category. There was a significant correlation between the increase of serum homocysteine levels and the decrease of ABI. The main risk factors for peripheral arterial disease in the subjects are hypertension, diabetes mellitus, and overweight/obesity. Serum homocysteine level is a predictor of peripheral arterial disease severity measured using ABI.

Introduction

At present, cardiovascular diseases (CVDs) are still becoming the main cause of mortality around the world. In 2016, 17.9 million people died due to CVDs, representing about 31% of total death with all causes (World Health Organization, 2017). The prevalence of cardiovascular diseases is predicted to continue to increase, especially in developing countries, since the prevalence of its risk factors is also increasing (Stewart, Manmathan, & Wilkinson, 2017). Basic Health Research (Riskesdas) survey conducted in 2013 showed that the prevalence of cardiovascular diseases in Indonesia, a representation of developing countries, is about 1.5% from the total population (Badan Penelitian dan Pengembangan Kesehatan, 2013). Similar to the number of death globally, about 37% of deaths in Indonesia is attributable to cardiovascular diseases (Cardiovascular

Division and Health Services Research Centre, 2017).

Cardiovascular diseases are a group of diseases consisted of coronary heart disease, cerebrovascular disease, and peripheral arterial disease. Peripheral artery disease (PAD) is the common manifestation of systemic atherosclerosis especially involving arteries other than the coronary artery, most common in lower extremity arteries (Kohlman-Trigoboff, 2019). Globally, its prevalence is high, especially in low-to-middle income countries, estimated at about 28.7% (Criqui & Aboyans, 2015). Since most of the PAD are unfortunately asymptomatic, it is commonly underdiagnosed and undertreated (Olin & Sealove, 2010). Peripheral arterial disease, either symptomatic or asymptomatic, is associated with increased risk of cardiovascular mortality in the cohort study and its early diagnosis and reduction risk

[✉] Correspondence Address:
Faculty of Medicine, Universitas Mataram, Indonesia
Email: yannaindrayana@unram.ac.id

management is important (Sartipy, Sigvant, Lundin, & Wahlberg, 2018). Therefore, the diagnosis of PAD in the primary care setting has important prognostic value (Diehm, et al., 2009). In clinical practice, screening and diagnosis of PAD can be conducted using a simple and non-invasive instrument named *ankle-brachial index* (ABI) (Campia, Gerhard-Herman, Piazza, & Goldhaber, 2019).

The major risk factors of PAD are similar to those of coronary artery disease and cerebrovascular disease, including older age, male, hypertension, diabetes mellitus, overweight/obesity, and cigarette smoking (Bennett, Silverman, Gill, & Lip, 2009). The prevalence of those risk factors is quite high among Indonesian population (Mihardja, Soetrisno, & Soegondo, 2014; Rachmi, Li, & Baur, 2017; Rosjidi, Isro'in, & Wahyuni, 2017; Harahap & Indrayana, 2018; Peltzer & Pengpid, 2018). Optimal management of those risk factors is crucial in the management of PAD and it may reduce overall cardiovascular mortality (Shammas, 2007). In recent years, some relatively new risk factors for PAD have been proposed, including homocysteine (Bennett, Silverman, Gill, & Lip, 2009).

Homocysteine is an intermediate product of the amino acid methionine metabolism produced via the transmethylation pathway. In normal conditions, homocysteine that is formed will be metabolized either via remethylation or transsulfuration pathway. In the remethylation pathway, homocysteine is converted back to methionine either via folate-dependent or folate-independent remethylation pathway. In the transsulfuration pathway, homocysteine is converted to cysteine (Cacciapuoti, 2018). The remethylation pathway through the folate-dependent pathway is dependent on folic acid, B2, and B12 vitamins, while the transsulfuration pathway is dependent on B6 vitamin (Barroso, Handy, & Castro, 2017). Therefore, the total plasma concentration of homocysteine (tHcy) is influenced by blood concentrations of folic acid, B12 and B6 vitamins (Hankey, 2018). In the blood circulation, a level of homocysteine <13 $\mu\text{mol/L}$ is considered normal, while a level $\geq 13 \mu\text{mol/L}$ is considered hyperhomocysteinemia (Cacciapuoti, 2018).

Since homocysteine was proposed

to be associated with the progression of atherosclerosis, now it is considered to be the risk factor of PAD. A meta-analysis of the epidemiologic study showed that the homocysteine level in PAD patients was higher than those healthy subjects (Khandanpour, Loke, Meyer, Jennings, & Armon, 2009). The role of homocysteine in the pathogenesis of PAD can be influenced by ethnic variation. This was the first study investigating the correlation between serum homocysteine level and severity of PAD determined based on ABI in the sub-population of Mataram, West Nusa Tenggara.

Method

This was a cross-sectional study involving 77 asymptomatic peripheral artery disease outpatients visiting Siti Hajar Hospital, Mataram, during the period of May to October 2019. The diagnosis of peripheral artery disease in those patients was based on their ankle-brachial index (ABI) <0.9. The exclusion criteria are the coexistence of some medical conditions, i.e chronic kidney disease, liver disease and hypothyroid based on medical records, and prior history of consumption folic acid, B6 and B12 in the last three months. This study was approved by Komisi Etik Penelitian Kesehatan Universitas Mataram (No. 233/UN18.F7/ETIK/2019).

The characteristics data of the subjects collected in this study were age, gender, hypertension, diabetes mellitus, and cigarette smoking based on their medical records. The data of body mass index (BMI) of each subject was also collected by measuring weight in kilograms (kg) divided by height in square meters (m^2) (Hicks, et al., 2018). The BMI was categorized into three groups, namely underweight ($\text{BMI} < 18.5 \text{ kg/m}^2$), norm weight ($\text{BMI} 18.5\text{-}24.9 \text{ kg/m}^2$), overweight/obese ($\text{BMI} \geq 25 \text{ kg/m}^2$). Since overweight and obesity are risk factors for PAD, they are grouped into one group, the overweight/obesity group respectively (Murata, et al., 2015).

The ABI was measured by comparing the ratio of higher of the systolic blood pressure of the two ankle arteries of the limb and higher of the two systolic blood pressure of the upper arms using handheld vascular Doppler Bestman BF-520T. The $\text{ABI} < 0.9$ was diagnostic for PAD (Rac-Albu, Iliuta, Guberna, & Sinescu, 2014).

The serum samples for the examination of homocysteine levels were obtained from 5cc of overnight (8 to 10 hours) fasting blood samples of the patients. The serum homocysteine level was measured in the Laboratory of Hepatika, Mataram, using FineTest® Human HCY (Homocysteine) ELISA Kit (Kit Product Code HCY161119).

The data of age was presented as mean (95%CI), while data of gender, hypertension, diabetes mellitus, cigarette smoking, and BMI category were presented as frequency. The statistical analysis of the correlation of the serum homocysteine level and ABI was based on the Spearman test. The result of statistical analysis was significant if $p < 0.05$.

Result and Discussion

The incidence of peripheral artery disease (PAD) is mostly determined by the existence of both major and relatively new risk factors. Homocysteine is a representative of relatively new risk factors for PAD which are getting more attention from the researchers in recent years. In the present study, the characteristic data of the subjects representing the major risk factors as well as serum homocysteine level representing

the relatively new risk factor for PAD were obtained (table 1). The results showed that most of the subjects are hypertensive, suffered from diabetes mellitus, overweight/obese, but with a small frequency of male gender and those who had cigarette smoking habits. The higher frequency of hypertension, diabetes mellitus, and overweight/obese among subjects with PAD in the present study was in accordance with the results of previous studies in different populations (Alvim, et al., 2018; Krishnan, Geevar, Mohanan, Venugopal, & Devika, 2018). The mean age of the subjects is 57 years old. The population of subjects aged 40 years and older in Indonesia is at high risk for cardiovascular disease (Maharani, Sujarwoto, Praveen, Delvac, Tampubolon, & Patel, 2019).

The present study showed that most of the subjects are female (79,2%). This result contrast with the existing reference which showed that male gender was the risk factor for PAD (Bennett, Silverman, Gill, & Lip, 2009; Alzamora, et al., 2010; Aggarwal, Loomba, & Arora, 2012). However, another study conducted by Rafie *et al.* showed similar results with the present study in which the female

Table 1. The characteristics of the subjects

Category	Mean (95%CI), unless otherwise stated
Gender, n(%)	
Male	16 (20.8)
Female	61 (79.2)
Age (years)	57.38(54.94 – 59.82)
Cigarette smoking, n(%)	
Yes	3 (3.9)
No	74 (96.1)
Hypertension, n(%)	
Yes	55 (71.4)
No	22 (28.6)
Diabetes mellitus, n(%)	
Yes	43 (55.8)
No	34 (44.2)
BMI category, n(%)	
Underweight (<18.5 kg/m ²)	4 (5.2)
Norm weight (18.5-24.9 kg/m ²)	29 (37.7)
Overweight/obese (≥25 kg/m ²)	44 (57.1)
ABI	0.79 (0.77 – 0.81)
Homocysteine (μmol/L)	4.79 (4.43 – 5.15)

CI=confidence interval; BMI=body mass index; ABI=ankle-brachial index

gender had a higher frequency (Rafie, et al., 2010). These different results may be due to the differences ethnic of the subjects studied. The differences in ethnic studied affect the role of various vascular risk factors in the pathogenesis of PAD (Bennett, Silverman, Gill, & Lip, 2009). The small frequency of subjects who had the cigarette smoking habit is mostly due to the small frequency of male subjects.

All of the well-known risk factors mentioned above as well as hyperhomocysteinemia contribute to the abnormal ankle-brachial index (ABI) (Campia, Gerhard-Herman, Piazza, & Goldhaber, 2019). The present study showed that the mean of the serum homocysteine level was in the normal range. It is interesting that there was no subject which was PAD patients that showed an elevation in serum homocysteine level (table 1). In the context of the existence of frequency of subjects with PAD who had mild-moderate to severe hyperhomocysteinemia, this result is contradictive with the results of a meta-analysis and previous studies conducted in different populations (Khandanpour, Loke, Meyer, Jennings, & Armon, 2009; Weragoda, Seneviratne, Weerasinghe, & Wijeyaratne, 2016; Preethi & Hemachandran, 2019). It can be proposed that diverse ethnic groups may have different serum homocysteine profiles. However, statistical analysis showed that there was a significant correlation between the increase of serum homocysteine level and the decrease of the ankle-brachial index (ABI) (table 2). It suggested that even in the normal range, the increase of serum homocysteine level is correlated with the progression of peripheral artery disease, but it still needs further investigation.

Table 2. The correlation between serum homocysteine level and ABI

	Mean (95%CI)	r	p-value
Homocysteine	4.79	-0.323	0.004*
ABI	0.79		

*Significant ($p < 0.05$)

The mechanism by which homocysteine is responsible for the progression of peripheral arterial disease is not well established yet.

Theoretically, its role is mediated via its effect on the progression of atherosclerotic plaque of the arterial wall and the generation of hypertension by several mechanisms. First, homocysteine may induce oxidative stress via activation of NADPH oxidase resulting in the generation of reactive oxygen species (ROS) (Omae, Nagaoka, Tanano, & Yoshida, 2013). This ROS is responsible for the decrease of bioavailability of nitric oxide (NO), a molecule important for vasodilatation, and this will lead to vasoconstriction and hypertension (Lai & Kan, 2015). In addition, NO which reacts to superoxide (O_2^-), a kind of ROS, to form peroxynitrite ($ONOO^-$) will facilitate further atherosclerotic plaque progression and decrease of vascular compliance resulting in arterial hypertension (Mury, Chirico, Mura, Millon, Canet-Soulas, & Pialoux, 2018). Second, the homocysteine-induced oxidative stress is also responsible for the activation of matrix metalloproteinases (MMPs), zinc-containing endopeptidases secreted by connective tissue as well as proinflammatory cells. In the atherosclerotic plaque, MMPs degrade elastin matrix separating the intima from the media and induce collagen synthesis. These process will result in smooth-muscle cells migration from the media to the intima, the increase of intima-media thickness, and the decrease of elastin/collagen ratio which eventually induce vascular hypertrophy and endothelial dysfunction, the hallmark of hypertension (Vacek, Rehman, Neamtu, Yu, & Givimani, 2015). Third, homocysteine alters the transsulfuration pathway, a kind of its metabolism pathway, resulting in a decrease of hydrogen sulfide (H_2S) production (Yang & He, 2019). Hydrogen sulfide (H_2S) is well known as a vasorelaxant molecule produced by vascular tissue and a decrease of its production will lead to vascular constriction and hypertension (Sen, Mishra, Tyagi, & Tyagi, 2010). However, further study is still needed to investigate whether the theories described above have a significant negative correlation with ABI, as shown by the result of the present study.

Conclusion

The serum homocysteine level negatively correlates to the ankle-brachial index (ABI). The identifiable risk factors of peripheral

arterial disease (PAD) were age, female gender, hypertension, diabetes mellitus, overweight/obesity, and serum homocysteine level, but not cigarette smoking. The role of ethnicity in influencing serum homocysteine level still needs further investigation.

References

- Aggarwal, S., Loomba, R. S., & Arora, R. (2012). Preventive aspects in peripheral artery disease. *Therapeutic Advances in Cardiovascular Disease*, 6 (2), 53-70.
- Alvim, R. O., Dias, F. A., de Oliveira, C. M., Horimoto, A. R., Ulbrich, A. Z., Krieger, J. E., et al. (2018). Prevalence of Peripheral Artery Disease and Associated Risk Factors in a Brazilian Rural Population: The Baependi Heart Study. *International Journal of Cardiovascular Sciences*, 31 (4), 405-413.
- Alzamora, M. T., Forés, R., Baena-díez, J. M., Pera, G., Toran, P., Sorribes, M., et al. (2010). The Peripheral Arterial disease study (PERART/ARTPER): prevalence and risk factors in the general population. *BMC Public Health*, 10, 38.
- Badan Penelitian dan Pengembangan Kesehatan. (2013). *Riset KESEHATAN DASAR 2013*. Jakarta: Badan Penelitian dan Pengembangan Kesehatan Kementerian Kesehatan RI.
- Barroso, M., Handy, D. E., & Castro, R. (2017). The Link Between Hyperhomocysteinemia and Hypomethylation: Implications for Cardiovascular Disease. *Journal of Inborn Errors of Metabolism & Screening*, 5, 1-15.
- Bennett, P. C., Silverman, S., Gill, P. S., & Lip, G. Y. (2009). Ethnicity and peripheral artery disease. *Q J Med*, 102, 3-16.
- Cacciapuoti, F. (2018). Poor re-Methylation of Homocysteine and Trans-Methylation of Methionine: Cause and Effect of Hyper-Homocysteinemia: Which Role for Folic Acid and Vitamins B-6-12 Supplementation? *Annals of Clinical & Experimental Motabolism*, 3 (1), 1026.
- Campia, U., Gerhard-Herman, M., Piazza, G., & Goldhaber, S. Z. (2019). Peripheral Artery Disease : Past, Present, and Future. *The American Journal of Medicine*, 132, 1133-1141.
- Cardiovascular Division and Health Services Research Centre. (2017). *Reducing the burden of Cardiovascular Disease in Indonesia*. Newton: George Institute for Global Health.
- Criqui, M. H., & Aboyans, V. (2015). Epidemiology of Peripheral Artery Disease. *Circulation Research*, 116, 1509-1526.
- Diehm, C., Allenberg, J. R., Pittrow, D., Mahn, M., Tepohl, G., Haberl, R. L., et al. (2009). Mortality and Vascular Morbidity in Older Adults With Asymptomatic Versus Symptomatic Peripheral Artery Disease. *Circulation*, 120, 2053-2061.
- Hankey, G. J. (2018). B vitamins for stroke prevention. *Stroke and Vascular Neurology*, 3, e000156.
- Harahap, H. S., & Indrayana, Y. (2018). Obstructive Sleep Apnea Risk Level Affect Executive Function Rather than Attention. *KEMAS*, 14 (2), 272-278.
- Hicks, C. W., Yang, C., Ndumele, C. E., Folsom, A. R., Heiss, G., Black III, J. H., et al. (2018). Associations of Obesity With Incident Hospitalization Related to Peripheral Artery Disease and Critical Limb Ischemia in the ARIC Study. *Journal of the American Heart Association*, 7, e008644.
- Khandanpour, N., Loke, Y. K., Meyer, F. J., Jennings, B., & Armon, M. P. (2009). Homocysteine and Peripheral Arterial Disease : Systematic Review and Meta-analysis. *European Journal of Vascular & Endovascular Surgery*, 38 (3), 316-322.
- Kohlman-Trigoboff, D. (2019). Update: Diagnosis and Management of Peripheral Arterial Disease. *The Journal for Nurse Practitioners*, 15 (1), 87-95.
- Krishnan, M. N., Geevar, Z., Mohanan, P. P., Venugopal, K., & Devika, S. (2018). Prevalence of peripheral artery disease and risk factors in the elderly: A community based cross-sectional study from northern Kerala, India. *Indian Heart Journal*, 70 (6), 808-815.
- Lai, W. K., & Kan, M. Y. (2015). Homocysteine-Induced Endothelial Dysfunction. *Annals of Nutrition and Metabolism*, 67, 1-12.
- Maharani, A., Sujarwoto, Praveen, D., Delvac, O., Tampubolon, G., & Patel, A. (2019). Cardiovascular disease risk factor prevalence and estimated 10-year cardiovascular risk scores in Indonesia: The SMARThealth Extend study. *PloS ONE*, 14 (4), e0215219.
- Mihardja, L., Soetrisno, U., & Soegondo, S. (2014). Prevalence and clinical profile of diabetes mellitus in productive aged urban Indonesians. *Journal of Diabetes Investigation*, 5, 507-512.
- Murata, N., Soga, Y., Iida, O., Yamauchi, Y., Hirano, K., Kawasaki, D., et al. (2015). Complex Relationship of Body Mass Index with Mortality in Patients with Critical Limb Ischemia Undergoing Endovascular

- Treatment. *European Journal of Vascular & Endovascular Surgery*, 49 (3), 297-305.
- Mury, P., Chirico, E. N., Mura, M., Millon, A., Canet-Soulas, E., & Pialoux, V. (2018). Oxidative Stress and Inflammation, Key Targets of Atherosclerotic Plaque Progression and Vulnerability: Potential Impact of Physical Activity. *Sports Medicine*, 48 (12), 2725-2741.
- Olin, J., & Sealove, B. (2010). Peripheral Artery Disease: Current Insight Into the Disease and Its Diagnosis and Management. *Mayo Clinic Proceedings*, 85 (7), 678-692.
- Omae, T., Nagaoka, T., Tanano, I., & Yoshida, A. (2013). Homocysteine Inhibition of Endothelium-Dependent Nitric Oxide-Mediated Dilation of Porcine Retinal Arterioles via Enhanced Superoxide Production. *Investigative Ophthalmology & Visual Science*, 54 (3), 2288-2295.
- Peltzer, K., & Pengpid, S. (2018). The Prevalence and Social Determinants of Hypertension among Adults in Indonesia: A Cross-Sectional Population-Based National Survey. *International Journal of Hypertension*, 2018, 5610725.
- Preethi, S. P., & Hemachandran, T. (2019). Serum homocysteine as a predictor of severity of peripheral arterial disease in association with Doppler ultrasonography. *International Surgery Journal*, 6 (11), 3977-3981.
- Rac-Albu, M., Iliuta, L., Guberna, S. M., & Sinescu, C. (2014). The Role of Ankle-Brachial Index for Predicting Peripheral Arterial Disease. *MAEDICA-a Journal of Clinical Medicine*, 9 (3), 295-302.
- Rachmi, C. N., Li, M., & Baur, A. (2017). Overweight and obesity in Indonesia: prevalence and risk factors - a literature review. *Public Health*, 147, 20-29.
- Rafie, A. H., Stefanick, M. L., Sims, S. T., Phan, T., Higgins, M., Gabriel, A., et al. (2010). Sex differences in the prevalence of peripheral artery disease in patients undergoing coronary catheterization. *Vascular Medicine*, 15 (6), 443-450.
- Rosjidi, C. H., Isro'in, L., & Wahyuni, N. S. (2017). Difference in Risk Factor of Cardiovascular Disease Risk on Rural and Urban Population. *KEMAS*, 13 (1), 69-76.
- Sartipy, F., Sigvant, B., Lundin, F., & Wahlberg, E. (2018). Ten Year Mortality in Different Peripheral Arterial Disease Stages : A Population Based Observational Study on Outcome. *European Journal of Vascular & Endovascular Surgery*, 55 (4), 529-536.
- Sen, U., Mishra, P. K., Tyagi, N., & Tyagi, S. C. (2010). Homocysteine to Hydrogen Sulfide or Hypertension. *Cell Biochemistry and Biophysics*, 57 (2-3), 49-58.
- Shammas, N. W. (2007). Epidemiology, classification, and modifiable risk factors of peripheral arterial disease. *Vascular Health and Risk Management*, 3 (2), 229-234.
- Stewart, J., Manmathan, G., & Wilkinson, P. (2017). Primary prevention of cardiovascular disease: A review of contemporary guidance and literature. *Journal of the Royal Society of Medicine Cardiovascular Disease*, 6, 1-9.
- Vacek, T. P., Rehman, S., Neamtu, D., Yu, S., & Givimani, S. T. (2015). Matrix metalloproteinases in atherosclerosis: role of nitric oxide, hydrogen sulfide, homocysteine, and polymorphisms. *Vascular Health and Risk Management*, 11, 173-183.
- Weragoda, J., Seneviratne, R., Weerasinghe, M. C., & Wijeyaratne, S. M. (2016). Risk factors of peripheral arterial disease: a case control study in Sri Lanka. *BMC Research Notes*, 9, 508.
- World Health Organization. (2017, May 17). *Cardiovascular Diseases (CVDs)*. Retrieved January 15, 2020, from World Health Organization: [https://www.who.int/news-room/fact-sheets/detail/cardiovascular-diseases-\(cvds\)](https://www.who.int/news-room/fact-sheets/detail/cardiovascular-diseases-(cvds))
- Yang, Q., & He, G. W. (2019). Imbalance of Homocysteine and H₂S: Significance, Mechanisms, and Therapeutic Promise in Vascular Injury. *Oxidative Medicine and Cellular Longevity*, 2019, 7629673.