

Effect of Mutations Induction on Vegetative and Generative Characters of G16 Rice (*Oryza sativa* L.)

by Sri Suliartini Ni Wayan

Submission date: 02-Jun-2023 08:52PM (UTC-0500)

Submission ID: 2107818094

File name: 0531_JPPIPA_Effect_of_mutations_induction_sri_suliartini_dkk.pdf (623.03K)

Word count: 7656

Character count: 37334



Effect of Mutations Induction on Vegetative and Generative Characters of G16 Rice (*Oryza sativa* L.)

Ni Wayan Sri Suliartini^{1*}, Shinta Adekayanti¹, Anak Agung Ketut Sudharmawan¹, I Gusti Putu Muliarta Aryana¹

52

¹ Agroecotechnology Study Program, Faculty of Agriculture, University of Mataram, Mataram, Indonesia.

Received: March 28, 2023

Revised: May 18, 2023

Accepted: May 26, 2023

Published: May 31, 2023

Corresponding Author:

Ni Wayan Sri Suliartini

sri.suliartini@gmail.com

DOI: [10.29303/jppipa.v9i5.3650](https://doi.org/10.29303/jppipa.v9i5.3650)

51

© 2023 The Authors. This open access article is distributed under a (CC-BY License)



63

Abstract: The aim of this research was to identify the vegetative and generative characters of G16 rice mutants due to gamma-ray irradiation. The experiment was conducted in Saribaye Village, Lingsar District, West Lombok Regency. The experimental design used was an Augmented Design, using three comparison plants, namely the G16 line (parent), Inpago Unram 1 variety, and IPB 3S variety. The 18 mutant populations tested were mutant doses of 200, 300, 400 and 500 Gy. The results showed that the induction of gamma ray mutations affected the character of plant height, flag leaf angle, flowering age, panicle length, number of filled grains per panicle, harvest age, and grain weight per clump. Comparison plants and mutant plants showed an interaction on all observed characters, except the number of total tillers and the number of productive tillers. Wide genetic diversity was shown by all characters, except flag leaf angle, total tiller number, weight of 100 grains, and weight of empty grain per clump. High heritability was obtained on the character of plant height (0.97), flag leaf angle (0.74), flowering age (0.99), number of productive tillers (0.57), panicle length (0.86), number of filled grain per panicle (0.92), number of empty grains per panicle (0.55), age at harvest (1.00), and weight of 100 grains (0.99).

39

Keywords: Gamma ray irradiation; Genetic diversity; Heritability; Mutant rice

Introduction

Rice (*Oryza sativa* L.) is staple food of more than half of the world's population (Ningrat et al., 2021; Tang et al., 2022) and a food crop native to Asia and West Africa. Rice is a very important agricultural commodity, especially in the East Asian region. Ninety percent of the world's rice production is produced by countries in Asia, one of which is Indonesia. According to BPS Statistics Indonesia (BPS.go.id, 2022), rice production in 2021, which is 54.42 million tons of milled dry grain, has decreased by 233.91 thousand tons or 0.43 percent compared to rice production in 2020 which amounted to 54.65 million tons of milled dry grain. This condition exacerbated by a decrease in land area and the conversion of agricultural land to non-agriculture (Ayun et al., 2020).

In order to maintain national rice production, one of the appropriate ways to use is the use of high-yielding varieties of rice. The development of high-yielding

varieties of rice has been emphasized more on improving local rice since 1970, especially in terms of shortening the life of plants so that in one year two to three harvests can be carried out.

In Indonesia, especially in the West Nusa Tenggara region, there are quite a lot of genes for local rice cultivars, one of which is the red rice cultivar (Arinta et al., 2018), is anthocyanin-rich cultivars (Dwiatmini et al., 2018; Suliartini et al., 2020). Red rice can prevent various types of diseases, including cancer, cholesterol, heart disease, constipation and high blood pressure (Castañeda-Ovando et al., 2019). The development of red rice varieties is an important breakthrough to maintain public health. This can certainly provide new hope for breeding activities to get a strain of hope before being released into a high-yielding variety. One of the strains produced from breeding activities is red rice G16 (Umam et al., 2018).

G16 line is red rice which is rich in anthocyanins. This line has a low number of filled grains per panicle

How to Cite:

Suliartini, N.W.S., Adekayanti, S., Sudharmawan, A.A.K., & Aryana, I.G.P.M. (2023). Effect of Mutations Induction on Vegetative and Generative Characters of G16 Rice (*Oryza sativa* L.). *Jurnal Penelitian Pendidikan IPA*, 9(5), 3790-3799. <https://doi.org/10.29303/jppipa.v9i5.3650>

(about 150 grains). Number of filled grains per panicle has a correlation with yield (Suliantini et al., 2021). A low number of filled grains will cause a low production potential.

In order to improve weaknesses, increase genetic diversity, and increase the economic value of G16 red rice, breeding activities have been carried out, namely induction of mutations and producing a basic population in the form of M1. M1 plants have high diversity, but segregation occurs in M2 populations so selection can be made in that generation. Mutations are changes in genetic material that generally occur suddenly and are inherited. Mutations cause phenotype changes in both vegetative and generative character. According to Riviello-Flores et al. (2022), plant genetic mutations can be carried out, one of which is by induction of gamma ray radiation. The usual plant parts in the radiation are seeds or other plant parts that can be g

The results of the study of Suhesti et al. (2021) showed phenotypic changes in gamma-ray-irradiated 77 arcana mutant. Diversity in germination percentage, plant height, number of flowers, and number of fruits in tomato (*Solanum lycopersicum* L.) obtained at irradiation doses 150-450 Gy (Zafar et al., 2022). Genetic diversity obtained through gamma ray irradiation (Parlaongan et al., 2022) is very important for further breeding programs. Basic population genetic diversity is needed for selection activities or combining good characters to produce superior varieties. This is supported by Sudharmawan et al. (Sudharmawan et al., 2022), that genetic diversity in vegetative and generative characters is very important for producing new superior varieties. Selection is carried out on yield characteristics and yield support to obtain superior varieties with high production potential. The higher the genetic diversity, the greater the chance of obtaining new superior varieties with the desired characters. Suliantini et al. (2018) added that broad genetic diversity will 16 fect the success of selection. Based on this statement, this study has been carried out to determine changes in the character of vegetation f and generative plants as well as heritability in rice mutants due to gamma ray irradiation.

Method

This research used an experimental method in the field, in Saribaye Village, Lingsar District, West Lombok Regency, from February 3 to June 14, 2020. The materials used are, among others: G16 line rice seeds , Inpago Unram-1 variety rice seeds, IPB 3S variety rice seeds, vermicompost, atonic growth regulator, cruiser 350 FS, Gandasil D 5 g/5 liters, insecticide Dumil 40 SP for walang sangit and grasshopper pests, fungicide Nativo 75 WG 150 g/ha (6.32 g), Petroganic fertilizer 1000 kg/ha

(42.1 kg), urea (45% N) at a dose of 300 kg/ha (12.6 kg on a land area of 20 m x 21 m), and NPK Phonska 300 kg/ha (12.6 kg), plastic pouches, sacks, plastic, label paper, rafia rope, small mine ropes, markers, and nameplates. The tools used are, among others: tractors, nets, sprayers, sickles, bamboo, tubs, hoes, analytical scales, cellphone cameras, and writing stationery.




Figure 1. Flowchart of research implementation of effect of mutations induction on vegetative and generative characters of G16 rice (*Oryza sativa* L.)

The experimental design used is augmented design. The genotype (g) tested was repeated once (without repeat) and the tester (c) was repeated 3 times.

The flow of research implementation in the field can be seen in Figure 1. Data were analyzed using Augmented Design Anova (Syahril, 2018). Further tests are carried out if there are noticeable differences between treatments using the Duncan Multiple Test (DMRT). The heritability value of the meaning of breadth is calculated according to Allard (1960) and the criteria of the heritability value of the meaning of broad is determined according to Stanfield (1991). Genetic

diversity and phenotypic diversity are obtained from the values in the calculation of heritability.

Result and Discussion

Vegetative and Generative Characters

Research result shows that the higher the dose of irradiation used, the lower the percentage of plants that live (Table 1). Mutant plant (seedling age 2 weeks after planting/wap), with a dose treatment of 200 Gy had a larger percentage of plants growing (89.8%) compared to other dose treatments. Low-dose irradiation treatment caused phenotypic changes in plants but it did not have a lethal effect. This was confirmed by Wahyuni et al. (2022) that certain doses are needed to produce effective mutations. Low dose irradiation treatment but capable of causing large genetic effects will be more effective than high dose irradiation treatment but causing a lot of damage. Gamma irradiation research on Begonia by Wahyuni et al. (2019) also showed irradiation treatment at low doses resulting in more than 80% germination of seeds but phenotypic observations showed variations in stem diameter, plant height, plant width, leaf width, leaf length, petiole diameter, petiole length, and number of leaves. The 500 Gy dose treatment has the lowest percentage of growing plants, which is 24.2% with the number of seedlings growing as many as 121 seedlings. These results are in accordance with the

opinion of Wulandari et al. (2019) that the number of abnormal or dead plants will be higher in line with the higher the dose of irradiation given.

Table 1. Percentage of Living Rice Mutants at Various Gamma-ray Irradiation Doses

Treatment	Percentage of Live Plants	Information
Dose of 200 Gy	89.81%	Grow
Dose of 300 Gy	81.00%	Grow
Dose of 400 Gy	45.22%	Grow
Dose of 500 Gy	24.23%	Grow

Description: Dose of 500Gy: only 11 growing plants remain

All treatments, both comparison plants (checks) and mutant plants (genotypes) were transplanted at the age of 2 wap, except for the 500 Gy dose treatment which was transferred to 5 wap seedlings. This is because at the age of 2 wap mutant seedlings dose 500 Gy on average only 1 cm high, the plant is still weak, if transplanted planting it will quickly wither and easily die if there are many factors that interfere with the growth and development of seedlings in the field, such as being carried away by water and conch pest attacks. At the age of 3 wap only 11 seedlings remain. This event is because the higher the dose used causes higher physiological damage that occurs and can cause death (Fadli et al., 2018).

Table 2. Advanced Test Results of Plant Height, Flowering Age, Number of Productive Tillers, Total Number of Tillers

Character	Treatment	Average		Min Value	Max Value	Range
PH	G16	138.5	ef	126.2	148.7	22.5
	Inpago U-1	128.2	cd	114.0	139.5	25.5
	IPB 3S	146.3	g	137.9	158.4	20.5
	D200	133.5	e	117.0	151.5	34.5
	D300	125.2	c	90.3	182.4	92.1
	D400	116.7	b	11.6	139.5	127.9
AF	D500	88.4	a	24.6	138.5	113.9
	Treatment	Average		Min Value	Max Value	Range
	G16	69.3	c	69.0	70.0	1.0
	Inpago U-1	59.3	a	58.0	60.0	2.0
	IPB 3S	62.3	b	62.0	63.0	1.0
	D200	73.1	d	67.0	82.0	15.0
SPT	D300	81.7	ef	59.0	95.0	36.0
	D400	81.297	e	72.0	95.0	23.0
	D500	88.857	g	83.0	94.0	11.0
	Treatment	Average		Min Value	Max Value	Range
	G16	18.97	b	11.0	30.0	19.0
	Inpago U-1	26.00	c	13.0	46.0	33.0
STT	IPB 3S	11.10	a	6.0	15.0	9.0
	D200	16.75	tn	6.0	30.0	24.0
	D300	18.89	tn	6.0	34.0	28.0
	D400	17.69	tn	3.0	41.0	38.0
	D500	23.57	tn	4.0	46.0	42.0
	Treatment	Average		Min Value	Max Value	Range
G16	23.43	b	11.0	24.0	13.0	
Inpago U-1	17.97	a	12.0	36.0	24.0	
IPB 3S	10.37	c	6.0	15.0	9.0	

Character	Treatment	Average		Min Value	Max Value	Range
	D200	16.11	tn	6.0	28.0	22.0
	D300	18.36	tn	6.0	34.0	28.0
	D400	17.08	tn	4.0	41.0	37.0
	D500	22.86	tn	4.0	46.0	42.0

Description: PH = Plant Height, AF = Age Flowering, SPT = Sum of Productive Tiller, STT= Sum of Total Tiller, D200 = irradiation dose 200 Gy, D300 = irradiation dose 300 Gy, D400 = irradiation dose 400 Gy, D500 = irradiation dose 500 Gy, and tn = unreal

Table 3. Test Results of Harvest Age, Panicle Length, Sum of Filled Grain per Panicle, and Sum of Empty Grain per Panicle

Character	Treatment	Average		Min Value	Max Value	Range
HA	G16	101.33	c	101.00	102.00	1.00
	Inpago U-1	93.67	a	93.00	94.00	4.00
	IPB 3S	97.33	b	97.00	98.00	1.00
	D200	107.28	d	102.00	115.00	13.00
	D300	116.15	ef	94.00	130.00	36.00
	D400	115.88	ef	107.00	130.00	23.00
PL	D500	123.86	g	118.00	129.00	11.00
	Treatment	Average		Min Value	Max Value	Range
	G16	25.21	b	22.13	26.80	4.67
	Inpago U-1	22.61	def	15.57	26.97	11.40
	IPB 3S	30.72	a	26.57	33.03	6.47
	D200	24.87	bc	20.97	28.03	7.07
SFGP	D300	24.37	bcd	18.33	33.63	15.30
	D400	23.46	bcde	14.63	30.37	15.73
	D500	17.92	g	11.87	26.57	14.70
	Treatment	Average		Min Value	Max Value	Range
	G16	128.59	b	100.67	159.67	59.00
	Inpago U-1	93.61	c	55.33	156.00	100.67
SEGP	IPB 3S	229.18	a	147.67	300.33	152.67
	D200	71.79	d	2.33	137.00	134.67
	D300	29.41	e	0.00	139.33	139.33
	D400	2.46	f	0.00	25.33	25.33
	D500	0.00	f	0.00	0.00	0.00
	Treatment	Average		Min Value	Max Value	Range
G16	27.63	ab	14.33	88.00	73.67	
Inpago U-1	19.11	a	5.33	36.00	30.67	
IPB 3S	82.60	cde	48.67	143.00	94.33	
D200	64.26	c	12.67	158.67	146.00	
D300	96.93	def	21.00	170.67	149.67	
D400	110.05	f	24.00	172.67	148.67	
D500	71.05	cd	26.00	155.67	129.67	

Description: HA = Harvest Age, PL = Panicle Length, SFGP = Sum of Filled Grain Per Panicle; SEGP = Sum of Empty Grain Per Panicle, D200 = irradiation dose of 200 Gy, D300 = irradiation dose of 300 Gy, D400 = irradiation dose of 400 Gy, D500 = irradiation dose of 500 Gy, and tn = unreal

Gamma radiation treatment causes the plants to be shorter than their parents (Table 2). The higher the dose given, the shorter the plant. The shortest plant is in mutant plants with a dose of 500 Gy, which is 88.43. The result is supported by the opinion of Astuti et al. (2019), an increase in the dose of irradiation can cause an increase in plant sensitivity so that there is a decrease in plant growth. This is due to reduced amounts of *endogenous* growth hormone in plants. This statement is

supported by the opinion of Kupchishin et al. (2019), that high doses of irradiation can cause the destruction of chemical bonds of a plant compound so that the death of meristematic cells occurs in the area of the plant growing point, characterized by the absence of plant height increase every week and even causing death. According to Oh et al. (2018) and Kuzmić (2018), the inhibition of cell metabolism or cell damage or death occurs due to the presence of RNA synthesis disorders, causing

inhibition of the synthesis of enzymes needed in the growth process, such as enzymes that stimulate budding. The presence of disturbances in the structure of DNA can cause the resulting enzymes to lose their function.

Plants that are not too tall are expected in plant breeding activities. The ideal height of rice plants according to Kuzmanović et al. (2021) is 90 cm to 100 cm. Rice plants that are too high are easy to fall over. Falling down will reduce rice yields (Sadimantara et al., 2018). Meanwhile, rice plants that are too short will make it difficult to harvest. Dhaka et al. (2021), further confirmed that plant height is correlated with the yield degree of rice plants. This makes the plant height character is one of the characters that determines farmers' acceptance of new superior varieties.

Flowering age (Table 2) and harvest age (Table 3) increased after gamma irradiation. The increase in flowering age and harvest age is in line with the increasing dose of irradiation given. Successively, the flowering age and the longest harvest age occurred in mutant plants with doses of 500 Gy, namely 88.86 and 123.86.

This is suspected because the irradiation treatment can damage plant cells, resulting in disruption of plant growth including flowering age and harvest age. Mutations may cause changes in DNA and chromosomes that result in changes in the duration of flowering and harvest duration. Furthermore, research by Purwanto et al. (2019) produced rice mutants with a longer flowering and harvesting age compared to their parents.

The results showed that gamma-ray irradiation had no effect on the total number of tillers and the number of productive tillers (Table 2). This is due to situation of diploic selection in mutations, as happened with the irradiation of grapevine clones (Vondras et al., 2019). Diploic selection is a condition where mutant cells are inferior to other cells around them, so that in the next development the cells will return to normal. On the other hand, if the cells affected by mutants are able to disrupt normal cells, then in the next generation the plant will grow into a generation of mutants.

On the other hand, gamma-ray irradiation is random so gene mutations are also random (Mardiyah et al., 2021). Some characters undergo changes but other characters may not change. It is suspected that the genes controlling the total tillers number and the productive tillers number did not mutate, so that the radiation treatment had no effect on the two characters.

The length of the panicle of mutant plants has decreased compared to the parent plants (G16). The 500 Gy dose mutant population has shortest panicle of 17.92 cm (Table 3). Panicle gets shorter as the dose of irradiation increases. Research by Arinta et al. (2018) proved that there was no correlation between the length

of the panicles and the number of grains per panicle due to the different shape and arrangement of the panicles. The different panicle shapes between cultivars can be influenced by genetic factors.

The variable amount of grain per panicle shows the higher the amount of filled grain in the panicle, the higher the productivity of the crop, and conversely the higher the amount of empty grain, the lower the productivity. The amount of filled grain per panicle in irradiated plants has decreased compared to parent. The 500 Gy mutant plant has the lowest average amount of filled grain, which is 0.00. The results showed that the higher the dose of irradiation given, the lower the amount of filled grain per panicle. This statement is in accordance with the results of research by (Mardiyah et al. (2022) showed a decrease in the filled grain amount at doses of 200 Gy to 400 Gy compared to control. There is an increase in the amount of empty grain in irradiated plants compared to parent plants. The 400 Gy mutant plant has the highest amount of empty grain of per panicle which is 110.05 grains.

These results indicate that the irradiation dose not only causes a higher number of empty grain, but also a lower number of panicles and panicle length. Radiation treatment caused a decrease in the reproductive capacity of plants and increased the amount of empty grain. According to Rahayu et al. (2020), the decrease in the number of fertile grain is caused by one of the chromosomal aberrations. Yunus et al. (2018), added that panicle length can affect the productivity of rice plants, both the number of filled grain and the number of empty grain. The number of panicles that are too large can reduce the filling to one panicle for another, thus the limited amount of nutrients will increase the number of empty panicles. Tumanggor et al. (2022), added that the presence of gamma-ray radiation treatment causes sterilization of panicles. The higher the irradiation dose given, the higher the occurrence of damage and even cell death that causes rice flowers to become sterile.

The death of somatic cell populations due to irradiation can occur directly or indirectly. According to Kumar et al. (2021), death can directly occur due to the degradation of enzymes that play a role in the IAA biosynthesis process as well as an increase in DNA and chromosomal damage which is directly proportional to the increase in the dose of irradiation given.

According to Susila et al. (2019), death indirectly occurs due to toxic influences from the results of water radiolysis in the form of free radicals H_2O_2 and $OH\cdot$. Water is the material that undergoes the most irradiation, then decomposes into H_2O^+ and e^- . The reaction further forms free radicals which then combine with peroxides. When free radicals and peroxides react with other molecules, compounds will be formed that can affect the plant biological system.

The results showed that gamma-ray mutation treatment resulted in a decrease in the weight of 100 grains mutant plants compared to parents (Table 4). Parent plant (G16) has an average weight of the highest 100 grains (3.04 g) while the mutant plant 500 Gy has an average weight value of the lowest 100 grains (0.00 g). This shows that the higher the dose given, the higher

decreases by 100 grains plants. The same results were shown by Karera's (2019) research that the treatment of an irradiated dose of 400 Gy caused a very noticeable difference in losing weight of 100 butir. According to Karera (2019), this happens because higher doses result in greater damage in inhibiting generative character in plants.

Table 4. Weight of 100 Grains, Weight of Filled Grain Per Clump, Weight of Empty Grain Per Clump

Character	Treatment	Average		Min Value	Max Value	Range
WG	G16	3.04	a	2.59	3.37	0.78
	Inpago U-1	2.72	bcd	2.25	3.09	0.84
	IPB 3S	2.74	bc	1.62	3.00	1.38
	D200	2.82	b	2.00	3.31	1.31
	D300	2.15	e	0.00	3.80	3.80
	D400	1.16	f	0.00	3.17	3.17
	D500	0.00	g	0.00	0.00	0.00
WFGC	Treatment	Average		Min Value	Max Value	Range
	G16	36.43	ab	13.21	71.77	58.56
	Inpago U-1	33.47	bc	12.56	62.80	50.24
	IPB 3S	43.94	a	21.36	88.20	66.84
	D200	14.42	d	0.30	47.03	46.73
	D300	7.33	de	0.00	71.58	71.58
	D400	0.33	e	0.00	2.28	2.28
WEGC	D500	0.00	e	0.00	0.00	0.00
	Treatment	Average		Min Value	Max Value	Range
	G16	3.67	ab	1.47	5.81	4.34
	Inpago U-1	3.09	a	1.10	5.72	4.62
	IPB 3S	4.91	c	2.91	9.15	6.24
	D200	6.03	d	0.43	12.15	11.72
	D300	7.49	e	2.00	17.12	15.12
WEGC	D400	6.78	de	0.22	16.84	16.62
	D500	6.53	de	0.16	19.27	19.11

Description: WG = Weight of 100 Grains, WFGC = Weight of Filled Grain per Clump, WEGC = Weight of Empty Grain per Clump, D200 = irradiation dose of 200 Gy, D300 = irradiation dose of 300 Gy, D400 = irradiation dose of 400 Gy, D500 = irradiation dose of 500 Gy, and tn = unreal.

The weight of grain per clump indicates how much grain is produced in one clump and also one panicle of both the main panicle and the branch of the panicle. Based on Table 4, the weight of grain filled with clumps has decreased in plants that have been irradiated when compared to parent plants. Parent plants have an average of 36.43 g, while mutant plants of 500 Gy have the lowest average filled grain weight of per clump, which is 0.00 g.

The weight character of the empty grain per clump shows an increase along with an increase in the dose of irradiation given. The occurrence of diversity weight of grain per clump indicates a mutation in plants so that there is an increase in plant diversity. Wu et al. (2019), succeeded in proving that irradiation causes high genetic diversity based on the Jaccard difference coefficient of 0.337 to 1.000.

The decrease in yield and yield support characteristics was caused by exposure to gamma rays on the seeds. Damage to genes, chromosomes and cells

increased with increasing doses of irradiation. Hong et al. (2022), explained that the higher the dose of irradiation, the more free radicals are formed. Excess ROS due to gamma irradiation treatment cannot be removed due to decreased antioxidant enzyme activity. This will inhibit plant growth. Furthermore, Gudkov et al. (2019) and Duarte et al. (2023) added that high doses of irradiation would inhibit plant physiological processes. The results of research by Wu et al. (2019) on chrysanthemum 'Pinkling' showed that frequency of chromosomal aberrations increased according to the irradiation dose.

Based on the explanation that has been described above, it clearly shows that rice plants that have previously been given gamma-ray radiation treatment at various dose levels show a diverse character compared to comparison plants and abnormal growth compared to parent plants. The more mutated genes, the higher the potential for the formation of new gene combinations, so that this gene combination will increase the diversity in the population.

Mutations cannot be observed in the M1 generation. The presence of mutations can be determined in the M2 generation and beyond. This is due to physiological damage after the seeds are irradiated with gamma rays. The damage is as a result of the formation of free radicals that are very labile in the reaction process resulting in mutational changes at the DNA, cell, or tissue level (Hong et al., 2022). However, this damage is not derived. This is due to diplontic selection towards recovery or improvement of the function of the enzyme system which is disturbed due to gamma ray irradiation. Physiological damage occurs only in M1, while gene mutations and chromosomal mutations will be passed down in later generations.

Variety of Genotypes and Varieties of Phenotypes

Gamma-ray irradiation treatment causes the emergence of diversity in each treatment based on range values (the value of the difference between the

maximum and minimum values). The highest diversity in plant height character was found in the 400 Gy irradiation dose treatment (127.9), the total number of tillers variable was found in the 500 Gy dose treatment (42), the variable number of productive tillers was found in the 500 Gy dose treatment (42), the panicle length variable was found in the 400 Gy dose treatment (15.73), the filled grain amount variable was found in the 300 Gy dose treatment (139.33) but lower when compared to the IPB 3S comparison plant, the variable amount of empty grain was found in the 300 Gy dose treatment (149.67), the filled grain weight variable was found in the 300 Gy dose treatment (71.58), the empty grain weight variable was found in the 500 Gy dose treatment (19.11), the 100 grain weight variable was found in the 300 Gy dose treatment (3.80), the flowering age variable was in the 300 Gy treatment (36.00), and the harvest age variable was in the 300 Gy dose treatment (36.00) (Table 2-4).

Table 5. Variety of Genotypes and Varieties of Phenotypes

Response Variables	σ^2g	σ^2p	σ_{σ^2g}	σ_{σ^2p}	Ket.	
					σ^2g	σ^2p
PH (cm)	379.61	389.59	19.48	19.74	L	L
AF (dap)	127.01	128.01	11.27	11.31	L	L
STT (tillers)	1.99	16.33	1.41	4.04	S	L
SPT (tillers)	6.46	11.35	2.54	3.37	L	L
PL (cm)	9.53	11.08	3.09	3.33	L	L
SFGP (grain)	1064.07	1156.68	32.62	34.01	L	L
SEGP (grain)	329.61	599.37	18.16	24.48	L	L
HA (dap)	122.08	122.63	11.05	11.07	L	L
WG (g)	1.50	1.52	1.23	1.23	S	S
WFGC (g)	29.83	63.37	5.46	7.96	L	L
WEGC (g)	0.17	0.56	0.42	0.75	S	S

Description: σ^2g = genetic variety, σ^2p = phenotypic variety, σ_{σ^2g} = standard deviation of genetic variety, σ_{σ^2p} = standard deviation of phenotypic variety, L = broad diversity, S = narrow diversity, dap = days after planting

The highest genetic variety value is found in the character of the amount of filled grain per panicle, while the lowest genetic variety value was found in the character of the weight of empty grain per clump. Based on the comparison between the variety value and the 2sd value, it can be seen that the genetic variety value on the character of plant height, flowering age, number of productive tillers, panicle length, number of filled grain per panicle, number of empty grain per panicle, harvest age, and weight of filled grain per clump have a value greater than 2sd so that these characters fall into the criteria of broad genetic diversity (Table 5). This means that these criteria have a chance at genetic improvement. The result is supported by the opinion of Widyapangesthi et al. (2022) that characters that have genetic diversity with broad categories occur because genetic factors are the ones that have a great influence on the visual appearance of a plant when compared to environmental factors. The more diverse the plant characters in the population, the higher the desired gene frequency. Furthermore, Widyapangesthi et al. (2022)

emphasized that broad genetic diversity increases the chances of successful selection getting higher. ³⁴

The total number of tillering characters, the weight of 100 grains, and the weight of the empty grain per clump have a narrow genetic diversity meaning that the characters have no chance of improvement. This is supported by the opinion of Wahyuni et al. (2019), if a trait has a broad genetic diversity then selection can be carried out on the plant population. Conversely, if a trait has narrow genetic diversity, then selection activities cannot be carried out because individuals in that population are relatively uniform.

Heritability

³⁴ The character of plant height, flowering age, number of productive tillers, panicle length, number of filled grain per panicle, number of empty grain, harvest age, and weight of 100 grains have a relatively high heritability value (Table 6). High heritability indicates that the genetic influence on phenotypes is greater than that of environmental influences (Wahyuni et al., 2019).

This is supported by the statement of Taneva et al. (2019), that the high heritability indicates the large number of additive genes that contribute to the trait so that it can be inherited in later generations.

Table 6. Heritability

Character	Heritability Value	Information
PH (cm)	0.97	T
FA (hst)	0.99	T
STT (tillers)	0.12	R
SPT (tillers)	0.57	T
PL (cm)	0.86	T
SFGP (grain)	0.92	T
SEGP (grain)	0.55	T
HA (hst)	1.00	T
WG (g)	0.99	T
WFGC (g)	0.47	S
WEGC (g)	0.31	S

Description: T = high heritability, S = imedium tability, and R = low heritability

40

Conclusion

Based on the results of the study, the following conclusions can be drawn: (1). The 400 Gy dose treatment caused high diversity in plant height variable (127.9), filled grain amount (139.33), grain content weight (71.58), weight of 100 grains (3.80), flowering age (36.00), and harvest age (36.00). The 400 Gy dose treatment caused diversity in the panicle length variable (15.73). The 500 Gy dose treatment caused diversity in the variables of total tiller number (42.00), number of productive tillers (42.00), and empty grain weight (19.11); (2). Selection for the improvement of mutant plant character can be made on the character of plant height, flowering age, number of productive tillers, panicle length, number of filled grain per panicle, number of empty grain per panicle, and harvest age.

Acknowledgments

The researcher expressed his gratitude to the Ministry of Education and Technology for the funds provided through the PNPB research of Mataram University in Capacity Building Research.

Author Contributions

Conceptualization, methodology, Ni Wayan Sri Suliartini; software, Shinta Adekayanti; validation, Ni Wayan Sri Suliartini, and Anak Agung Ketut Sudharmawan; formal analysis, Shinta Adekayanti; investigation, I Gusti Putu Muliarta Aryana; resources, Ni Wayan Sri Suliartini, and Shinta Adekayanti; data curation, I Gusti Putu Muliarta Aryana; writing—original draft preparation, Ni Wayan Sri Suliartini, and Shinta Adekayanti; writing—review and editing, Ni Wayan Sri Suliartini; supervision, Ni Wayan Sri Suliartini, Anak Agung Ketut Sudharmawan, and I Gusti Putu Muliarta Aryana; project administration, Ni Wayan Sri Suliartini; funding acquisition, Ni Wayan Sri Suliartini. All

authors have read and agreed to the published version of the manuscript.

Funding

This research was funded by the PNPB research of Mataram University in Capacity Building Research.

Conflicts of Interest

No Conflicts of interest.

References

- Allard, R. W. (1960). *Principles of Plant Breeding*. John Wiley and Sons, Inc.
- Arinta, K., & Lubis, I. (2018). Pertumbuhan dan produksi beberapa kultivar padi lokal Kalimantan. *Bul. Agrohorti*, 6(2), 270–280. <https://doi.org/10.29244/agrob.6.2>
- Astuti, D., Sulistyowati, Y., & Nugroho, S. (2019). Uji radiosensitivitas sinar gamma untuk menginduksi keragaman genetik sorgum berkadar lignin tinggi. *Jurnal Ilmiah Aplikasi Isotop Dan Radiasi*, 15(1), 1–6. <https://doi.org/10.17146/jair.2019.15.1.5480>
- Ayun, Q., Kurniawan, S., & Saputro, W. A. (2020). Perkembangan konversi lahan pertanian di bagian negara agraris. *VIGOR: Jurnal Ilmu Pertanian, Pika Dan Subtropika*, 5(2), 38–44. Retrieved from <https://jurnal.untidar.ac.id/index.php/vigor/article/view/3040>
- BPS.go.id. (2022). *Produksi Padi Tahun 2021 Turun 0,43 persen (Angka Tetap) Badan Pusat Statistik*. Retrieved from <https://www.bps.go.id/pressrelease/2022/03/01/1909/produksi-padi-tahun-2021-turun-0-43-persen-angka-tetap.html>
- Castañeda-Ovando, A., Pacheco-Hernández, M. L., Páez-Hernández, M. E., Rodríguez, J. A., & Galán-Vidal, C. A. (2019). Chemical studies of anthocyanins: A review. *Food Chemistry*, 113(4), 859–871. <https://doi.org/10.1016/j.foodchem.2008.09.001>
- Dhaka, K., Yadaw, R. B., Baral, B. R., Pokhrel, K. R., & Rasaily, S. (2021). Agro-morphological and genotypic diversity among rice germplasms under rainfed lowland condition. *AGRIVITA Journal of Agricultural Science*, 43(3), 466–478. <https://doi.org/10.17503/agrivita.v43i3.3026>
- Duarte, G. T., Volkova, P. Y., Fiengo Perez, F., & Horemans, N. (2023). Chronic Ionizing Radiation of Plants: An Evolutionary Factor from Direct Damage to Non-Target Effects. *Plants*, 12(5), 1178. <https://doi.org/10.3390/plants12051178>
- Dwiatmini, K., & Afza, H. (2018). Anthocyanin Content Characterization on Pigmented Local Rice as Genetic Resources of Functional Food. *Buletin Plasma Nutrafah*, 24(2), 125. <https://doi.org/10.21082/blpn.v24n2.2018.p125->

- 3 134
Fadli, N., Syarif, Z., Satria, B., & Akhir, N. (2018). The Effect of Gamma Cobalt-60 Ray Irradiation on Cultivar Growth in Taro White (*Xanthosoma Sagittifolium* L.). *International Journal of Environment Agriculture and Biotechnology*, 3(6), 2020-2025. <https://doi.org/10.22161/ijeab/3.6.9>
- 12 Gudkov, S. V., Grinberg, M. A., Sukhov, V., & Vodeneev, V. (2019). Effect of ionizing radiation on physiological and molecular processes in plants. *Journal of Environmental Radioactivity*, 202, 8-24. <https://doi.org/10.1016/j.jenvrad.2019.02.001>
- 2 Hong, M. J., Kim, D. Y., Jo, Y. D., Choi, H.-I., Ahn, J.-W., Kwon, S.-J., Kim, S. H., Seo, Y. W., & Kim, J.-B. (2022). Biological Effect of Gamma Rays According to Exposure Time on Germination and Plant Growth in Wheat. *Appl. Sci*, 12(3208), 1-14. <https://doi.org/10.3390/app12063208>
- Kumar, G., & Mishra, M. (2021). Mutagenic potential of gamma rays on somatic cell division and morphological parameters in *Foeniculum vulgare* Mill. *Current Botany*, 12, 66-71. <https://doi.org/10.25081/cb.2021.v12.6608>
- 7 Kupchishin, A. I., Taipova, B. G., Lisitsyn, V. M., & Niyazov, M. N. (2019). Study of the influence of the electron irradiation dose on the deformation of mylar films taking into account the processes of destruction and crosslinking. *IOP Conference Series: Materials Science and Engineering*, 510, 012025. <https://doi.org/10.1088/1757-899X/510/1/012025>
- 20 Kuzmanović, L., Giovenali, G., Ruggeri, R., Rossini, F., & Celoni, C. (2021). Small "nested" introgressions from wild *Thinopyrum* species, conferring effective resistance to fusarium diseases, positively impact durum wheat yield potential. *Plants*, 10(3), 1-16. <https://doi.org/10.3390/plants10030579>
- 43 Kuzmić, M. (2018). *Role of protein and DNA damage in biological response to radiation and aging*. Dissertation. Laboratoire de recherche sur les effets des radionucléides sur les écosystèmes. University Of Split, School of Medicine, Split.
- Mardiyah, A., Marnita, Y., & Syahril, M. (2021). Keragaan dan Produksi Padi Gogo Lokal Aceh Kultivar Gogo Generasi M1 Hasil Iradiasi Sinar Gamma. *Jurnal Ilmiah Aplikasi Isotop Dan Radiasi*, 17(1), 11-16. Retrieved from <https://jurnal.batan.go.id/index.php/jair/article/view/6231>
- Mardiyah, A., Wandira, A., & Syahril, M. (2022). Variabilitas Dan Heritabilitas Populasi Padi Gogo Kultivar Aarias Kuning Generasi Mutan-1 Hasil Iradiasi Sinar Gamma. *Jurnal Inovasi Penelitian*, 3(2), 4827-4838. Retrieved from <https://stpmataram.ejournal.id/JIP/article/view/1746>
- Ningrat, M. A., Mual, C. D., & Makabori, Y. Y. (2021). Pertumbuhan dan Hasil Tanaman Padi (*Oryza sativa* L.) pada Berbagai Sistem Tanam di Kampung Desay, Distrik Prafi, Kabupaten Manokwari. In *Prosiding Seminar Nasional Pembangunan dan Pendidikan Vokasi Pertanian* (Vol. 6, Issue 1, pp. 325-332). <https://doi.org/10.47687/snppvp.v2i1.191>
- 30 Oh, Y. J., Kwak, M. S., & Sung, M. H. (2018). Protection of radiation-induced DNA damage by functional cosmetic poly-gamma-glutamate. *Journal of Microbiology and Biotechnology*, 28(4), 527-533. <https://doi.org/10.4014/jmb.1712.12016>
- Parlaongan, A., Supriyanto, & Wulandari, A. S. (2022). Effects of Gamma Ray Irradiation to Induce Genetic Variability of Teak Planlets (*Tectona grandis* Linn. F.). *Journal of Sylva Indonesiana*, 5(01), 10-21. <https://doi.org/10.32734/jsi.v5i01.6166>
- 10 Purwanto, E., Nandariyah, Yuwono, S. S., & Yunindanova, M. B. (2019). Induced mutation for genetic improvement in black rice using gamma-ray. *Agrivita*, 41(2), 213-220. <https://doi.org/10.17503/agrivita.v41i2.876>
- Rahayu, S., Destavany, V., & Dasumiati. (2020). Keragaan Malai Mutan Padi Generasi M1 Hasil Iradiasi Gamma. *Aplikasi Isotop Dan Radiasi*, 16(2), 59-66. <https://doi.org/10.17146/jair.2020.16.2.5763>
- 14 Riviello-Flores, M. L., Cadena-Iñiguez, J., Ruiz-Posadas, L. M., Arévalo-Galarza, M. L., Castillo-Juárez, I., Hernández, M. S., & Castillo-Martínez, C. R. (2022). Use of gamma radiation for the genetic improvement of underutilized plant varieties. *Plants (Basel)*, 11(9), 1161. <https://doi.org/10.3390/plants11091161>
- 8 Sadimantara, G. R., Nuraida, W., Suliartini, N. W. S., & Muhidin. (2018). Evaluation of some new plant type of upland rice (*Oryza sativa* L.) lines derived from cross breeding for the growth and yield characteristics. *IOP Conf. Series: Earth and Environment Science*, 157, 12048. <https://doi.org/10.1088/1757-1315/157/1/012048>
- Stansfield, W. D. (1991). *Genetika* (M. Affandi & L. T. H. Erlangga (trans.)). Erlangga.
- Sudharmawan, A. A. K., Aryana, I. G. M., Suliartini, N. W. S., & Purnama, S. A. (2022). Genetic diversity of red rice (*Oryza sativa* L.) population M2 results of G16 rice genotype mutations with 200gy and 300gy gamma ray irradiation. *Jurnal Biologi Tropis*, 22(4), 1340-1346. <https://doi.org/10.29303/jbt.v22i4.4425>
- Suherman, S., Syukur, M., Husni, A., & Hartati, R. S. (2021). Increased genetic variability of sugarcane through gamma ray irradiation. *IOP Conf. Series: Earth and Environmental Science*, 653, 12134. <https://doi.org/10.1088/1757->

- 27 1315/653/1/012134
Suliartini, N. W. S., Aryana, I. G. P. M., 27 Itharmawan, A. A. K., & Wangiyana, W. (2021). Yield Potential Improvement of Some Lowland Red Rice Lines in F4 Pedigree Selection. *Russian Journal of Agricultural and Socio-Economic Sciences*, 120(12), 179-184. <https://doi.org/10.18551/rjoas.2021-12.18>
- Suliartini, N. W. S., Aryana, I. G. P. M., Wangiyana, I. W., 16 Awit, I. K., Muhidin, & Rakian, T. C. (2020). Identification of upland red rice mutant lines (RYZA SATIVA L.) high yield potential. *International Journal of Scientific and Technology Research*, 9(3), 4690-4692. Retrieved from <https://www.ijstr.org/final-print/mar2020/Identification-Of-Upland-Red-Rice-Mutant-Lines-oryza-Sativa-L-High-Yield-Potential.pdf>
- 31 Suliartini, N. W. S., Wijayanto, T., Madiki, A., Boer, D., Muhidin, & Juniawan. (2018). Relationship of some upland rice genotype after gamma irradiation. *IOP Conf. Series: Earth and Environmental Science*, 122, 12033. <https://doi.org/10.1088/1755-1315/122/1/012033>
- 18 Susila, E., Susilowati, A., & Yunus, A. (2019). The morphological diversity of Chrysanthemum resulted from gamma ray irradiation. *BIODIVERSITAS*, 20(2), 463-467. <https://doi.org/10.13057/biodiv/d200223>
- Syahril, M. (2018). Rancangan bersekut (augmented design) untuk penelitian bidang pemuliaan padi. *Jurnal Penelitian Agrosamudra*, 5(1), 63-66. Retrieved from <https://jurnal.unsam.ac.id/index.php/jagrs/article/view/848>
- 19 Taneva, K., Bozhanova, V., & Petrova, I. (2019). Variability, heritability and genetic advance of some grain quality traits and grain yield in durum wheat genotypes. *Bulgarian Journal of Agricultural Science*, 25(2), 288-295. Retrieved from <https://www.agrojournal.org/25/02-10.pdf>
- 21 Tang, L., Risalat, H., Cao, R., Hu, Q., Pan, X., Hu, Y., & Zhang, G. (2022). Food Security in China: A Brief View of Rice Production in Recent 20 Years. *Foods*, 11(21), 3324. <https://doi.org/10.3390/foods11213324>
- Tumanggor, G. E., Iswahyudi, & Mardiyah, A. (2022). Pertumbuhan, produksi dan karakter genetik padi kultivar Silesa generasi M-2 hasil iradiasi sinar gamma. *Jurnal Penelitian Agrosamudra*, 9(2), 31-40. Retrieved from <https://ejournalunsam.id/index.php/jagrs/article/view/6519/3786>
- Umam, R., Sudharmawan, & Sumarjan. (2018). Tampilan Sifat Kuantitatif Beberapa Galur F7 Padi Beras Merah (*Oryza sativa* L.) Hasil Silang Ganda Indica Dengan Javanica. *Crop Agro*, 11(1), 40-47. Retrieved from <https://cropagro.unram.ac.id/index.php/caj/article/view/188/157>
- 1 Vondras, A. M., Minio, A., Blanco-Ulate, B., Figueroa-Balderas, R., Penn, M. A., Zhou, Y., Seymour, D., Ye, Z., Liang, D., Espinoza, L. K., Anderson, M. M., Walker, M. A., Gaut, B., & Cantu, D. (2019). The genomic diversification of grapevine clones. *BMC Genomics*, 20(1), 972. <https://doi.org/10.1186/s12864-019-6211-2>
- 11 Wahyuni, H., Swasti, E., & Yusniwati, Y. (2019). Genetic Diversity Of Age, Plant Height And Number Of Grain Per Panicle Characters Of F3 Generation Derived From Crossing Silopuk With Fatmawati Varieties. *Jurnal Indonesian Journal of Crop Science*, 1(2), 36-46. <https://doi.org/10.25077/jijcs.1.2.36-46.2019>
- Wahyuni, S., Siregar, H. M., Isnaini, Y., Widiarsih, S., & Dwimahyani, I. (2022). Keragaman Morfologi Hibrid *Begonia sagaensis* Wiriad x *Begonia galeolopsis* Ardi & D.C. Rhomas Hasil Iradiasi Sinar Gamma. *Buletin Kebun Raya*, 25(1), 22-33. <https://doi.org/10.55981/bkr.2022.743>
- Widyapangesthi, D. A., Moeljani, I. R., & Soedjarwo, D. P. (2022). Keragaman Genetik Dan Heritabilitas M1 Mentimun (*Cucumis sativus* L.) Lokal Madura Hasil Iradiasi Sinar Gamma 60CO. *Jurnal Agrium*, 19(2), 191. <https://doi.org/10.29103/agrium.v19i2.7841>
- 26 Wu, J. H., Zhang, J., Lan, F., Fan, W. F., & Li, W. (2019). Morphological, cytological, and molecular variations induced by gamma rays in ground-grown chrysanthemum 'Pinkling.' *Can. J. Plant Sci*, 100, 68-77. <https://doi.org/10.1139/cjps-2019-0064>
- 13 Wulandari, Y. A., Sobir, & Aisyah, S. I. (2019). Studi radiosensitivitas dan analisis keragaman M1 kacang tunggak (*Vigna unguiculata* L) hasil induksi mutasi. *Jurnal Agrosains Dan Teknologi*, 4(1), 1-9. <https://doi.org/10.24853/jat.4.1.1-9>
- Yunus, A., Parjanto, N., & Wulandari, S. (2018). Performance of Mentik Wangi rice (*Oryza sativa* L.) M2 generation from gamma ray irradiation. *IOP Conf. Ser.: Earth Environ. Sci*, 142, 12050. <https://doi.org/10.1088/1755-1315/142/1/012050>
- 4 Zafar, S. A., Aslam, M., Albaqami, M., Ashraf, A., Hassan, A., Iqbal, J., Maqbool, A., Naeem, M., Al-Yahyai, R., & Zuan, A. T. K. (2022). Gamma rays induced genetic variability in tomato (*Solanum lycopersicum* L.) germplasm. *Audi Journal of Biological Sciences*, 29(5), 3300-3307. <https://doi.org/10.1016/j.sjbs.2022.02.008>

Effect of Mutations Induction on Vegetative and Generative Characters of G16 Rice (*Oryza sativa* L.)

ORIGINALITY REPORT

23%

SIMILARITY INDEX

20%

INTERNET SOURCES

17%

PUBLICATIONS

10%

STUDENT PAPERS

PRIMARY SOURCES

- 1 Massimo Gardiman, Mirko De Rosso, Fabiola De Marchi, Riccardo Flamini. "Metabolomic profiling of different clones of *vitis vinifera* L. cv. "Glera" and "Glera lunga" grapes by high-resolution mass spectrometry", *Metabolomics*, 2023
Publication 1%
- 2 Mohd Rafi Wani, Bharati Patel. "Photosynthetic performance of *Terminalia arjuna* Roxb towards different doses of ^{60}Co γ -irradiation", *Biocatalysis and Agricultural Biotechnology*, 2022
Publication 1%
- 3 jurnal.ikipmataram.ac.id
Internet Source 1%
- 4 arastirma.tarimorman.gov.tr
Internet Source 1%
- 5 dspace.lib.ntua.gr
Internet Source 1%

6	jurnal.uisu.ac.id Internet Source	1 %
7	project.abaiuniversity.edu.kz Internet Source	1 %
8	www.pathofscience.org Internet Source	1 %
9	www.biotech-asia.org Internet Source	1 %
10	S H Y Saragih, K D Sitanggang, W Lestari, K Rizal. "Growth Response of Labuhanbatu Local Rice With Gamma Ray Irradiation in M1 Generation", IOP Conference Series: Earth and Environmental Science, 2022 Publication	1 %
11	repository.uin-malang.ac.id Internet Source	1 %
12	Hui Liu, Huaizhu Li, Guiyan Yang, Guoyun Yuan, Yanping Ma, Ting Zhang. "Mechanism of early germination inhibition of fresh walnuts (<i>Juglans regia</i>) with gamma radiation uncovered by transcriptomic profiling of embryos during storage", Postharvest Biology and Technology, 2021 Publication	1 %
13	sinta3.ristekdikti.go.id Internet Source	<1 %

14	Milan Stanković. "10th Anniversary of Plants—Recent Advances and Further Perspectives", Plants, 2023 Publication	<1 %
15	researchportal.sckcen.be Internet Source	<1 %
16	www.ijteee.org Internet Source	<1 %
17	talenta.usu.ac.id Internet Source	<1 %
18	v2.smujo.id Internet Source	<1 %
19	ir.haramaya.edu.et Internet Source	<1 %
20	Submitted to University College London Student Paper	<1 %
21	Submitted to Wageningen University Student Paper	<1 %
22	www.updatepublishing.com Internet Source	<1 %
23	Submitted to Australian National University Student Paper	<1 %
24	www.ejplantbreeding.org Internet Source	<1 %

25	conference.um.ac.id Internet Source	<1 %
26	medicine.dp.ua Internet Source	<1 %
27	opac.wsb.torun.pl Internet Source	<1 %
28	agrosainstek.ubb.ac.id Internet Source	<1 %
29	repository.upi.edu Internet Source	<1 %
30	Ji-Young Hong, Sung-Ho Son, Sang-Pil Hong, Sung-Hun Yi, Sun Hee Kang, Na-Kyoung Lee, Hyun-Dong Paik. "Production of β -glucan, glutathione, and glutathione derivatives by probiotic <i>Saccharomyces cerevisiae</i> isolated from cucumber jangajji", LWT, 2018 Publication	<1 %
31	Muhidin, E Syam'un, Kaimuddin, Y Musa, G R Sadimantara, S Leomo, D N Yusuf, T C Rakian. "The bacterial mixture effect of <i>Azotobacter</i> and <i>Azospirillum</i> on nitrogen content and harvest date of upland red rice cultivar", IOP Conference Series: Earth and Environmental Science, 2021 Publication	<1 %
32	hal-normandie-univ.archives-ouvertes.fr	

Internet Source

<1 %

33

Submitted to Politeknik Negeri Lampung

Student Paper

<1 %

34

Gusmiatun Gusmiatun, Neni Marlina. "Peran Pupuk Organik dalam Mengurangi Pupuk Anorganik pada Budidaya Padi Gogo",
Agrikan: Jurnal Agribisnis Perikanan, 2019

Publication

<1 %

35

Submitted to Syiah Kuala University

Student Paper

<1 %

36

www.scopus.com

Internet Source

<1 %

37

Submitted to University of Calgary

Student Paper

<1 %

38

ejurnalunsam.id

Internet Source

<1 %

39

scholars.direct

Internet Source

<1 %

40

www.ajol.info

Internet Source

<1 %

41

www.irsn.fr

Internet Source

<1 %

42

kasetartjournal.ku.ac.th

Internet Source

<1 %

43

theses.hal.science

Internet Source

<1 %

44

N Rahman, Sriharti, C E W Anggara, H M Astro et al. "The effectiveness of beyonic startmix organic biological fertilizer on growth and yield of Inpari rice 30", IOP Conference Series: Earth and Environmental Science, 2023

Publication

<1 %

45

academic-accelerator.com

Internet Source

<1 %

46

journal.ummat.ac.id

Internet Source

<1 %

47

pkht.ipb.ac.id

Internet Source

<1 %

48

Darmawan, Kazutake Kyuma, Arsil Saleh, H. Subagjo, Tsugiyuki Masunaga, Toshiyuki Wakatsuki. "Effect of green revolution technology during the period 1970–2003 on sawah soil properties in Java, Indonesia: II. Changes in the chemical properties of soils", Soil Science and Plant Nutrition, 2006

Publication

<1 %

49

N Kasim, R Sjahril, M Riadi, F Arbie. " The Phenotype performance of M3 red rice mutant (L.) ", IOP Conference Series: Earth and Environmental Science, 2018

Publication

<1 %

50

Submitted to Universitas Jenderal Soedirman

Student Paper

<1 %

51

eprints.unm.ac.id

Internet Source

<1 %

52

ijeab.com

Internet Source

<1 %

53

Wei-Wei Yang, Xi-Long Chen, Marc Saudreau, Xiao-Yun Zhang, Man-Rang Zhang, Hang-Kong Liu, Evelyne Costes, Ming-Yu Han. "Canopy structure and light interception partitioning among shoots estimated from virtual trees: comparison between apple cultivars grown on different interstocks on the Chinese Loess Plateau", *Trees*, 2016

Publication

<1 %

54

Yusi Nurmalita Andarini, Willy Bayuardi Suwarno, Hajrial Aswidinnoor, Hakim Kurniawan. "Genetic relationship of pigmented rice (*Oryza sativa* L.) collected from Eastern Indonesia based on morpho-agronomical traits and SSR markers", AIP Publishing, 2022

Publication

<1 %

55

Z.A. Abdulhamed, N.M. Abood, A.H. Noaman. "Genetic Path Analysis and Correlation Studies of Yield and Its Components of Some

<1 %

56

repository.radenfatah.ac.id
Internet Source

<1 %

57

Muthukumaravel Chinnusamy, Brahma Dutta Kaushik, Radha Prasanna. "Growth, Nutritional, and Yield Parameters of Wetland Rice as Influenced by Microbial Consortia Under Controlled Conditions", Journal of Plant Nutrition, 2007
Publication

<1 %

58

Ni Wayan Sri Suliartini, Kuswanto Kuswanto, Nur Basuki, Andy Soegianto. "Superior Lines Candidates Evaluation of Two Local Red Rice Southeast Sulawesi cultivars (Indonesia) Derived from Gamma rays Irradiation Techniques", International Journal of Plant Biology, 2017
Publication

<1 %

59

ejurnal.budiutomomalang.ac.id
Internet Source

<1 %

60

journal.ibrahimy.ac.id
Internet Source

<1 %

61

turcomat.org
Internet Source

<1 %

62

Internet Source

<1 %

63

www.ejurnal.litbang.pertanian.go.id

Internet Source

<1 %

64

A Khaeruni, T Wijayanto, M H Musa, M Rahayu, A Rahman, V N Satrah. "

Effectiveness of the composition of bokashi organic materials on the ability of biofresh biological agents in inducing resistance of three maize varieties to sheath blight disease () ", IOP Conference Series: Earth and Environmental Science, 2020

Publication

<1 %

65

Grazielle Ribeiro Borges, Mostafa Galal Aboelkheir, Fernando Gomes de Souza Junior, Kassia Cristina Waldhelm et al. "Poly (butylene succinate) and derivative copolymer filled with Dendranthema grandiflora biolarvicide extract", Environmental Science and Pollution Research, 2020

Publication

<1 %

66

I M Sudantha, Sudirman, N M L Ernawati. "The effect of method and dosage application of biofungicide extract of Legundi leaf fermented with Trichoderma harzianum fungus for control of Fusarium wilt disease on shallots", IOP Conference Series: Earth and Environmental Science, 2021

<1 %

67

Kartika Yurlisa. "Review: Komposisi Vegetasi dan Keragaman Gulma di Lahan Padi Sawah", Prosiding Seminar Nasional Pembangunan dan Pendidikan Vokasi Pertanian, 2021

Publication

<1 %

68

Nurussanatil Alfaini, Nurussakinah Daulay. "The Effect of Self-Efficacy and Well-Being on Smartphone Addiction", EDUKATIF : JURNAL ILMU PENDIDIKAN, 2022

Publication

<1 %

69

Submitted to University of Hertfordshire

Student Paper

<1 %

70

fp.univamedan.ac.id

Internet Source

<1 %

71

ojs.umb-bungo.ac.id

Internet Source

<1 %

72

www.ornamentalhorticulture.com.br

Internet Source

<1 %

73

E Panjaitan, N S Napitupulu, K Afrianta. "The effect of organic fertilizer and cobgn vmpost of volcanic ash on growth and yield of red chili (Capsicum annum L.)", IOP Conference Series: Earth and Environmental Science, 2018

Publication

<1 %

74

I G M Kusnarta, D Rahmadhanti, N W D Dulur, W Wangiyana. "Soil chemical characteristics and yield of red rice under aerobic irrigation system as affected by intercropping with peanut and application of organic wastes on permanent raised-beds", IOP Conference Series: Earth and Environmental Science, 2021

Publication

<1 %

75

Submitted to Nexford Learning Solutions

Student Paper

<1 %

76

Sutardi, Yayan Apriyana, Popi Rejekiningrum, Annisa Dhienar Alifia et al. "The Transformation of Rice Crop Technology in Indonesia: Innovation and Sustainable Food Security", Agronomy, 2022

Publication

<1 %

77

W Nasira, R Adawiyah, Muhidin, G R Sadimantara, D N Yusuf. "Effect of liquid organic fertilizer derived from moringa on growth of upland red rice lines crosses from SE Sulawesi", IOP Conference Series: Earth and Environmental Science, 2021

Publication

<1 %

78

cgspace.cgiar.org

Internet Source

<1 %

79

faculty.nuist.edu.cn

Internet Source

<1 %

80	gjournals.org Internet Source	<1 %
81	jurnal.stmikroyal.ac.id Internet Source	<1 %
82	jurnal.umberau.ac.id Internet Source	<1 %
83	koreascience.kr Internet Source	<1 %
84	repositorio.utp.edu.co Internet Source	<1 %
85	repository.unibos.ac.id Internet Source	<1 %
86	www.gjournals.org Internet Source	<1 %
87	Dyah Marganingrum, Mochamad Furqon Azis Ismail, Diana Rahayuning Wulan. "Assessment of shallow groundwater contamination on Pari Island, Indonesia", Environmental Monitoring and Assessment, 2022 Publication	<1 %
88	Tresta Nurina Ciptaningayu, Arie Dipareza Syafei, Muhayatun Santoso, Joni Hermana, Rachmat Boedisantoso, Abdu Fadli Assomadi. "Identification of Metal Pollutants in Ambient	<1 %

Air(Case Study: West Surabaya, East Java, Indonesia)", Research Square, 2020

Publication

89

W Wangiyana, I G P M Aryana, N W D Dular. "Effects of mycorrhiza biofertilizer on anthocyanin contents and yield of various red rice genotypes under aerobic irrigation systems", Journal of Physics: Conference Series, 2021

Publication

<1 %

90

Yirui Li, Li Chen, Xiaodie Zhan, Liang Liu, Feihong Feng, Zihua Guo, Dan Wang, Hao Chen. " Biological effects of gamma-ray radiation on tulip (L.) ", PeerJ, 2022

Publication

<1 %

91

R P Bachtari, S Listyawati, Sutarno. " Starch, amylose and amylopectin levels of M5 and M6 generations of black rice irradiated by gamma Co ray ", Journal of Physics: Conference Series, 2020

Publication

<1 %

Exclude quotes Off

Exclude matches Off

Exclude bibliography Off