

## About Legume Research

*Legume Research* is a **journal** covering the technologies/fields/categories related to **Agronomy and Crop Science (Q2); Plant Science (Q2); Soil Science (Q2)**. It is published by **Agricultural Research Communication Centre**. The overall rank of Legume Research is **12698**. According to **SCImago Journal Rank (SJR)**, this journal is ranked **0.391**. SCImago Journal Rank is an indicator, which measures the scientific influence of journals. It considers the number of citations received by a journal and the importance of the journals from where these citations come. SJR acts as an alternative to the Journal Impact Factor (or an average number of citations received in last 2 years). This journal has an **h-index** of **18**. The **best quartile** for this journal is **Q2**.

The ISSN of **Legume Research** journal is **02505371, 09760571**. An International Standard Serial Number (ISSN) is a unique code of 8 digits. It is used for the recognition of journals, newspapers, periodicals, and magazines in all kind of forms, be it print-media or electronic. **Legume Research** is cited by a total of **471** articles during the last 3 years (Preceding 2022).

### Indexing Services :


BIOSIS Preview, ISI Citation Index, Biological Abstracts, **Elsevier (Scopus and Embase)**, AGRICOLA, Google Scholar, CrossRef, CAB Abstracting Journals, Chemical Abstracts, Indian Science Abstracts, EBSCO Indexing Service.

**Note:** The **impact score** or **impact index** shown here is equivalent to the average number of times documents published in a journal/conference in the past two years have been cited in the current year (i.e., Cites / Doc. (2 years)). It is based on Scopus data and can be a little higher or different compared to the impact factor (IF) produced by Journal Citation Report. Please refer to the **Web of Science data source** to check the exact journal impact factor <sup>TM</sup> (Thomson Reuters) metric.

## Important Metrics

Title	Legume Research
Abbreviation	Legume Res.
Publication Type	Journal
Subject Area, Categories, Scope	Agronomy and Crop Science (Q2); Plant Science (Q2); Soil Science (Q2)
h-index	18
Overall Rank/Ranking	12698
SCImago Journal Rank (SJR)	0.391
Impact Score	0.84
Publisher	Agricultural Research Communication Centre
Country	India
ISSN	02505371, 09760571
Best Quartile	Q2
Coverage History	2008-2022

Year	SJR
2023/2024	Coming Soon
2022	0.391
2021	0.349
2020	0.397
2019	0.371
2018	0.343
2017	0.319
2016	0.197
2015	0.154
2014	0.218



## Table of contents

### Genetics and Breeding

[Genetic and biochemical properties of \*Cicer\* spp reveal distinction between wild and cultivated chickpea genotypes](#)

Feyza Nur Kafadar, Ali Özkan, Canan Can, Yagmur Kar, Dürdane Mart, Ercan Ceyhan

[Heterosis and combining ability analysis for yield and yield contributing traits in velvet bean \*Mucuna pruriens\* \(L.\) DC](#)

A. Chinapolaiah, K. Hima Bindu, G.N. Manjesh, N. Hariprasad Rao, S. Sudheer Kumar, T. Vasantha Kumar

[Exploring the genetic diversity and population structure of Turkish common bean germplasm by the iPBS-retrotransposons markers](#)

Muhammet Fatih Aydin, Faheem Shehzad Baloch

[Gamma rays induced urdbbean \[\*Vigna mungo\* \(L.\) Hepper\] mutants with YMV resistance, good batter quality and bold seeded type](#)

C. Vanniarajan, S. Ganeshram, J. Souframanien, K. Veni, S. Anandhi Lavanya, V. Kuralarasan

[Molecular dissection of genetic diversity in pigeonpea \[\*Cajanus cajan\* \(L.\) Millsp.\] minicore collection](#)

S. Muniswamy, R. Lokesha, Rachit Saxena, B. Fakrudin, Kishan Patel, Yamanura

[Physico-chemical, engineering and functional properties of two soybean cultivars](#)

Jagbir Rehal, Vinod Beniwal, B.S. Gill

[Essential selection criteria for dual uses of red pea \(\*Lathyrus cicera\* L.\)](#)

A. Adak, H. Canci, N. Ertoy Inci, F. Oncu Ceylan, D. Sari, H. Sari, T. Yildirim, C. Toker

[Genetic studies in mungbean \[\*Vigna radiata\* \(L.\) Wilczek\]](#)

P. Khajudparn, O. Poolsawat, P.A. Tantasawat

[Successful Interspecific hybridization between mungbean \[\*Vigna radiata\* \(L.\) Wilczek\] and ricebean \[\*V. umbellata\* \(Thunb.\) Ohwi & Ohashi\]](#)

Lestari Ujianto, Nur Basuki, Kuswanto, Astanto Kasno

### Plant Physiology

[Effect of water stress on morpho-physiological parameters of fenugreek \(\*Trigonella foenum-graecum\* L.\) genotypes](#)

S.N. Saxena, R.K. Kakani, L.K. Sharma, D. Agarwal, S. John, Y. Sharma

[Variations in osmo-regulatory and anti-oxidative status and their links with total nitrogen and phosphorus concentrations in lucerne during re-growth](#)

Zhennan Wang, Mei Yang, Jiaoyun Lu, Huimin Yang

[The effects of plant growth promoting rhizobacteria on antioxidant activity in chickpea \(\*Cicer arietinum\* L.\) under salt stress](#)

Hilal Yilmaz, Haluk Kulaz

### Agronomy

[Role of irrigation and mulch in chickpea \(\*Cicer arietinum\* L.\) growth, productivity and moisture extraction pattern in Alluvial Zone of West Bengal, India](#)

S. Sarkar, A. Sarkar

[Foliar applications for amelioration of iron deficiency in peanut \(\*Arachis hypogaea\* L.\)](#)

*Shamim Akhtar, Nazneen Bangash, Armghan Shahzad, Sammer Fatima, Muhammad Akbar, Muhammad Sajjad Iqbal, Muhammad Arshad, Fayyaz-Ul Hassan*

[Improving productivity of groundnut \(\*Arachis hypogaea\* L.\) under drip and micro sprinkler fertigation system](#)

*Jeetendra Kumar Soni, N. Asoka Raja, Vimal Kumar*

[Insight into floral biology and ancillary characteristics of underutilized legume-Bambara groundnut \[\*Vigna subterranea\* \(L.\)Verdc.\]](#)

*Kailash Chandra, R. Nandini, R. Gobu, Pranesh, Chitti Bharat Kumar, R. Muthuraju*

[Productivity and profitability of legume-cereal forages under different planting ratio and nitrogen fertilization](#)

*Akanksha Tamta, Rakesh Kumar, Hardev Ram, R.K. Meena, V.K. Meena, M.R. Yadav, D.J. Subrahmanya*

[Response of soil arthropods population to different farming systems under cotton integrated with groundnut and pigeonpea crops](#)

*R. Chethan, R.K. Patil, B. Halappa*

[Performance of rajma \(\*Phaseolus vulgaris\*\) cultivars under organic mulches in Meghalayan Plateau of North Eastern India](#)

*Y. Marwein, Lala I.P. Ray*

## Editorial Board

### Chief Editor

J. S. Sandhu  
Vice-Chancellor  
Sri Karan Narendra Agri. Uni., Jobner, Rajasthan, INDIA

### Associate Chief Editor

**Kadambot Siddique,**  
Chair in Agriculture and Director,  
The UWA Institute of Agriculture,  
Australia

### Editors

**H.A. Shimelis**  
Deputy Director  
African Center for Crop Improvement

**Pei Xu**  
Professor  
China Jiliang University

**Bao Lam Huynh**  
Assistant Professional Researcher  
University of California

### Managing Editor

**R.D. Goel,**  
A.R.C.C.,  
Karnal

### Members

**Shiv Kumar Agrawal,** Syria  
**Paras Nath,** Fiji Islands  
**Harvir Singh,** Hyderabad  
**Uma Sah,** Kanpur  
**K.B. Saxena,** Patancheru  
**K. Thiyagarajan,** Aduthurai  
**S. Jebaraj,** Madurai  
**Om Vir Singh,** Jodhpur  
**G. S. Sidhu,** U.S.A.  
**Z. Abraham,** Thrissure  
**B. B. Gupta,** Jammu

**Vilas A. Tonapi,** New Delhi

**Suk-Ha Lee,** Korea

**S.K. Dubey,** New Delhi  
**Murari Singh,** Syria  
**A.K. Bera,** Mohanpur  
**E.V.D. Sastry,** Jobner  
**G.R. Korwar,** Hyderabad  
**V.K. Khanna,** Pantnagar  
**K.R. Sreeramalu,** Bangalore  
**C.R. Ananda Kumar,** Madurai  
**A. Vijay Kumar,** Coimbatore  
**B.R. Ranwah,** Udaipur  
**H. Kumar,** Varanasi  
**Mustafa YILMAZ,** TÜRKİYE  
**Henry T. Nguyen,** Columbia

Please write to :

**The Managing Editor**  
*Agricultural Research  
Communication Centre  
1130, Sadar, Karnal - 132 001  
E mail : [arccgoel@yahoo.co.in](mailto:arccgoel@yahoo.co.in)  
Phone No : +91-184-2255080*

**Re: Interspecific hybridization**  
**ARCC Journals <contact@arccjournals.com>**

Kam, 31 Mei 2018,  
16.35

Dear Dr. Ujianto,

Thank you for sending the article, it is with Editor for first opinion to check for its quality to process further.

Further it is to inform you that there are processing and printing charges of US\$ 450 for the article which author need to remit at the time of approval. This is for your information and confirmation.

Please feel free to contact for any other assistance.

-----  
Best Regards

Gaurav Gupta

General Managing Editor

Agricultural Research Communication Centre |

1130, Sadar Bazar | Karnal - 132001 | Haryana (INDIA) |

Web: [www.arccjournals.com](http://www.arccjournals.com) | E-mail: [gaurav@arccjournals.com](mailto:gaurav@arccjournals.com) / [contact@arccjournals.com](mailto:contact@arccjournals.com)

On Mon, May 21, 2018 at 7:30 PM, <[contact@arccjournals.com](mailto:contact@arccjournals.com)> wrote:

<b>Journal</b>	Legume Research
<b>Title</b>	Successful Interspecific hybridization between mungbean ( <i>Vigna radiata</i> L. Wilczek) and ricebean ( <i>V. umbellata</i> (Thunb.) Ohwi & Ohashi)
<b>Article type</b>	Research Article
<b>Area of research</b>	Plant Breeding
<b>Country</b>	Indonesia
<b>Subject</b>	Interspecific hybridization
<b>Message</b>	Please let me know if there is something to be improved
<b>No of author</b>	4
<b>Corresponding author</b>	Lestari Ujianto
<b>Corresponding author affiliation</b>	1Faculty of Agriculture University of Mataram
<b>Corresponding author email</b>	<a href="mailto:ujianto@unram.ac.id">ujianto@unram.ac.id</a>
<b>Corresponding author contact</b>	+6281805756649
<b>Corresponding author address</b>	1Faculty of Agriculture University of Mataram, Jl. Majapahit 62 Mataram, 83121, NTB, Indonesia

**Acknowledgement of Article "LR-433" "Successful Interspecific hybridization between mungbean (Vigna radiata L. Wilczek) and ricebean (V. umbellata (Thunb.) Ohwi & Ohashi)"**

**ARCC JOURNALS** <contact@arccjournals.com>

5 Jun 2018,  
16.29

**Dear Dr. Lestari Ujianto,**

Greetings!!

Thank you for submitting the article. It is to acknowledge the receipt of the article for "Legume Research" entitled "Successful Interspecific hybridization between mungbean (Vigna radiata L. Wilczek) and ricebean (V. umbellata (Thunb.) Ohwi & Ohashi)". For any correspondence, please mention Reference Number **LR-433** for this article in your all email.

Submit the attached certificate that this article or its data has not been/will not be sent to any other journal for publication. Also, inform the subject area of specialization of the article to send it the right area of expert for Review Process. Printing charges will be informed with the approval of the article. Acknowledge the receipt of this email and submit the attached certificate duly filled with for taking further action. Should you require any further clarification, feel free to contact. Check the article status any time at <http://arccjournals.com/>.

Warm Regards

Gaurav Gupta

General Managing Editor

Agricultural Research Communication Centre |

1130, Sadar Bazar | Karnal - 132001 | Haryana (INDIA) |

Tel: +91 (0) 9034233744 |

Web: [www.arccjournals.com](http://www.arccjournals.com) | E-mail: [gaurav@arccjournals.com](mailto:gaurav@arccjournals.com) /[contact@arccjournals.com](mailto:contact@arccjournals.com)

**lestari ujianto** <ujianto@unram.ac.id>

5 Jun 2018,  
22.00

Dear Sir/Madam, we are happy to submit article submission certificate (attached file). The subject area of specialization of the article (LR-433) is plant breeding. Thank you very much.

Best regard,

Dr. Ujianto

**ARCC Journals** <contact@arccjournals.com>

Jum, 8 Jun 2018,  
17.45

Dear Dr. Ujianto,

Thank you for sending the certificate, soon article will be processed further for review.

-----  
Best Regards

Gaurav Gupta

General Managing Editor

Agricultural Research Communication Centre |

1130, Sadar Bazar | Karnal - 132001 | Haryana (INDIA) |

Web: [www.arccjournals.com](http://www.arccjournals.com) | E-mail: [gaurav@arccjournals.com](mailto:gaurav@arccjournals.com) /[contact@arccjournals.com](mailto:contact@arccjournals.com)

**Comments of Reviewer on your Article "LR-433" "Successful Interspecific hybridization between mungbean (*Vigna radiata* L. Wilczek) and ricebean (*V. umbellata* (Thunb.) Ohwi & Ohashi)"**  
ARCC JOURNALS <contact@arccjournals.com>

Min, 19 Agu 2018,  
19.10

**Dear Dr. Lestari Ujianto,**

It is with reference to your article no. ARCC/LR-433 entitled "**Successful Interspecific hybridization between mungbean (*Vigna radiata* L. Wilczek) and ricebean (*V. umbellata* (Thunb.) Ohwi & Ohashi)**" which you have sent for publication in "**Legume Research**", kindly find the attachment of comments by reviewer on your article, you are requested to revise the article as per comments and follow guidelines to author and sample paper attached on the website [www.arccjournals.com](http://www.arccjournals.com). Submit the corrected complete article including all author names and addresses for further process. You are requested to share complete details of all the authors in the article with email id

**Kindly give point wise reply of the reviewer comments in Word File.**

"It is requested to you to please consider reading relevant studies and reviews published in ARCC journals and insert at least 3 to 5 references from any of the ARCC Journals if found suitable. You can find complete open access archive of our previous issues at <http://arccjournals.com/archive>"

Revise the article and references according to the Format of the journal. Click here for Sample Paper:-  
<http://arccjournals.com/images/Sample%20Article.pdf>

Guidelines for Authors:-

<http://arccjournals.com/author-guidelines>

1. Full-length articles (not exceeding 3,000 words including references but excluding Tables and Figures).
2. Short theoretical articles (not exceeding 2,000 words including references).
3. Review Articles (not exceeding 6000 words)

Note:- No further authors will be added after receiving the revised article.

Resubmit your revised article as the reply of this email.

Regards

Editor

Agricultural Research Communication

1130, Sadar Bazar | Karnal

- 132001 | Haryana

Centre |  
(INDIA) |

Web: [www.arccjournals.com](http://www.arccjournals.com) | E-mail: [editor@arccjournals.com](mailto:editor@arccjournals.com) / [contact@arccjournals.com](mailto:contact@arccjournals.com)

**lestari ujianto** <ujianto@unram.ac.id>

Jum, 31 Agu 2018,  
23.52

Dear Sir / Madam, I have revised my article as per comments of reviewer and follow guide lines to author. I have added 3 references from any of the ARCC Journals. If there is something I have not completed, please let me know. Thank you very much.

Sincerely,

Lestari Ujianto



---

**Approval of article "LR-433" "Successful Interspecific hybridization between mungbean (*Vigna radiata* L. Wilczek) and ricebean (*V. umbellata* (Thunb.) Ohwi & Ohashi)"**

Kam, 13 Sep 2018,  
15.29

**ARCC JOURNALS** <contact@arccjournals.com>

**Dear Dr. Lestari Ujianto,**

We are glad to inform you that your article no. **LR-433** entitled "**Successful Interspecific hybridization between mungbean (*Vigna radiata* L. Wilczek) and ricebean (*V. umbellata* (Thunb.) Ohwi & Ohashi)**" has been recommended by the Editor and approved for publication as **Full Article** in "**Legume Research**".

We would like to thank you for choosing ARCC Journals to publish your article.

**"Please consider reading relevant studies and reviews published in ARCC group of journals and consider citing in your recent and future publications, if found suitable."**

Equally to increase more scope of ARCC Journals; please recommend our Journals to you Institute/university Library which will result in Global exposure of articles published in ARCC Journals. Subscription details are available at [www.arccjournals.com](http://www.arccjournals.com).

Kindly remit **USD 550 \$** as printing and processing charges for your article by Wire Bank Transfer, Western Union and PayPal. Invoice is enclosed.

[Click here](#) for Bank Details

[Click here](#) for PayPal

Western Union Details:-

Name: - **Akash Gupta**

Address: - Karnal, Haryana (INDIA)

Please feel free to contact for any further assistance you may require.

Best Regards

Gaurav Gupta

General Managing Editor

Agricultural Research Communication Centre |

1130, Sadar Bazar | Karnal - 132001 | Haryana (INDIA) |

Tel: +91 92555 40308 |

Web: [www.arccjournals.com](http://www.arccjournals.com) | E-mail: [gaurav@arccjournals.com](mailto:gaurav@arccjournals.com) /[contact@arccjournals.com](mailto:contact@arccjournals.com)

**lestari ujianto** <ujianto@unram.ac.id>

Rab, 19 Sep 2018,  
22.30

Dear Gaurav Gupta

General Managing Editor

Agricultural Research Communication Centre

Thank you very much for approving my article no. **LR-433** entitled "**Successful Interspecific hybridization between mungbean (*Vigna radiata* L. Wilczek) and ricebean (*V. umbellata* (Thunb.) Ohwi & Ohashi)**" for publication as **Full Article** in "**Legume Research**".

I have paid 550 Dollar for printing and processing charges for my article. Proof of payment is enclosed.

Sincerely,

Lestari Ujianto

**ARCC Journals** <contact@arccjournals.com>

20 Sep 2018,  
04.48

Thank you for the email, it normally takes 5 working days by the bank to reflect the charges in our account.

As soon we receive it, we will send acknowledgement to you with final acceptance letter.

-----  
Best Regards

Gaurav Gupta

General Managing Editor

Agricultural Research Communication Centre |

1130, Sadar Bazar | Karnal - 132001 | Haryana (INDIA) |

Web: [www.arccjournals.com](http://www.arccjournals.com) | E-mail: [gaurav@arccjournals.com](mailto:gaurav@arccjournals.com) / [contact@arccjournals.com](mailto:contact@arccjournals.com)

lestari ujianto <ujianto@unram.ac.id>

20 Sep 2018,  
06.58

Thank you very much

**Acknowledgement of Printing Fees and Acceptance Letter**

**ARCC JOURNALS** <contact@arccjournals.com>

Min, 23 Sep 2018,  
04.23



# Successful Interspecific hybridization between mungbean [*Vigna radiata* (L.) Wilczek] and ricebean [*V. umbellata* (Thunb.) Ohwi & Ohashi]

Lestari Ujianto\*<sup>1</sup>, Nur Basuki<sup>2</sup>, Kuswanto<sup>2</sup> and Astanto Kasno<sup>3</sup>

Faculty of Agriculture University of Mataram,  
Jl. Majapahit 62 Mataram, 83121, NTB, Indonesia.

Received: 05-06-2018

Accepted: 23-09-2018

DOI: 10.18805/LR-433

## ABSTRACT

Successful interspecific hybridization between several cultivars of mungbean (*Vigna radiata*) and ricebean (*V. umbellata*) was achieved using mungbean as the female parent. Hybridization was not achieved if ricebean was used as the female parent. Seeds of hybrids were viable and grew and developed normally, producing viable seeds. The gene action of all observed quantitative characters was partially dominant, implying that it should be possible to produce stable hybrids. The progeny had desirable characteristics of both parents suggesting that interspecific hybridization of the two species could be used to produce a new, superior cultivar.

**Key words:** Gene action, Hybrids, Interspecific hybridization, Mungbean, Ricebean.

## INTRODUCTION

Mungbean (*Vigna radiata*) is one of the main crops in Indonesia. It is an excellent protein source and easily digestible. It is an important short-duration legume crop with wide adaptability, requires low fertilizer inputs and has the ability to improve soil fertility by fixing nitrogen from the air (Arshad *et al.*, 2009, Fery, 2002, Turkova and Klozova, 1985). The main problems of mungbean cropping systems are low yields and susceptibility to many pests. The yield and resistance to pests and diseases can be improved through plant breeding programs, but mungbean has a low genetic background (Choudhary *et al.*, 2017, Das *et al.*, 2015, Somta *et al.*, 2007). According to Kumar *et al.* (20s15), Fasoula and Fasoula (2002), genetic variability is essential in plant breeding programs to achieve crop improvement. Ricebean (*V. umbellata*) has superior characters with respect to pod number per plant and resistance to some pests that are not found in mungbean.

This paper reports the interspecific hybridization between mungbean and ricebean and we report the production of hybrids with superior characters. This was achieved by enhancing the number of pods per plant using this genetic trait from ricebean (Ghafoor *et al.*, 2002, Gopinathan and Babu, 1985).

## MATERIALS AND METHODS

The hybridization work was conducted from May 2010 to August 2011 in pots set up in a shade house at the Agricultural Faculty, Mataram University. Field evaluation was conducted in irrigated lowland. Planting was conducted

for 3 weeks at weekly intervals for both the mungbean and ricebean cultivars to maximize the possibility of both species attaining flowering at the same time. The plants were prepared for hybridization by removing one third of the flowers and all of the anthers in the afternoon of the day before the flower was anticipated to flower. Pollination was conducted in the morning by placing pollen from the donor on the pistil of emasculated flowers. The interspecific hybridization between mungbean and ricebean to form the F1 and BC1.1 populations was conducted according to the IITA Research Guide 42 Hand Crossing of cowpea (Myers, 1996) except that pollination is done by hooking the stamen of ricebean to the stigma of the mungbean. Hybrids were evaluated using a randomized completely block design with four replications. The data were analyzed using analysis of variance, Duncan's Multiple Range Test, cross ability degree and estimation of gene action.

## RESULTS AND DISCUSSION

Interspecific hybridization between mungbean as female parent and ricebean as male parent was successfully carried out as assessed by the degree of cross ability (%) and change of the offspring characters. Eight F1 populations were obtained using four mungbean genotypes which have different characteristics and two genotypes of ricebean (Table 1). The highest cross ability degree was achieved between the Manyar variety of mungbean with the yellow seed ricebean genotype (65%). The lowest cross ability degree occurred between the Merak variety of mungbean and the yellow seed ricebean (44%).

\*Corresponding author's e-mail: [ujianto@unram.ac.id](mailto:ujianto@unram.ac.id)

<sup>1</sup>Faculty of Agriculture University of Mataram, Jl. Majapahit 62 Mataram, 83121, NTB, Indonesia.

<sup>2</sup>Faculty of Agriculture University of Brawijaya, Jl. Veteran, Malang, 65145, Indonesia.

<sup>3</sup>Research Institute for Nuts and Tubers, Jl. Raya Kendalpayak km 8, Malang, 65101 Indonesia.

**Table 1:** The degree of cross ability of hybridization between 4 mungbean genotypes and 2 ricebean genotypes.

Type of crossing	Number of crossed flowers	Harvested pods number	Degree of cross ability (%)
Manyar x RBK	240	156	65.0
Manyar x RBM	246	158	64.2
Merak x RBK	232	103	44.4
Merak x RBM	230	110	47.8
Sampeong x RBK	248	157	63.3
Sampeong x RBM	246	159	64.6
Vima x RBK	234	113	48.3
Vima x RBM	236	122	51.7

RBK- yellow seed ricebean, RBM- red seed ricebean.

Hybridization was successful with mungbean as the female receptor but was unsuccessful if the ricebean was used as the female receptor (Table 2). The incompatibility with ricebean as the female receptor could be because it lack a compound suitable for the germination of mungbean pollen (Thiyagu *et al.*, 2008 and Somta *et al.*, 2006). Another possibility could be the relative length of the stylus in mungbean and ricebean, where that of mungbean is shorter, being 1.6 mm compared with 1.4 for ricebean. Barathi *et al.* (2006) demonstrated that even though pollen can germinate, fertilization of the ovum may not be possible if the stylus is too long. Hence fertilization of the ova of ricebean may be difficult for this reason.

The different cross ability level of *Vigna* species has been reported by Bharathi *et al.* (2006) and Chen *et al.*

**Table 2:** The degree of cross ability of mungbean as female receptor and ricebean as female receptor.

Type of crossing	Degree of cross ability (%)	
	Mungbean as female	Ricebean as female
Manyar x RBK	63.9	0
Manyar x RBM	61.7	0
Merak x RBK	44.4	0
Merak x RBM	49.1	0
Sampeong x RBK	60.6	0
Sampeong x RBM	62.5	0
Vima x RBK	46.2	0
Vima x RBM	50.6	0

RBK- yellow seed ricebean, RBM- red seed ricebean.

**Table 3:** The degree of cross ability of backcross between hybrids and mungbean as female parent.

Type of crossing	Number of crossed flowers	Harvested pods number	Degree of cross ability (%)
(Manyar x RBK) x Manyar	192	118	61.5
(Manyar x RBM) x Manyar	191	113	59.2
(Merak x RBK) x Merak	188	72	38.3
(Merak x RBM) x Merak	183	74	40.4
Merak x (Merak x RBM)	190	86	45.3
(Sampeong x RBK)xSampeong	198	115	58.1
(Sampeong x RBM)xSampeong	193	120	62.2
(Vima X RBK) x Vima	184	85	46.2
(Vima X RBM) x Vima	189	87	46.0
Vima x (Vima x RBM)	188	83	44.2

Note : RBK = yellow seed ricebean, RBM = red seed ricebean.

(1977). Some researchers have successfully crossed mungbean and ricebean but they have reported a low level of cross compatibility (Chowdhury and Chowdhury, 1978, Ahn and Hartmann 1978, Thiyagu *et al.*, 2008). The relatively high cross compatibility that we report in Table 1 implies that the two species can be successfully hybridized, provided that the timing of anthesis and stigma receptivity is the same for both parents.

Backcrossing between hybrids and mungbean as female parent was also successful (Table 3). The cross ability degree of backcrossing was lower than for interspecific hybridization between mungbean and ricebean (Table 1). The highest degree of cross ability was achieved between hybrids and their female parent (62% for (Sampeong x RBM) x Sampeong) and 61.5% for (Manyar x RBK) x Manyar). The lowest degree of cross ability occurred with hybrids from crosses between the Merak variety of mungbean and yellow seed ricebean and the female parent (Merak x RBK) x Merak (38.3 %).

Hybridization is considered successful if the offspring characteristics are different from their parents, especially the female parent. The quantitative and qualitative characters of the hybrids were different from those of the parents (Table 4). This provides further evidence that interspecific hybridization between mungbean and ricebean has been successful. In this regard, both species have the same chromosome number ( $2n = 2x = 22$ ), an important factor in facilitating successful interspecies crossing (Gopinathan and Babu, 1985).

**Table 4:** The characteristics of parents and hybrids of interspecific hybridization between mungbean and ricebean.

Characteristics	Mungbean as female parent	Ricebean as male parent	Hybrids (F1)
Seed color	Green	Red	Brown
Plant height (cm)	58	106	83
Number of pod plant <sup>1</sup>	23	64	46
Maturity date (days)	62	86	68
Protein content (%)	27	18	29
Carbohydrate content (%)	16	54	28
Resistance to bruchus pest	Susceptible	Tolerant	Tolerant

The production of hybrids with characteristics that vary from their parents has also been reported between cowpea having short and stiff pods and longbean having long and soft pods producing offspring with characteristic combinations of both parents (Ujianto *et al.*, 2012). The research results of Chen *et al.* (1982) indicated that characteristics of interspecific hybridization among four species of vigna food legumes showed incorporation of their parent characteristics.

Tables 5 and 6 show that all offspring from the population crosses showed high seed germination ability (more than 80% required for certification standard). Seed germination of hybrids (F1) from crosses between the Manyar variety of mungbean and yellow seed ricebean (MR x RBK) or with red seed ricebean (MR x RBM) was not significantly different from that of their parents. Seed germination of hybrids from crosses between the Sampeong variety of mungbean with red seed ricebean (F1 S x RBM) was not significantly different from the male parent. Likewise, with backcross offspring between the first offspring with the female parent, seed germination was not significantly different.

Table 6 shows that the seed germination ability among female parents, Merak (MK) and Vima (V) varieties of mungbean, with red seed ricebean (RBM) as male parent is not significantly different either in the first or second generation. Seed germination in offspring from crosses of the Merak variety mungbean and red seed ricebean in F2 (MK x RBM) and the Vima variety mungbean with red seed ricebean in F2 (V x RBM) is higher than in the BC1 backcross except with V x (F1 (V x RBM)). Seed germination of offspring from all cross combinations was more than 90%. This suggests that the offspring from crosses seeds have high viability.

Table 7 shows that the potential ratio values were between 0.25 and 0.75. None of the values were less than 0.25, indicative of low dominance, nor in the range 0.75 – 1.25, typical of high dominance (Ullah *et al.*, 2010, Peter and Frey, 1966). Therefore the data in Table 7 imply partial dominance and that one parent has greater influence than the other parent. As there were progeny having character

**Table 5:** Seed germination of progeny population of result of crossing between mungbean (Manyar and Sampeong varieties) and ricebean (yellow and red seeds).

Genotype	Seed germination (%)	
	Mean	Standard error
Manyar (MR)	95.50 abcd	1.60
Sampeong (S)	89.17 a	1.59
Yellow seed ricebean (RBK)	94.17 bcde	1.60
Red seed ricebean (RBM)	95.56 cde	0.91
F1 (MR x RBK)	90.83 ab	1.59
F1 (MR x RBM)	93.33 abcde	1.93
F1 (S x RBK)	94.17 bcde	0.84
F1 (S x RBM)	92.50 abcd	0.83
BC1 (F1(MR x RBK) x MR)	94.44 bcde	0.64
BC1 (F1(MR x RBM) x MR)	94.44 bcde	0.64
BC1 (F1(S x RBK) x S)	91.11 abc	0.91
BC1 (F1(S x RBM) x S)	92.78 abcd	1.40
F2 (MR x RBK)	97.98 e	0.41
F2 (MR x RBM)	97.50 de	0.30
F2 (S x RBK)	96.90 de	0.46
F2 (S x RBM)	96.55 de	0.49

The numbers followed by the same letter in the same column indicates no significant difference based on Duncan's Multiple Range Test at the 5% significance level.

**Table 6:** Seed germination of progeny population of the result of crossing between mungbean (Merak and Vima varieties) and ricebean (red seed).

Genotype	Seed germination (%)	
	Mean	Standard error
Merak (MK)	95.56 ab	0.91
Vima (V)	97.22 ab	1.06
Red seed ricebean(RBM)	95.56 ab	0.91
F1 (MK x RBM)	95.00 ab	0.56
F1 (V x RBM)	96.11 ab	1.06
BC1 (F1(MK x RBM) x MK)	94.58 a	1.05
BC1 (MK x F1(MK x RBM))	94.58 a	1.25
BC1 (F1V x RBM) x V	94.58 a	1.42
BC1 V x (F1V x RBM)	95.00 ab	0.68
F2 (MK x RBM)	98.33 b	0.28
F2 (V x RBM)	98.33 b	0.38

The numbers followed by the same letter in the same column indicates no significant difference based on Duncan's Multiple Range Test at the 5% significance level.

**Table 7:** Gene action of some observed quantitative traits of some crossing types.

	Crossing types	Low parent	Parent mean	High parent	F1	Ratio
Plant	MR x RBK	68.60	99.15	129.70	75.60	0.385
Height (cm)	MR X RBM	68.60	92.90	117.20	73.90	0.391
	MK x RBM	62.40	87.70	113.00	51.00	0.725
	S x RBK	68.40	99.05	129.70	72.20	0.438
	S x RBM	68.40	92.80	117.20	78.70	0.289
	V x RBM	41.80	79.55	117.30	54.50	0.332
Maturity Date	MR x RBK	68.86	81.83	94.80	71.20	0.410
	MR X RBM	68.86	82.28	95.70	71.50	0.402
	MK x RBM	61.91	76.79	91.66	63.26	0.455
	S x RBK	70.56	82.68	94.80	74.50	0.337
	S x RBM	70.56	83.13	95.70	74.40	0.347
Number of pods Plant <sup>-1</sup>	V x RBM	59.21	75.44	91.66	64.14	0.348
	MR x RBK	18.70	36.95	55.20	21.40	0.426
	MR X RBM	18.70	37.15	55.60	20.70	0.446
	MK x RBM	15.61	34.15	52.69	17.40	0.452
	S x RBK	15.40	35.30	55.20	22.80	0.314
Seed Weight Plant <sup>-1</sup> (g)	S x RBM	15.40	35.50	55.60	22.80	0.316
	V x RBM	14.62	33.66	52.69	17.60	0.422
	MR x RBK	9.63	19.22	28.80	12.40	0.356
	MR X RBM	9.63	18.22	26.80	10.60	0.444
	MK x RBM	8.40	18.95	29.50	9.40	0.453
	S x RBK	7.90	18.35	28.80	10.80	0.361
	S x RBM	7.90	17.35	26.80	11.70	0.299
	V x RBM	7.60	18.55	29.50	10.60	0.363

The gene actions for all crosses were partly dominant.

combinations of both parents the production of a new superior variety should be possible.

## CONCLUSION

Hybridization was successful with mungbean as the female receptor but was unsuccessful if the ricebean was used as the female receptor. Backcrossing between hybrids and mungbean as female parent was also successful. The cross ability degree of backcrossing was lower than for interspecific hybridization between mungbean and ricebean.

## REFERENCES

- Ahn, C.S. and Hartmann, R.W. (1978). Interspecific hybridization between mungbean [*Vigna radiata* (L.) Wilczek]. and adzuki bean (*Vigna angularis* (Wild) Ohwi & Ohashi. *An American Society of Horticultural Science*, **103**:3-6.
- Arshad, M., Aslam, M., and Irshad, M. (2009). Genetic variability and character association among morphological traits of mungbean, *Vigna radiata* L. wilczek genotypes. *Agricultural Research*, **47**:121-126.
- Bharathi, A., Selvaraj, K.S.V., Veerabhadhiran, P., and Lakshmi, B.S. (2006). Crossability barriers in mungbean [*Vigna radiata* (L.) Wilczek]: with its wild relatives. *Indian Journal of Crop Science*, **1**:120-124.
- Chen, N.C., Parrot, J.F., Jacobb, T., Baker, L.R., and Carlson, P.S. (1977). Interspecific hybridization of food legumes by unconventional methods of plant breeding. Proceeding of first International Mungbean Symposium. Asian Vegetable Research and Development Center, Shanhun, Taiwan:247-252.
- Chen, N. C., Baker, L.R., and Honma, S. (1982). Interspecific crossability among four species of vigna food legumes. *Euphytica*, **32**:925-937.
- Choudhary, P., Payasi, S.K., and Patle, N.K. (2017). Genetic study and selection indices for grain yield of mungbean. *Legume Research*, **40**(5): 836-841.
- Chowdhury, R.K., and Chowdury, J.B. (1978). Intergeneric hybridization between *Vigna mungo* (L.) Hepper and *Phaseolus calcaratus* Roxb. *Indian Journal of Agricultural Science*, **48**:117-121.
- Das, S., Dewanjee, Maji, A., Sarkar, K.K. (2015). Breeding mungbean [*Vigna radiata* (L.) Wilczek] for development of high yielding protein rich lines with low trypsin inhibitor. *Legume Research*, **38**(4): 447-4451.
- Fasoula, V. A., and Fasoula, D.A. (2002). Principles underlying genetic improvement for high and stable crop yield potential. *Field Crop Research*, **75**:191-209.

All offspring from the population crosses showed high seed germination ability. The gene action of all observed quantitative characters was partially dominant.

## ACKNOWLEDGEMENT

This study was founded by Directorate of Higher Education, the Ministry of Education and Culture. Special thanks are dedicated to Dr. John W. Anderson, Pat Anderson, Prof. Taufik Fauzi, and Prof. Sunarpi for helping and supporting in the writing of this paper.

- Fery, F.L. (2002). New opportunities in *Vigna*. In: Trends in New Crops and New Uses, J. Janick and A. Whipkey (Eds.), ASHS Press, (Alexandria, VA.) 424–428.
- Ghafoor, A., Ahmad, Z., Qureshi, A.S., Bashir, M. (2002). Genetic relationship in *Vigna mungo* (L.) Hepper and *V. radiata* (L.) R.Wilczek based on morphological traits and SDS-PAGE. *Euphytica*, **123**:367–378.
- Gopinathan, M.C. and Babu, C.R (1985). Meiotic studies of the F1 hybrid between ricebean (*Vigna umbellata*) and its wild relative *V. minima*. *Genetica*, **71**:115-117.
- Kumar, B.K., Prakash, M., Gokulakrishnan, J. (2015). Genetic studies on biometric, biochemical, biophysical and morpho-physiological traits in mungbean [*Vigna radiata* (L.) Wilczek]. *Legume Research*, **38(4)**: 457-460.
- Myers, G.O. (1996). IITA Research Guide 42: Hand crossing of cowpea. IITA Ibadan Nigeria.
- Peter, F.C. and Frey, K.I., (1966). Genotype correlation, dominance and heritability of quantitative characters in oat. *Crop Science*, **6**:259-262.
- Somta, P., Ammaranan, C., Ooi, P.A.C., 424–428. Srinives, P. (2007). Inheritance of seed resistance to bruchids in cultivated mungbean [*Vigna radiata*, (L.) Wilczek]. *Euphytica*, **155**:47-55.
- Somta, P., Kaga, A., Tomooka, N., Kashiwaba, K., Isemura, T., Chaitieng, B., Srinives, P., and Vaughan, D.A. (2006). Development of an interspecific *Vigna* linkage map between *Vigna umbellata* (Thumb) Ohwi & Ohashi dan *Vigna nakashimae* (Ohwi) Ohwi & Ohasi and its use in analysis of bruchid resistance and comparative genomics. *Plant Breeding*, **125**:77-84.
- Thiyagu, K., Jayamani, P., and Nadarajan, N. (2008). Polen pistil interaction in interspecific crosses of *Vigna* sp. *Cytologia*, **73**:251-257.
- Turkova, V. and Klozova, E. (1985). Comparison in seed protein in some representatives of genus *Vigna*. *Biologia Plantarum*, **27**:70-73.
- Ujjianto, L., Idris, Yakop, U.M. (2012). Heritability and heterosis study on hybrids of crossing between cowpea and longbean. *Plasma Nutrafah Bulletin*, **18**:9-17.
- Ullah, S., Khan, A.S., Raza, A., and Sidiqie, A. (2010). Gene action analysis of yield related traits in spring wheat (*Triticum aestivum*). *International Journal of Agricultural Biology*, **12**:125-128.